

# Novel Conjugative Plasmid from *Escherichia coli* of Swine Origin That Coharbors the Multiresistance Gene *cfr* and the Extended-Spectrum- $\beta$ -Lactamase Gene *bla*<sub>CTX-M-14b</sub>

Wan-Jiang Zhang,<sup>a</sup> Xiu-Mei Wang,<sup>a</sup> Lei Dai,<sup>b</sup> Xin Hua,<sup>a</sup> Zhimin Dong,<sup>a</sup> Stefan Schwarz,<sup>c</sup> Siguo Liu<sup>a</sup>

State Key Laboratory of Veterinary Biotechnology, Harbin Veterinary Research Institute, Chinese Academy of Agricultural Sciences, Harbin, China<sup>a</sup>; Department of Veterinary Microbiology and Preventive Medicine, College of Veterinary Medicine, Iowa State University, Ames, Iowa, USA<sup>b</sup>; Institute of Farm Animal Genetics, Friedrich-Loeffler-Institut (FLI), Neustadt-Mariensee, Germany<sup>c</sup>

Two porcine *Escherichia coli* isolates harbored the *cfr* gene on conjugative plasmids of 38,405 bp (pGXEC6) and 41,646 bp (pGXEC3). In these two plasmids, the *cfr* gene was located within a 4,612-bp region containing a *tnpA*-IS26-*cfr*-IS26- $\Delta$ *hyp* element. Plasmid pGXEC3 was almost identical to pGXEC6 except for a 3,235-bp *ISEcp1*-*bla*<sub>CTX-M-14b</sub> insertion. The colocation of the multiresistance *cfr* gene with an extended-spectrum- $\beta$ -lactamase gene on a conjugative plasmid may support the dissemination of these genes by coselection.

During the past decade, the transferable multiresistance gene *cfr*, which confers resistance to phenicols, lincosamides, oxazolidinones, pleuromutilins, and streptogramin A (1), has been identified mainly in Gram-positive bacteria, including *Staphylococcus* spp., *Bacillus* spp., *Enterococcus* spp., *Macrococcus caseolyticus*, *Jeotgalicoccus pinnipedialis*, and *Streptococcus suis* (2, 3). However, a small number of reports have described the presence of the gene *cfr* in Gram-negative bacteria, such as *Proteus vulgaris* (4) and *Escherichia coli* (5–7). The global spread of this gene in bacteria of human and animal origin has gained particular public health attention, since *cfr* confers resistance to linezolid, the last-resort antibiotic against vancomycin-resistant enterococci, methicillin-resistant *Staphylococcus aureus*, and penicillin-resistant pneumococci (2, 8). In *Enterobacteriaceae* from human and animal sources, extended-spectrum- $\beta$ -lactamase (ESBL) genes have emerged during recent years globally (9). Since the year 2000, ESBL genes of the type *bla*<sub>CTX-M</sub> have become the predominant ESBL genes (10, 11). Similar to the *cfr* gene (2), ESBL genes are often located on plasmids, including plasmids that carry additional resistance genes which may facilitate the dissemination and persistence of the ESBL genes (8, 11–13).

So far, a single conjugative IncA/C broad-host-range plasmid of *E. coli*, designated pSCEC2 (6), has been described, in which the *cfr* gene was colocated with the resistance genes *tet*(A) (tetracycline resistance), *strA*-*strB* (streptomycin resistance), and *sul2* (sulfonamide resistance). In the present study, two novel *cfr*-carrying plasmids, pGXEC6 and pGXEC3, were identified, one of which also harbored a *bla*<sub>CTX-M-14b</sub> gene. These two plasmids were completely sequenced to gain insight into their genetic relatedness and understand how they may have developed.

Thirty-one pericardial fluid swab samples collected from diseased swine from June 2013 to January 2014 in four different commercial pig farms in the Guangxi Province, China, were inoculated on brain heart infusion agar plates supplemented with 10 mg/liter florfenicol. Two *cfr*-positive *E. coli* isolates, designated GXEC6 and GXEC3, were obtained from samples from two different pig farms, farm A (a breeding pig farm) and farm B (a fattening pig farm), respectively. The antimicrobial usage records for these two pig farms indicated that aside from ceftriaxone, which was used for prophylactic purposes at farm A, other anti-

microbial agents, such as ampicillin, florfenicol, sulfamethoxazole, amikacin, enrofloxacin, lincomycin, and tiamulin, were extensively used to prevent or treat bacterial infections at the two pig farms. The two *E. coli* isolates were typed by multilocus sequence typing (MLST) (see <http://mlst.warwick.ac.uk>) and subjected to pulsed-field gel electrophoresis (PFGE) (14). Conjugation experiments by filter mating and S1-PFGE combined with Southern blotting (15) were performed as described previously (15). PCR-based replicon typing (PBRT) followed previously published recommendations (16). The two *cfr*-positive original *E. coli* isolates and their transconjugants were tested for their susceptibility to amikacin, ampicillin, ceftazidime, chloramphenicol, enrofloxacin, florfenicol, gentamicin, sulfamethoxazole, and tetracycline by broth microdilution according to CLSI recommendations (17, 18). ESBL confirmatory tests also followed CLSI recommendations (18). The *cfr*-carrying plasmids of the transconjugants were sequenced using the 454 Life Sciences GS FLX system (Roche). The gaps between the contigs were closed by PCR, and the whole-plasmid sequences were draft assembled using the SeqMan software from DNAStar (Lasergene, Madison, WI, USA). The AatII, BstEII, ScaI, and SnaBI restriction profiles of *cfr*-carrying plasmids were determined and compared with the plasmid sequences to confirm the correct assembly of the contigs. The plasmid sequences were annotated using the Vector NTI program, and the putative coding sequences were identi-

Received 28 October 2014 Returned for modification 16 November 2014

Accepted 20 November 2014

Accepted manuscript posted online 24 November 2014

Citation Zhang W-J, Wang X-M, Dai L, Hua X, Dong Z, Schwarz S, Liu S. 2015.

Novel conjugative plasmid from *Escherichia coli* of swine origin that cohabors the multiresistance gene *cfr* and the extended-spectrum- $\beta$ -lactamase gene *bla*<sub>CTX-M-14b</sub>. *Antimicrob Agents Chemother* 59:1337–1340.

[doi:10.1128/AAC.04631-14](https://doi.org/10.1128/AAC.04631-14).

Address correspondence to Siguo Liu, [siguo\\_liu@hvri.ac.cn](mailto:siguo_liu@hvri.ac.cn), or Stefan Schwarz, [stefan.schwarz@fli.bund.de](mailto:stefan.schwarz@fli.bund.de).

W.-J.Z. and X.-M.W. contributed equally to this article.

Copyright © 2015, American Society for Microbiology. All Rights Reserved.

[doi:10.1128/AAC.04631-14](https://doi.org/10.1128/AAC.04631-14)

TABLE 1 MICs for *E. coli* GXEC3 and GXEC6 and the corresponding *cfr*-positive *E. coli* J53 transconjugants

Strain	MIC (mg/liter) of <sup>a</sup> :								
	AMP	TAZ	GEN	AMK	CHL	FFC	ENO	TET	SMZ
GXEC3	≥256	64	64	16	256	128	≥32	128	32
<i>E. coli</i> J53 (pGXEC3) <sup>b</sup>	256	32	2	2	128	128	0.06	2	16
GXEC6	8	1	64	8	256	128	≥32	128	≥512
<i>E. coli</i> J53 (pGXEC6) <sup>c</sup>	4	0.06	2	2	128	128	0.06	2	16
<i>E. coli</i> J53	4	0.06	1	2	2	1	0.03	2	8

<sup>a</sup> AMP, ampicillin; TAZ, ceftazidime; GEN, gentamicin; AMK, amikacin; CHL, chloramphenicol; FFC, florfenicol; ENO, enrofloxacin; TET, tetracycline; SMZ, sulfamethoxazole.<sup>b</sup> *E. coli* J53 transconjugants of GXEC3.<sup>c</sup> *E. coli* J53 transconjugants of GXEC6.

fied using the ORF Finder program (<http://www.ncbi.nlm.nih.gov/projects/gorf/>). Sequence comparisons were performed using BLASTN and BLASTP (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>).

Although isolates GXEC6 and GXEC3 originated from different farms, PFGE analysis revealed two related XbaI patterns. This observation may be explained by the fact that fattening pig farm B, from

which isolate GXEC3 originated, imports weaned piglets from pig-breeding farm A, from which isolate GXEC6 was obtained. MLST identified each isolate as sequence type (ST) 746, and according to the MLST database, this ST was found in *E. coli* isolates of human and horse origins from Egypt and Germany, respectively. Moreover, a recent study reported an ST746 chicken *E. coli* isolate from China coharboring the *bla*<sub>CTX-M-55</sub> and *bla*<sub>CTX-M-123</sub> genes (19). The *cfr*




FIG 1 Linear comparison of plasmids pGXEC6 and pGXEC3 with *E. coli* (GenBank accession no. JN982327) and pSAM7 from *E. coli* (GenBank accession no. JX981514). The arrows indicate the positions and directions of the transcription of the genes. Gray shading indicates >99% nucleotide sequence identity. Direct repeats are shown in boxes. Δ indicates a truncated gene. A distance scale in kb is displayed below the plasmid sequences.

gene was transferable by conjugation from these two original *E. coli* isolates to the recipient *E. coli* strain J53. Transconjugants were obtained with a high frequency ( $\sim 2 \times 10^{-3}$  to  $3 \times 10^{-3}$  per donor cell). S1-PFGE of the transconjugants revealed approximate sizes of the *cfr*-carrying plasmids of 38 kb (pGXEC6) and 41 kb (pGXEC3). None of the two *cfr*-carrying plasmids produced amplicons by PBRT using the 18 primer pairs described previously (16). While the pGXEC6-carrying transconjugant conferred resistance to only chloramphenicol and florfenicol, the pGXEC3-transconjugant was also resistant to ampicillin and ceftazidime and exhibited an ESBL phenotype (Table 1).

Plasmid pGXEC3 had a size of 41,646 bp, while that of pGXEC6 was 38,405 bp. Comparative sequence analysis of the two plasmids revealed that a 3,241-bp fragment was inserted upstream of the *hicA* gene in pGXEC3 (Fig. 1), whereas the remaining parts of the two plasmids were virtually identical except for several nucleotide substitutions. The insertion included an intact 3,235-bp *ISEcp1-bla<sub>CTX-M-14b</sub>* transposition unit and one 6-bp direct repeat (Fig. 1). The detection of a functionally intact *bla<sub>CTX-M-14b</sub>* gene explained the observed ESBL phenotype. In these two plasmids, the *cfr* gene was located in a 4,612-bp segment, which included a 1,545-bp *cfr*-carrying central region that was identical to the corresponding sequence of plasmid pEC-01 from porcine *E. coli* (5) (Fig. 1). Moreover, comparative analysis of the 13,122-bp *cfr*-carrying fragment of pGXEC6/pGXEC3 and the 12,390-bp *cfr*-carrying fragment of pEC-01 revealed >99% nucleotide sequence identity, except for a 732-bp insertion in pGXEC6/pGXEC3 at positions 5041 to 5772. The *cfr* gene was flanked by two identical copies of the insertion sequence IS26 located in the same orientation. Further database searches revealed that pGXEC3 and pGXEC6 were related to the recently described IncX4 *bla<sub>CTX-M-14b</sub>*-carrying plasmid pSAM7 from bovine *E. coli*, isolated in 2008 in the United Kingdom (20), except for the *cfr*-carrying segment, the *repB* plasmid replication gene region, and a segment containing the *stbDE* toxin/antitoxin genes (Fig. 1). It is tempting to assume that the 4,612-bp region consisting of *tnpA-IS26-cfr-IS26-Δhyp* may represent a transposition unit which has integrated into a pSAM7-like backbone to give rise to pGXEC3. Since *ISEcp1* is able to mobilize itself and adjacent resistance genes, such as *bla<sub>CTX-M</sub>*-type genes (21–23), it is possible that plasmids pGXEC3 and pGXEC6 have developed from one another by either a gain or loss of the *ISEcp1-bla<sub>CTX-M-14b</sub>* segment.

To the best of our knowledge, this is the first report of the *cfr* gene and an ESBL gene (*bla<sub>CTX-M-14b</sub>*) being colocated on a conjugative plasmid in *E. coli*. Given the widespread use of penicillins in veterinary medicine (24), the occurrence and dissemination of plasmids carrying *cfr* and ESBL genes among Gram-negative bacteria need further surveillance.

**Nucleotide sequence accession numbers.** The sequences of plasmids pGXEC3 and pGXEC6 have been deposited in GenBank under accession no. KM580532 and KM580533, respectively.

## ACKNOWLEDGMENTS

This work was funded by the national 973 program (grant 2012CB518801), the National Natural Science Foundation of China (grant 31201862), and the Central Public-Interest Scientific Institution Basal Research Fund (grant 0302014016). The contribution of S.S. was funded by the German Federal Ministry of Education and Research (BMBF) through the German Aerospace Center (DLR), grant 01KI1313D (RESET II) and grant 01KI1301D (MedVet-Staph II).

## REFERENCES

- Long KS, Poehlsgaard J, Kehrenberg C, Schwarz S, Vester B. 2006. The Cfr rRNA methyltransferase confers resistance to phenicols, lincosamides, oxazolidinones, pleuromutilins, and streptogramin A antibiotics. *Antimicrob Agents Chemother* 50:2500–2505. <http://dx.doi.org/10.1128/AAC.00131-06>.
- Shen J, Wang Y, Schwarz S. 2013. Presence and dissemination of the multiresistance gene *cfr* in Gram-positive and Gram-negative bacteria. *J Antimicrob Chemother* 68:1697–1706. <http://dx.doi.org/10.1093/jac/dkt092>.
- Wang Y, Li D, Song L, Liu Y, He T, Liu H, Wu C, Schwarz S, Shen J. 2013. First report of the multiresistance gene *cfr* in *Streptococcus suis*. *Antimicrob Agents Chemother* 57:4061–4063. <http://dx.doi.org/10.1128/AAC.00713-13>.
- Wang Y, Wang Y, Wu CM, Schwarz S, Shen Z, Zhang W, Zhang Q, Shen JZ. 2011. Detection of the staphylococcal multiresistance gene *cfr* in *Proteus vulgaris* of food animal origin. *J Antimicrob Chemother* 66:2521–2526. <http://dx.doi.org/10.1093/jac/dkr322>.
- Wang Y, He T, Schwarz S, Zhou D, Shen Z, Wu C, Wang Y, Ma L, Zhang Q, Shen J. 2012. Detection of the staphylococcal multiresistance gene *cfr* in *Escherichia coli* of domestic animal origin. *J Antimicrob Chemother* 67:1094–1098. <http://dx.doi.org/10.1093/jac/dks020>.
- Zhang WJ, Xu XR, Schwarz S, Wang XM, Dai L, Zheng HJ, Liu S. 2014. Characterization of the IncA/C plasmid pSCEC2 from *Escherichia coli* of swine origin that harbours the multiresistance gene *cfr*. *J Antimicrob Chemother* 69:385–389. <http://dx.doi.org/10.1093/jac/dkt355>.
- Deng H, Sun J, Ma J, Li L, Fang LX, Zhang Q, Liu YH, Liao XP. 2014. Identification of the multi-resistance gene *cfr* in *Escherichia coli* isolates of animal origin. *PLoS One* 9:e102378. <http://dx.doi.org/10.1371/journal.pone.0102378>.
- Witte W, Cuny C. 2011. Emergence and spread of *cfr*-mediated multiresistance in staphylococci: an interdisciplinary challenge. *Future Microbiol* 6:925–931. <http://dx.doi.org/10.2217/fmb.11.69>.
- Smet A, Martel A, Persoons D, Dewulf J, Heyndrickx M, Herman L, Haesebrouck F, Butaye P. 2010. Broad-spectrum beta-lactamases among *Enterobacteriaceae* of animal origin: molecular aspects, mobility and impact on public health. *FEMS Microbiol Rev* 34:295–316. <http://dx.doi.org/10.1111/j.1574-6976.2009.00198.x>.
- Bush K, Jacoby GA. 2010. Updated functional classification of beta-lactamases. *Antimicrob Agents Chemother* 54:969–976. <http://dx.doi.org/10.1128/AAC.01009-09>.
- Naseer U, Sundsfjord A. 2011. The CTX-M conundrum: dissemination of plasmids and *Escherichia coli* clones. *Microb Drug Resist* 17:83–97. <http://dx.doi.org/10.1089/mdr.2010.0132>.
- Wang J, Stephan R, Power K, Yan Q, Hächler H, Fanning S. 2014. Nucleotide sequences of 16 transmissible plasmids identified in nine multidrug-resistant *Escherichia coli* isolates expressing an ESBL phenotype isolated from food-producing animals and healthy humans. *J Antimicrob Chemother* 69:2658–2668. <http://dx.doi.org/10.1093/jac/dku206>.
- Wu G, Day MJ, Mafura MT, Nunez-Garcia J, Fenner JJ, Sharma M, van Essen-Zandbergen A, Rodríguez I, Dierikx C, Kadlec K, Schink AK, Wain J, Helmuth R, Guerra B, Schwarz S, Threlfall J, Woodward MJ, Woodford N, Coldham N, Mevius D. 2013. Comparative analysis of ESBL-positive *Escherichia coli* isolates from animals and humans from the UK, The Netherlands and Germany. *PLoS One* 8:e75392. <http://dx.doi.org/10.1371/journal.pone.0075392>.
- Hansen KH, Bortolaia V, Damborg P, Guardabassi L. 2014. Strain diversity of CTX-M-producing *Enterobacteriaceae* in individual pigs: insights into the dynamics of shedding during the production cycle. *Appl Environ Microbiol* 80:6620–6626. <http://dx.doi.org/10.1128/AEM.01730-14>.
- Zhang WJ, Lu Z, Schwarz S, Zhang RM, Wang XM, Si W, Yu S, Chen L, Liu S. 2013. Complete sequence of the *bla<sub>NDM-1</sub>*-carrying plasmid pNDM-AB from *Acinetobacter baumannii* of food animal origin. *J Antimicrob Chemother* 68:1681–1682. <http://dx.doi.org/10.1093/jac/dkt066>.
- Carattoli A, Bertini A, Villa L, Falbo V, Hopkins KL, Threlfall EJ. 2005. Identification of plasmids by PCR-based replicon typing. *J Microbiol Methods* 63:219–228. <http://dx.doi.org/10.1016/j.mimet.2005.03.018>.
- Clinical and Laboratory Standards Institute. 2013. Performance standards for antimicrobial disk and dilution susceptibility tests for bacteria isolated from animals—4th ed. Approved standard VET01-A4. Clinical and Laboratory Standards Institute, Wayne, PA.

18. Clinical and Laboratory Standards Institute. 2013. Performance standards for antimicrobial disk and dilution susceptibility tests for bacteria isolated from animals, 2nd informational supplement. CLSI VET01-S2. Clinical and Laboratory Standards Institute, Wayne, PA.
19. He D, Partridge SR, Shen J, Zeng Z, Liu L, Rao L, Lv L, Liu JH. 2013. CTX-M-123, a novel hybrid of the CTX-M-1 and CTX-M-9 group beta-lactamases recovered from *Escherichia coli* isolates in China. *Antimicrob Agents Chemother* 57:4068–4071. <http://dx.doi.org/10.1128/AAC.00541-13>.
20. Stokes MO, Abuoun M, Umur S, Wu G, Partridge SR, Mevius DJ, Coldham NG, Fielder MD. 2013. Complete sequence of pSAM7, an IncX4 plasmid carrying a novel *bla*<sub>CTX-M-14b</sub> transposition unit isolated from *Escherichia coli* and *Enterobacter cloacae* from cattle. *Antimicrob Agents Chemother* 57:4590–4594. <http://dx.doi.org/10.1128/AAC.01157-13>.
21. Toleman MA, Walsh TR. 2010. ISCR elements are key players in IncA/C plasmid evolution. *Antimicrob Agents Chemother* 54:3534. <http://dx.doi.org/10.1128/AAC.00383-10>.
22. Freire Martin I, AbuOun M, Reichel R, La Ragione RM, Woodward MJ. 2014. Sequence analysis of a CTX-M-1 IncI1 plasmid found in *Salmonella* 4,5,12:i-, *Escherichia coli* and *Klebsiella pneumoniae* on a UK pig farm. *J Antimicrob Chemother* 69:2098–2101. <http://dx.doi.org/10.1093/jac/dku098>.
23. Wang Y, Song C, Duan G, Zhu J, Yang H, Xi Y, Fan Q. 2013. Transposition of IS<sub>Ecp1</sub> modulates *bla*<sub>CTX-M-55</sub>-mediated *Shigella flexneri* resistance to cefalothin. *Int J Antimicrob Agents* 42:507–512. <http://dx.doi.org/10.1016/j.ijantimicag.2013.08.009>.
24. Silley P, Simjee S, Schwarz S. 2012. Surveillance and monitoring of antimicrobial resistance and antibiotic consumption in humans and animals. *Rev Sci Tech* 31:105–120.