

## **The incorporation of elemental sulfur as fertilizer coating material increased yields of wheat crop**

**D.L. Bouranis<sup>1\*</sup>, D. Argyros<sup>3</sup>, G. Georgoulas<sup>3</sup>, D. Petrakos<sup>3</sup>, A. Varnas<sup>3</sup>, D. Gasparatos<sup>2</sup>, M. Margetis<sup>1</sup>, F. Maniou<sup>1</sup>, S.N. Chorianopoulou<sup>1</sup>, H. Mavrogiannis<sup>3</sup> and D. Benardos<sup>3</sup>**

*<sup>1</sup>Plant Physiology and Morphology Laboratory, Crop Science Department, Agricultural University of Athens, Iera Odos 75, 11855 Athens, Greece (E-mail: bouranis@aua.gr); <sup>2</sup>Soil Science Laboratory, Department of Hydraulics, Soil Science and Agricultural Engineering, School of Agriculture, Faculty of Agriculture, Forestry and Natural Environment, Aristotle University of Thessaloniki, 54124 Thessaloniki, Greece; <sup>3</sup>Sulphur Hellas S.A., Leoforos Athinon 142, 104 42 Athens, Greece*

In a preliminary set of field trials, a number of 11 crops of durum wheat (Simeto, Meridiano and Quantrato cultivars) were established in different areas of central and northern Greece and soil fertility levels were determined prior to crop establishment. Sowing and harvesting took place mid November 2015 and end of June 2016, respectively. Each crop (roughly 2-3 acres each) was divided into two parts: one of them served as the reference crop that received conventional fertilization scheme, whilst the other one (the treatment SG-crop) received the same combination of fertilizers additionally coated with 2% elemental sulfur (Sulfogrow® by Sulphur Hellas S.A.).

Six of the fields were containing no calcium carbonate or traces of it, with pH ranging from 6.20 to 7.74. In five of them, all characterized by adequate fertility levels, the yields of the SG-crops presented relative increases from 8% to 27%. In the field characterized by very low soil initial phosphate content coupled with low potassium content, the yield of the SG-crop presented a relative decrease by 11%.

The other five fields were containing moderate or high calcium carbonate, with pH ranging from 7.96 to 8.20. In three of them, those with adequate fertility levels, the yields of the SG-crops presented relative increases from 3% to 8%. The other two cases of this category presented relative decreases in the yield by 27% and 3.5%, respectively. The rhizospheric soil of these cases was characterized by very low initial phosphate content coupled with marginal iron content, whilst in the former one the rhizospheric soil additionally presented a very low concentration of humic substances and very low sand content.

Bearbeitet von/ Compiled by:

Luit J. De Kok<sup>2</sup>, Silvia Haneklaus<sup>1</sup>, Ewald Schnug<sup>1</sup>

**25<sup>th</sup> International Symposium of the Scientific  
Centre for Fertilizers “Significance of Sulfur in  
High-Input Cropping Systems”  
Groningen (Netherlands), September 5-8, 2017**

<sup>1</sup>Julius Kühn-Institut (JKI), Bundesforschungsinstitut für Kulturpflanzen  
Institut für Pflanzenbau und Bodenkunde

<sup>2</sup>University of Groningen  
Faculty of Science and Engineering

**Berichte aus dem Julius Kühn-Institut**

**191**



**Kontaktadresse/ Contact**

Dr. Silvia Haneklaus  
Federal Research Centre for Cultivated Plants  
Institute for Crop and Soil Science  
Bundesallee 50  
38116 Braunschweig  
Germany

Telefon +49 (0) 531 596 2121  
Telefax +49 (0) 531 596 2199

Wir unterstützen den offenen Zugang zu wissenschaftlichem Wissen.  
Die Berichte aus dem Julius Kühn-Institut erscheinen daher als OPEN ACCESS-Zeitschrift.  
Alle Ausgaben stehen kostenfrei im Internet zur Verfügung:  
<http://www.julius-kuehn.de> Bereich Veröffentlichungen – Berichte.

We advocate open access to scientific knowledge. Reports from the Julius Kühn Institute are therefore published as open access journal. All issues are available free of charge under <http://www.julius-kuehn.de> (see Publications – Reports).

**Herausgeber / Editor**

Julius Kühn-Institut, Bundesforschungsinstitut für Kulturpflanzen, Braunschweig, Deutschland  
Julius Kühn Institute, Federal Research Centre for Cultivated Plants, Braunschweig, Germany

**Vertrieb**

Saphir Verlag, Gutsstraße 15, 38551 Ribbesbüttel  
Telefon +49 (0)5374 6576  
Telefax +49 (0)5374 6577

**ISSN 1866-590X**

**DOI 10.5073/berjki.2017.191.000**



Dieses Werk ist lizenziert unter einer Creative Commons – Namensnennung – Weitergabe unter gleichen Bedingungen – 4.0 Lizenz.  
This work is licensed under a Creative Commons – Attribution – ShareAlike – 4.0 license.