

Linking genes to field performance: adventures in sulfur research

Malcolm J. Hawkesford

Rothamsted Research, West Common, Harpenden, Hertfordshire, AL5 2JQ, U.K. (E-mail: malcolm.hawkesford@rothamsted.ac.uk)

The development of widespread sulfur deficiency in crops became evident in the 1980's and 90's. It is well documented that this arose due to reduced aerial inputs due to decreased pollution as well as the use of low S fertilizer formulations. Negative impacts were seen on high S-requiring crops such as *Brassica napus* (oilseed rape, colza, canola) and also for cereals such as wheat. Both yield and quality were affected, and for example a detrimental impact of S deficiency on the content of S-rich storage proteins of wheat, which are essential for dough rheology, was observed (Zhao et al. 1999). In addition, low S availability has substantial impacts on uptake and accumulation of Se and Mo, which may have substantial implications for nutritional quality (Shinmachi et al. 2010; Stroud et al. 2010). Many studies demonstrated that remedial application of S fertilizers successfully restored yield and quality parameters. One area of importance was accurate diagnosis (Blake-Kalff et al. 2000; Blake-Kalff et al. 2001). Remediation of crop S deficiencies was a recent success story in relation to applied plant nutrition studies coupled with appropriate agronomy.

More fundamental studies have focused on the sulfur uptake and assimilatory pathways and their regulation as a paradigm for plant and indeed crop responses to mineral nutrition (Hawkesford and De Kok 2006). The rationale for such studies is a genetic solution for producing crops with increased efficiency for uptake and utilization of S, or for reduced physiological requirements for S. The most significant advances at the molecular level have been achieved using the model plant *Arabidopsis thaliana*, although such studies have more recently been extended into a number of crops including rice, wheat and Brassica, aided by rapid recent advances in resources for genomics in these crops (Buchner et al. 2004). Achievements over the past 20 years have resulted in the elucidation of molecular aspects of uptake and assimilation pathways, the existence of large multigene families, with isoforms having defined roles, pathways for regulation and inter connections with a range of metabolic pathways and physiological responses. A suggestion that manipulation of feedback mechanisms which restrict uptake could enable luxury uptake during time of plenty (immediately following fertilizer applications) and subsequent effective internal management of this reserve S, could ensure the most effective use of fertilizer applied S. A wide range of roles and requirements for adequate S-nutritional status have been established indicating little opportunity for decreasing crop S demands. One exception has been in identifying prospects for improving effective remobilization of S within the plant particularly, between senescing and newly developing organs (Blake-Kalff et al. 1998; Dubousset et al. 2009). Brassica species have proved to be a useful model for understanding physiological aspects of regulation, having a high S requirement and being

particularly amenable to experimental manipulation of S sources for growth (Aghajanzadeh et al. 2016; Hawkesford and De Kok 2006).

Recent developments in high resolution crop monitoring and phenotyping offer opportunities to closely monitor impacts of S nutrition on crop growth and development, and hence performance in the field. As well as being able to follow crop establishment and determine yield components, opportunities exist to monitor subtle effect of fertilizer on developmental processes such as flowering or other physiological factors such as resistance to stress. Application of an automated field phenotyping platform to monitor crop performance at Rothamsted Research will be presented.

Acknowledgments: Rothamsted Research receives funding from the Biotechnology and Biological Sciences Research Council of the UK.

- Aghajanzadeh T., Hawkesford M.J. and De Kok L.J. (2016) Atmospheric H₂S and SO₂ as sulfur sources for *Brassica juncea* and *Brassica rapa*: Regulation of sulfur uptake and assimilation. *Environ. Exp. Bot.* 124: 1-10.
- Blake-Kalff M.M.A., Harrison K.R., Hawkesford M.J., Zhao F.J. and McGrath S.P. (1998) Distribution of sulfur within oilseed rape leaves in response to sulfur deficiency during vegetative growth. *Plant Physiol.* 118: 1337-1344.
- Blake-Kalff M.M.A., Hawkesford M.J., Zhao F.J. and McGrath S.P. (2000) Diagnosing sulfur deficiency in field-grown oilseed rape (*Brassica napus* L.) and wheat (*Triticum aestivum* L.). *Plant Soil* 225: 95-107.
- Blake-Kalff M.M.A., Zhao F.J., Hawkesford M.J. and McGrath S.P. (2001) Using plant analysis to predict yield losses caused by sulphur deficiency. *Ann. Appl. Biol.* 138: 123-127.
- Buchner P., Prosser I.M., Hawkesford M.J. (2004) Phylogeny and expression of paralogous and orthologous sulphate transporter genes in diploid and hexaploid wheats. *Genome* 47: 526-534.
- Dubousset L., Abdallah M., Desfeux A.S., Etienne P., Meuriot F., Hawkesford M.J., Gombert J., Segura R., Bataille M.P., Reze S., Bonnefoy J., Ameline A.F., Ourry A., Le Dily F. and Avice J.C. (2009) Remobilization of leaf S compounds and senescence in response to restricted sulphate supply during the vegetative stage of oilseed rape are affected by mineral N availability. *J. Exp. Bot.* 60: 3239-3253.
- Hawkesford M.J. and De Kok L.J. (2006) Managing sulphur metabolism in plants. *Plant Cell Environ* 29: 382-395.
- Shinmachi F., Buchner P., Stroud J.L., Parmar S., Zhao F.J., McGrath S.P. and Hawkesford M.J. (2010) Influence of sulfur deficiency on the expression of specific sulfate transporters and the distribution of sulfur, selenium, and molybdenum in wheat. *Plant Physiol.* 153: 327-336.
- Stroud J.L., Zhao F.J., Buchner P., Shinmachi F., McGrath S.P., Abecassis J., Hawkesford MJ and Shewry P.R. (2010) Impacts of sulphur nutrition on selenium and molybdenum concentrations in wheat grain. *J. Cereal Sci.* 52: 111-113.
- Zhao F.J., Hawkesford M.J. and McGrath S.P. (1999) Sulphur assimilation and effects on yield and quality of wheat. *J. Cereal Sci* 30: 1-17.

Bearbeitet von/ Compiled by:

Luit J. De Kok², Silvia Haneklaus¹, Ewald Schnug¹

**25th International Symposium of the Scientific
Centre for Fertilizers “Significance of Sulfur in
High-Input Cropping Systems”
Groningen (Netherlands), September 5-8, 2017**

¹Julius Kühn-Institut (JKI), Bundesforschungsinstitut für Kulturpflanzen
Institut für Pflanzenbau und Bodenkunde

²University of Groningen
Faculty of Science and Engineering

Berichte aus dem Julius Kühn-Institut

191



Kontaktadresse/ Contact

Dr. Silvia Haneklaus
Federal Research Centre for Cultivated Plants
Institute for Crop and Soil Science
Bundesallee 50
38116 Braunschweig
Germany

Telefon +49 (0) 531 596 2121
Telefax +49 (0) 531 596 2199

Wir unterstützen den offenen Zugang zu wissenschaftlichem Wissen.
Die Berichte aus dem Julius Kühn-Institut erscheinen daher als OPEN ACCESS-Zeitschrift.
Alle Ausgaben stehen kostenfrei im Internet zur Verfügung:
<http://www.julius-kuehn.de> Bereich Veröffentlichungen – Berichte.

We advocate open access to scientific knowledge. Reports from the Julius Kühn Institute are therefore published as open access journal. All issues are available free of charge under <http://www.julius-kuehn.de> (see Publications – Reports).

Herausgeber / Editor

Julius Kühn-Institut, Bundesforschungsinstitut für Kulturpflanzen, Braunschweig, Deutschland
Julius Kühn Institute, Federal Research Centre for Cultivated Plants, Braunschweig, Germany

Vertrieb

Saphir Verlag, Gutsstraße 15, 38551 Ribbesbüttel
Telefon +49 (0)5374 6576
Telefax +49 (0)5374 6577

ISSN 1866-590X

DOI 10.5073/berjki.2017.191.000



Dieses Werk ist lizenziert unter einer [Creative Commons – Namensnennung – Weitergabe unter gleichen Bedingungen – 4.0 Lizenz](https://creativecommons.org/licenses/by-sa/4.0/).
This work is licensed under a [Creative Commons – Attribution – ShareAlike – 4.0 license](https://creativecommons.org/licenses/by-sa/4.0/).