

# Soil innate microbiome shows the potential to protect crops against human pathogens

Jasper Schierstaedt<sup>1,2</sup>, Helena J. Barkowski<sup>2</sup>, Sven Jechalke<sup>3</sup>, Abhishek Shrestha<sup>2</sup>, Kornelia Smalla<sup>2</sup>, Rita Grosch<sup>1</sup> and Adam Schikora<sup>2</sup>

<sup>1</sup> Leibniz Institute of Vegetable and Ornamental Crops, Department Plant Health, Grossbeeren

<sup>2</sup> Julius Kühn-Institut, Institute for Epidemiology and Pathogen Diagnostics, Braunschweig

<sup>3</sup> JLU Gießen, Institute of Phytopathology, Gießen

E-mail of corresponding author: jasper.schierstaedt@julius-kuehn.de

Foodborne diseases are increasingly associated with fresh fruits and vegetables. Serovars of *Salmonella* were the second most frequent cause in 2015. The biological diversity of soil plays a major role in the establishment of *Salmonella* in the plant production environment.

Antagonists and plant beneficial microbes negatively affect the establishment of *Salmonella* in the rhizosphere. Here we analyzed the tripartite interactions between the human pathogen *Salmonella enterica*, the soil microbiome and the crop plants tomato, lettuce and corn salad grown under greenhouse conditions.

We observed that *Salmonella* persisted in the rhizosphere of lettuce and tomato. In contrast, its numbers declined in the rhizosphere of corn salad. Very important was the observation that reduction of microbial diversity in soil increased the ability of *Salmonella* to persist in this en-

vironment. These results clearly show a dependency between the microbial diversity and the potential of *Salmonella* to colonize the rhizosphere as well as the high physiological plasticity of *Salmonella*. In the following, we focused on the impact of resistance induced in crop plants on the establishment in plant production environment and colonization of plants. In greenhouse experiments, crop plants were primed for induced resistance with the bacterium *Ensifer meliloti*. This bacterium produces *N*-acyl-homoserine-lactones, which are known to induce resistance.

Our results show that priming has a negative effect on the persistence of *Salmonella*. Primed plants express the defense-related genes earlier than unprimed plants and are able to close their stomata for longer period. These results indicate the potential of priming for enhanced resistance against *Salmonella*.



# 11<sup>th</sup> Young Scientists Meeting 2018

14<sup>th</sup> – 16<sup>th</sup> November  
in Braunschweig

- Abstracts -



Berichte aus dem Julius Kühn-Institut

# 200

## **Kontaktadresse/ Contact**

Dr. Anja Hühnlein  
Julius Kühn-Institut (JKI)  
Bundesforschungsinstitut für Kulturpflanzen  
Informationszentrum und Bibliothek  
Erwin-Baur-Straße 27  
06484 Quedlinburg  
Germany

Telefon +49 (0) 3946 47-123

Telefax +49 (0) 3946 47-255

Wir unterstützen den offenen Zugang zu wissenschaftlichem Wissen.  
Die Berichte aus dem Julius Kühn-Institut erscheinen daher als OPEN ACCESS-Zeitschrift.  
Alle Ausgaben stehen kostenfrei im Internet zur Verfügung:  
<http://www.jki.bund.de> Bereich Veröffentlichungen – Berichte.

We advocate open access to scientific knowledge. Reports from the Julius Kühn Institute are therefore published as open access journal. All issues are available free of charge under <http://www.jki.bund.de> (see Publications – Reports).

## **Herausgeber / Editor**

Julius Kühn-Institut, Bundesforschungsinstitut für Kulturpflanzen, Braunschweig, Deutschland  
Julius Kühn Institute, Federal Research Centre for Cultivated Plants, Braunschweig, Germany

## **Vertrieb**

Saphir Verlag, Gutsstraße 15, 38551 Ribbesbüttel  
Telefon +49 (0)5374 6576  
Telefax +49 (0)5374 6577

**ISSN 1866-590X**

**DOI 10.5073/berjki.2018.200.000**



Dieses Werk ist lizenziert unter einer [Creative Commons – Namensnennung – Weitergabe unter gleichen Bedingungen – 4.0 Lizenz](https://creativecommons.org/licenses/by-sa/4.0/).

This work is licensed under a [Creative Commons – Attribution – ShareAlike – 4.0 license](https://creativecommons.org/licenses/by-sa/4.0/).