

Two obesogenic diets differently modulate microbial dysbiosis and adipose tissue inflammation in C57Bl/6 male mice.

Krista A. Power^{1,2,3}, Kyle Robertson², Daniela Graf^{2,3,4}, Jennifer Monk², Dion Lepp³, Wenqing Wu³

¹School of Nutrition Sciences, University of Ottawa, Canada; ²Department of Human Health and Nutrition Science, University of Guelph, Canada; ³Guelph Research and Development Centre, Agriculture and Agri-food Canada; ⁴Department of Physiology and Biochemistry of Nutrition, Max Rubner-Institute, Germany

Objectives: Obesogenic diets are commonly used to induce obesity in rodent models, however, the nutritional composition (e.g. fat and CHO level) often varies dramatically which may result in variable effects on the obese phenotype. Therefore, the objective was to compare the effects of two commonly used obesogenic diets on the intestinal microenvironment and adipose tissue dysfunction in mice.

Methods: 3 week old male C57Bl/6 mice were fed diets comprised of 17% fat (%kcal; low fat diet; LFD), 45% fat (%kcal; 45HFD), or 60% fat (%kcal; 60HFD) for 12wks and aspects of the obese phenotype (weight gain, body mass index (BMI), oral glucose tolerance, homeostatic model assessment of insulin resistance (HOMA-IR), adipose dysfunction) and intestinal microenvironment (microbiota community structure (16S rRNA gene sequencing), mucosal inflammation and barrier integrity) were assessed.

Results: Mice consuming 60HFD gained more weight compared to 45HFD and LFD, resulting in significant differences in final BMI and white adipose tissue (WAT) weights. WAT dysfunction was more severe in the 60HFD group (e.g. increased crown-like structures (macrophage infiltration) and serum adipokines (leptin and resistin)) compared to the 45HFD and LFD. Blood glucose clearance following an oral glucose load and HOMA-IR were significantly higher in the 60HFD group compared to 45HFD and LFD. Cecal and fecal microbial dysbiosis was also more severe in the 60HFD group compared to the 45HFD and LFD, including reduced phylogenetic diversity and *Akkermansia muciniphila* abundance. Colon mRNA expression of TLR2 and IL-6 were significantly increased by 60HFD, while tight junction protein expression (JAM-A) was decreased, compared to LFD and 45HFD, whereas both HFDs increased TNF- α and MCP-1 expression. Serum lipopolysaccharide binding protein, a biomarker of endotoxemia, was increased by the 60HFD compared to 45HFD and LFD.

Conclusions: Despite interchangeable use of different obesogenic diets in obesity research, clear differences in the obese phenotype and intestinal microenvironment composition and function result. This study highlights the importance of properly defining the obesogenic diet composition to accurately model the human clinical phenotype in obesity research.

Ontario Research Fund: Research Excellence; Agriculture and Agri-Food Canada; German Research Foundation; uOttawa