

Suitability of hemp seed for reproduction of stored-product insects

Kim Stadnyk¹, Noel D.G. White¹, Fuji Jian², Paul G. Fields¹

¹Morden Research and Development Centre, Agriculture and Agri-Food Canada, kim.stadnyk@agr.gc.ca, noel.white@agr.gc.ca, paul.fields@agr.gc.ca

²Biosystems Engineering, University of Manitoba, Winnipeg, MB, Canada, jianf@cc.umanitoba.ca
DOI 10.5073/jka.2018.463.041

Extended Abstract

1. Introduction

Hemp, or industrial hemp, is a high value alternative crop that has seen major increases in acreage in Canada since commercial production was legalized in 1998. The term industrial hemp applies to non-psychoactive varieties of *Cannabis sativa* L. There have been reports of insect infestations on stored hemp seed in Manitoba. The moths *Plodia interpunctella* (Hübner) Indianmeal moth, and *Ephesia kuehniella* (Zeller) Mediterranean flour moth feed on hemp seed (Hagstrum and Subramanyam, 2009). Our objectives were to determine which stored-product beetles can reproduce on hemp and the effect of dockage and seed moisture content.

2. Materials and Methods

Twenty adult insects were placed on 15 g of hemp seed at two different moisture contents (~8% or ~15%) and two different dockage levels (~0% or ~15%) and held at 30°C and 60-70% relative humidity. The number of live and dead insects were counted at 3, 5, 7 and 9 weeks. Only live adults were returned to vials.

3. Results and Discussion

These insect populations increased over the 9 weeks; red flour beetle [*Tribolium castaneum* (Herbst)], drugstore beetle [*Lasioderma serricorne* (F.)] saw-toothed grain beetle [*Oryzaephilus surinamensis* (L.)], warehouse beetle (*Trogoderma variabile* Ballion). These insect populations did not increase: rusty grain beetle [*Cryptolestes ferrugineus* (Stephens)], lesser grain borer [*Rhyzopertha dominica* (F.)], rice weevil [*Sitophilus oryzae* (L.)], flour mill beetle (*Cryptolestes turcicus* (Grouvelle), confused flour beetle (*Tribolium confusum* Jacquelin du Val), cigarette beetle [*Stegobium paniceum* (L.)]. In general, higher dockage led to higher populations. The effect of moisture content was variable.

Keywords: *Cannabis sativa*, reproduction, dockage, moisture

Acknowledgements

We thank Colin Deminayk, Whitney Morse and Emily Hanuschuk for technical assistance.

References

HAGSTRUM, D.W., SUBRAMANYAM, B., 2009. Stored-product insect resource. American Association of Cereal Chemists, Inc (AACC).

The use of long-lasting insecticide netting to prevent dispersal of stored product insects

William R. Morrison III^{1*}, Rachel V. Wilkins²

¹ USDA, Agricultural Research Service, Center for Grain and Animal Health Research, Manhattan, KS 66502, USA

² Department of Entomology, Kansas State University, Manhattan, KS 66502, USA

*Corresponding author: William.morrison@ars.usda.gov

DOI 10.5073/jka.2018.463.042