



# ICTV Virus Taxonomy Profile: *Hepeviridae* 2022

Michael A. Purdy<sup>1</sup>, Jan Felix Drexler<sup>2</sup>, Xiang-Jin Meng<sup>3</sup>, Heléne Norder<sup>4,5</sup>, Hiroaki Okamoto<sup>6</sup>, Wim H. M. Van der Poel<sup>7</sup>, Gábor Reuter<sup>8</sup>, William M. de Souza<sup>9</sup>, Rainer G. Ulrich<sup>10</sup>, Donald B. Smith<sup>11,\*</sup> and ICTV Report Consortium

## Abstract

The family *Hepeviridae* includes enterically transmitted small quasi-enveloped or non-enveloped positive-sense single-stranded RNA viruses infecting mammals and birds (subfamily *Orthohepevirinae*) or fish (*Parahepevirinae*). Hepatitis E virus (genus *Paslahepevirus*) is responsible for self-limiting acute hepatitis in humans; the infection may become chronic in immunocompromised individuals and extrahepatic manifestations have been described. Avian hepatitis E virus (genus *Avihepevirus*) causes hepatitis-splenomegaly syndrome in chickens. This is a summary of the International Committee on Taxonomy of Viruses (ICTV) Report on the family *Hepeviridae*, which is available at [www.ictv.global/report/hepeviridae](http://www.ictv.global/report/hepeviridae).

**Table 1.** Characteristics of members of the family *Hepeviridae*

Example:	human hepatitis E virus Burma (M73218), species <i>Paslahepevirus balayani</i> , genus <i>Paslahepevirus</i>
Virion	Quasi-enveloped or non-enveloped, 27–34 nm diameter with a single capsid protein
Genome	6.4–7.2 kb capped positive-sense monopartite RNA containing three ORFs
Replication	Occurs in association with the host endoplasmic reticulum
Translation	From genomic (ORF1) and a subgenomic (ORF2 and ORF3) capped mRNA
Host range	Mammals ( <i>Chirohepevirus</i> , <i>Paslahepevirus</i> , <i>Rocahepevirus</i> ), birds ( <i>Avihepevirus</i> ) and salmonid fishes ( <i>Piscihepevirus</i> )
Taxonomy	Realm <i>Riboviria</i> , kingdom <i>Orthornavirae</i> , phylum <i>Kitrinoviricota</i> , class <i>Alsuviricetes</i> , order <i>Hepelivirales</i> : two subfamilies, multiple genera and species

## VIRION

Virions of human hepatitis E virus are icosahedral, quasi-enveloped (blood or tissue culture [1]) or non-enveloped (faeces), spherical particles with a diameter of 27–34 nm (Table 1, Fig. 1a). The capsid is formed by capsomeres consisting of homodimers of a single capsid protein, forming the virus shell. Each capsid protein contains three linear domains forming distinct structural elements: S (the continuous capsid), P1 (three-fold protrusions) and P2 (two-fold spikes). Neutralizing epitopes have been found in the P2 domain. Each domain contains a putative polysaccharide-binding site that may interact with cellular receptors [2].

## GENOME


Viral genomes (Fig. 1b) are positive-sense monopartite RNA of 6.4–7.2 kb, with three ORFs flanked by short 5'- and 3'-terminal non-coding regions: ORF2 (capsid protein) overlaps ORF3 but neither overlaps ORF1. The 5'-end is m<sup>7</sup>G-capped and the 3'-end is polyadenylated (Fig. 2). Non-structural proteins encoded by ORF1 have limited similarity with the 'alpha-like supergroup' of viruses and contain domains consistent with a methyltransferase, papain-like cysteine protease, macro domain, RNA helicase and RNA-directed RNA polymerase [3]. A small immunoreactive protein (12.5 kDa) encoded by ORF3 has been shown to

Received 02 June 2022; Accepted 11 June 2022; Published 28 September 2022

**Author affiliations:** <sup>1</sup>Centers for Disease Control and Prevention, Atlanta, GA, USA; <sup>2</sup>Charité-Universitätsmedizin Berlin, Corporate Member of Freie Universität Berlin, Humboldt-Universität zu Berlin, Institute of Virology, Berlin, Germany; <sup>3</sup>Virginia Polytechnic Institute and State University, Blacksburg, VA, USA; <sup>4</sup>University of Gothenburg, Gothenburg, Sweden; <sup>5</sup>Sahlgrenska University Hospital, Gothenburg, Sweden; <sup>6</sup>Jichi Medical University School of Medicine, Tochigi, Japan; <sup>7</sup>Wageningen University and Research, Lelystad, The Netherlands; <sup>8</sup>Department of Medical Microbiology and Immunology, Medical School, University of Pécs, Pécs, Hungary; <sup>9</sup>University of Texas Medical Branch at Galveston, Galveston, Texas, USA; <sup>10</sup>Friedrich-Loeffler-Institut, Greifswald-Insel Riems, Riems, Germany; <sup>11</sup>University of Oxford, England and University of Edinburgh, Scotland, UK.

\*Correspondence: Donald B. Smith, [donald.smith.mail@gmail.com](mailto:donald.smith.mail@gmail.com)

**Keywords:** hepatitis E virus; *Hepeviridae*; ICTV report; taxonomy.



**Fig. 1.** (a) Negative-contrast electron micrograph of human hepatitis E virus particles from a case stool collected in Nepal. (V) virion and (E) empty capsid. Bar: 100 nm (photograph M. Purdy). (b) Phylogeny of orthohepevirin methyltransferase domains (details: ICTV Report).

exhibit multiple functions associated with virion morphogenesis, egress and viral pathogenesis. The capsid and ORF3 proteins are translated from a subgenomic RNA.

## REPLICATION


The viral RNA-directed RNA polymerase associates with the host endoplasmic reticulum through residues encoding a predicted transmembrane domain. Replication involves temporal separation and alternating cycles of synthesis of positive- and negative-sense RNAs [3, 4].

## TAXONOMY

Current taxonomy: [ictv.global/taxonomy](http://ictv.global/taxonomy). Members of the subfamily *Orthohepevirinae* infect humans and domestic and wild mammals, (genera *Paslahepevirus* and *Rocahepevirus*), bats (*Chirohepevirus*), and birds (*Avihepevirus*) [5, 6]. Human hepatitis E virus can cause self-limiting acute hepatitis in humans and is transmitted by contaminated water, the consumption of undercooked or raw meat or iatrogenically through blood transfusion or organ transplantation. Globally, human hepatitis E virus is a major cause of acute hepatitis.

### References

1. Yin X, Ambardekar C, Lu Y, Feng Z. Distinct entry mechanisms for nonenveloped and quasi-enveloped hepatitis E viruses. *J Virol* 2016;90:4232–4242.
2. Mori Y, Matsuura Y. Structure of hepatitis E viral particle. *Virus Res* 2011;161:59–64.
3. Cao D, Meng XJ. Molecular biology and replication of hepatitis E virus. *Emerg Microbes Infect* 2012;1:e17.



**Fig. 2.** Genome organization of hepeviruses. There are short 5'- and 3'-non-coding regions. The ORF1 polyprotein includes the putative functional domains: MT, methyltransferase; P, a putative papain-like cysteine protease; HUD, hepevirus unique (or Z-) domain; PP, a hypervariable polyproline region; Macro, macro domain; Hel, helicase; and RdRP, RNA-directed RNA polymerase. ORF2 encodes the capsid protein (CP) and ORF3 a small phosphoprotein.

Chronic hepatitis E virus infection has increasingly become an important clinical problem in immunocompromised individuals. Cutthroat trout virus, a member of the subfamily *Parahepevirinae*, infects trout and salmon, although its pathogenicity and full host range are unknown.

## RESOURCES

Full ICTV Report on the family *Hepeviridae* : [www.ictv.global/report/hepeviridae](http://www.ictv.global/report/hepeviridae).

### Funding information

Production of this summary, the online chapter and associated resources was funded by a grant from the Wellcome Trust (WT108418AIA).

### Acknowledgements

Members of the ICTV Report Consortium are Stuart G. Siddell, Elliot J. Lefkowitz, Sead Sabanadzovic, F. Murilo Zerbini, Donald B. Smith, and Peter Simmonds.

### Conflicts of interest

The authors declare that there are no conflicts of interest.

### Disclaimer

M. Purdy - CDC/ATSDR has not formally disseminated the conclusions in this report; they do not represent and should not be construed to represent any agency determination or policy.

4. Varma SPK, Kumar A, Kapur N, Durgapal H, Acharya SK, et al. Hepatitis E virus replication involves alternating negative- and positive-sense RNA synthesis. *J Gen Virol* 2011;92:572–581.
5. Smith DB, Simmonds P, Jameel S, Emerson SU, et al. Consensus proposals for classification of the family *Hepeviridae*. *J Gen Virol* 2014;95:2223–2232.
6. Wang B, Meng XJ. Hepatitis E virus: host tropism and zoonotic infection. *Curr Opin Microbiol* 2021;59:8–15.