

RESEARCH

Open Access



# Modulation of the porcine intestinal microbiota in the course of *Ascaris suum* infection

Andrea Springer<sup>1</sup>, Liane Wagner<sup>1,2</sup>, Sarina Koehler<sup>1</sup>, Stefanie Klinger<sup>3</sup>, Gerhard Breves<sup>3</sup>, Dagmar A. Brüggemann<sup>2</sup> and Christina Strube<sup>1\*</sup>

## Abstract

**Background:** The porcine roundworm *Ascaris suum* impairs feed conversion and weight gain, but its effects on intestinal microbiota remain largely unexplored.

**Methods:** Modulation of the intestinal microbiota was assessed in pigs that were infected once with 10,000 *A. suum* eggs and pigs that received a trickle infection (1000 eggs/day over 10 days), compared with a non-infected control group. Six pigs each were sacrificed per group at days 21, 35 and 49 post-infection (p.i.). Faecal samples taken weekly until slaughter and ingesta samples from different intestinal compartments were subjected to next-generation sequencing of the bacterial 16S rRNA gene.

**Results:** The results revealed marked differences between the single- and the trickle-infected group. Single infection caused a remarkable but transient decrease in microbial diversity in the caecum, which was not observed in the trickle-infected group. However, an increase in short-chain fatty acid-producing genera in the caecum on day 21 p.i., which shifted to a decrease on day 35 p.i., was common to both groups, possibly related to changes in excretory–secretory products following the parasite’s final moult. Faecal microbial interaction networks were more similar between the single-infected and control group than the trickle-infected group. In addition, a lower degree of similarity over time indicated that *A. suum* trickle infection prevented microbiota stabilization.

**Conclusions:** These different patterns may have important implications regarding the comparability of experimental infections with natural scenarios characterized by continuous exposure, and should be confirmed by further studies.

**Keywords:** Ascarids, Roundworms, Trickle infection, Intestinal microbiota, Microbiome, Network analysis, Short-chain fatty acids

\*Correspondence: Christina.strube@tiho-hannover.de

<sup>1</sup> Institute for Parasitology, Centre for Infection Medicine, University of Veterinary Medicine Hannover, Buenteweg 17, 30559 Hanover, Germany  
Full list of author information is available at the end of the article



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article’s Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article’s Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

## Background

Gastrointestinal parasites share their habitat with a diverse community of bacteria, the gastrointestinal microbiota, which plays a profound role in host metabolism, physiological development, immune regulation and overall health [1]. Consequently, multilateral relationships exist between intestinal parasites, the microbiota and host cells, shaping the physiological consequences of infection [2].

The roundworm *Ascaris suum* is one of the economically most important endoparasites in commercial pig farming, and is closely related to or even conspecific with the human roundworm *Ascaris lumbricoides* [3], which affects approximately 800 million people worldwide [4]. After oral infection with embryonated eggs, *A. suum* larvae penetrate the caecal or colonic mucosa, enter the bloodstream and undergo a body migration via the liver and the lungs, before reaching the small intestine approximately 8–10 days post-infection (p.i.) [5]. In the jejunum, they continue their development; however, a large proportion of the fourth-stage larvae (L4) are eliminated between days 17 and 21 p.i., irrespective of the infection dose. This so-called self-cure phenomenon is associated with an increase in mucosal eosinophils, macrophages and T-cells and decreased gastrointestinal transit time [6]. The remaining larvae undergo a final moult between days 21 and 29 p.i. [7], and the onset of patency occurs approximately 6–8 weeks p.i. [8].

*Ascaris suum*-related production losses result from reduced weight gain and feed conversion efficiency, in addition to condemnation of livers showing characteristic lesions due to the parasite's body migration (so-called milk spots), costs for helminth control, and indirect, immunomodulatory effects of the parasite [9]. Although it has been difficult to demonstrate consistent changes in feed utilization and weight development of infected versus non-infected pigs, several experimental and field studies have reported significant differences (reviewed in [9]). For example, a 12% reduction in live weight gain was found in *A. suum*-infected versus anthelmintic-treated pigs in an outdoor system [10]. Reduced feed utilization efficiency is presumably due to a parasite-induced reduction in sodium-coupled intestinal glucose resorption [11–14]; however, alterations of the intestinal microbiota might also play a role. It has been suggested that *A. suum* actively manipulates its microbial environment by the release of potent antimicrobial proteins and peptides, thereby creating a favourable metabolic environment and immunoregulatory state [2, 15]. Furthermore, changes in intestinal physiology during infection, such as increased or disrupted mucus production, are likely to affect microbial populations in the gut, as suggested by an increase in *Mucispirillum* spp. in *Trichuris suis*-infected pigs [16].


Indeed, changes in the porcine intestinal microbiota have been observed during the early phase (day 14 p.i.) of *A. suum* infections [17, 18], as well as on day 54 p.i. [19]. The results were somewhat incongruent, since increased colonic microbial diversity was noted in the early phase [17], but a decrease was seen in the chronic phase of infection [19]. Furthermore, since *A. suum* resides in the small intestine, effects in other intestinal compartments may differ from those observed in the colon [18]. Therefore, the present study aimed to comprehensively characterize microbial changes during the course of *A. suum* infection by analysing faecal samples taken in weekly intervals until day 49 p.i., and ingesta samples from the jejunum, ileum, caecum and colon taken at days 21, 35 and 49 p.i. Furthermore, continuous, low-dose infections are common under natural conditions, and may have different consequences from those of the single-dose experimental infections mentioned above [20]. The current study therefore compared single-dose-infected pigs and pigs trickle-infected over the course of 10 days to an uninfected control group.

## Methods

### Experimental *A. suum* infections and sample collection

*Ascaris suum* eggs were obtained from adult worms collected at an abattoir and allowed to embryonate at 25 °C for 2 months. Embryonated eggs were subsequently kept at 4 °C until oral infection of pigs.

Infection experiments were performed in a project on intestinal nutrient transport [13]. Fifty-four helminth-free hybrid German Landrace pigs weighing approximately 10 kg and 5 weeks of age were acquired from the *A. suum*-free Ruthe Research and Education Farm of the University of Veterinary Medicine Hannover, Germany. Faecal samples were taken rectally on the day of the pigs' arrival and subjected to coproscopic analysis to verify they were helminth-free. The pigs were divided into three groups, which were kept in rooms with concrete flooring on straw bedding. They received a standard pig diet and water ad libitum (Deuka Ferkelstarter Primo, Deutsche Tiernahrung Cremer, Düsseldorf, Germany). After a minimum adaptation period of 1 week, 18 pigs each received either a single oral infection of 10,000 embryonated *A. suum* eggs or a trickle infection of 1000 eggs/day for 10 days (10,000 eggs in total), while a further 18 pigs served as a non-infected control. To ensure that all pigs had received the same overall infection dose by the time of investigation, the trickle infection was not spread over more than 10 days. Pigs were kept until slaughter on days 21, 35 and 49 p.i. ( $N=6$  per group and time point), respectively. Serum samples were taken weekly from each pig and subjected to anti-*Ascaris* antibody detection to verify successful infection [human anti-*Ascaris*



*lumbricoides* immunoglobulin G (IgG) enzyme-linked immunosorbent assay (ELISA, Abcam (Netherlands) B.V., Amsterdam, Netherlands]. In addition, macroscopically visible worms were collected from the intestines upon slaughter. To monitor weight gain, pigs were weighed upon arrival and on the day of necropsy.

Faecal samples for microbiota analysis were taken rectally 3 days prior to infection and on days 3, 7, 14, 21, 28, 35, 42 and 49 p.i., and stored at  $-80^{\circ}\text{C}$  until further processing. Intestinal contents were sampled with sterile tools from the proximal jejunum (approximately 3 m distal to the pylorus), medial jejunum (approximately 6–8 m distal to the pylorus), ileum, caecum and colon ascendens (second coil) and frozen immediately in liquid nitrogen, before storage at  $-80^{\circ}\text{C}$ . The experimental setup and sampling scheme is visualized in Fig. 1.

#### DNA isolation, polymerase chain reaction (PCR) and sequence processing

DNA isolation from faecal samples was performed with the DNeasy<sup>®</sup> PowerLyzer<sup>®</sup> PowerSoil<sup>®</sup> kit (Qiagen, Hilden, Germany) with a homogenization step of  $2 \times 45$  s at 5000 rpm in a Precellys<sup>®</sup> 24 tissue homogenizer (Peqlab Biotechnologie GmbH, Erlangen, Germany). The procedure followed the manufacturer's instructions, except that centrifugation time at step 10 was increased to 2 min. For ingesta samples, the QIAamp<sup>®</sup> BiOstic<sup>®</sup> Bacteremia DNA Kit (Qiagen, Hilden, Germany) was used, as this kit has been shown to increase bacterial DNA yield in low-biomass (i.e. small intestine) samples [21]. All steps followed the manufacturer's instructions, except that bead-beating was performed with the Precellys<sup>®</sup> 24 as described above, and centrifugation time at step 7 was increased to 2 min. Eluted DNA was stored at  $-20^{\circ}\text{C}$  until shipment to a commercial laboratory (Microsynth AG, Balgach, Switzerland).

Illumina MiSeq sequencing, sequence processing, operational taxonomic unit (OTU) clustering and taxonomic assignment were performed by Microsynth AG (Balgach, Switzerland), according to Microsynth's standard operating procedures. After dilution of the samples to  $2.5 \text{ ng DNA}/\mu\text{l}$ , two-step Nextera PCR libraries of the V3–V4 region of the bacterial 16S rRNA gene were created using KAPA HiFi Mastermix (Roche Sequencing and Life Science, Indianapolis, IN, USA). The first-step PCR with primers 341F and 805R ( $4 \mu\text{m}$  each) [22] and the second-step PCR with index primers ( $4 \mu\text{m}$  each, including Illumina adapters) comprised 20 and 10 cycles of the following thermoprofile, respectively:  $95^{\circ}\text{C}$  3 min;  $20/10 \times 98^{\circ}\text{C}$  20 s,  $56^{\circ}\text{C}$  30 s,  $72^{\circ}\text{C}$  30 s;  $72^{\circ}\text{C}$  5 min. The second-step PCR products were bead-purified, quantified with fluorescence spectroscopy and pooled in equimolar amounts. The Illumina MiSeq platform and a v3 600-cycle kit (both Illumina Inc., San Diego, CA, USA) were used for sequencing. The resulting paired-end reads which passed Illumina's chastity filter were subjected to de-multiplexing and adaptor trimming using Illumina's real-time analysis software included in the MiSeq Reporter software v. 2.6 (no further refinement or selection). Read quality was checked with FastQC v. 0.11.8 (Babraham Institute, Cambridge, UK), and the locus-specific V3–V4 primers were trimmed with Cutadapt v. 2.8 [23], discarding reads if primer trimming failed. Further, the reads were trimmed from the original 300 base pairs (bp) to 250 bp from the 3' end to reduce sequencing noise. Trimmed forward and reverse reads were merged in silico, considering a minimum overlap of 15 bases using USEARCH v. 11.0.667 [24]. Merged sequences were quality-filtered, allowing a maximum of one expected error per merged read. Reads with ambiguous bases and outliers regarding amplicon size were

discarded. The remaining reads were denoised via the UNOISE algorithm [25] to form OTUs, discarding singletons and chimeras. The resulting OTU abundance table was filtered for possible bleed-in contaminations using the UNCROSS algorithm [26]. OTUs were compared against the reference sequences of the Ribosomal Database Project (RDP) 16S database [27], and taxonomies were predicted considering a minimum confidence threshold of 0.5 using the SINTAX algorithm implemented in USEARCH [28]. OTU clustering and taxonomic assignment were performed separately for faecal samples and for small intestine (jejunum, ileum) and large intestine (colon, caecum) samples, resulting in three separate datasets.

### Statistical analyses

Statistical analyses were conducted in R v. 4.1.0 [29], using the packages phyloseq [30], vegan [31], DESeq2 [32] and lmerTest [33]. The average daily weight gain until day 21, day 35 and day 49 p.i. was compared between the experimental groups by Kruskal–Wallis tests, followed by Dunn’s post hoc tests.

Prior to calculating microbial alpha and beta diversity indices, the faecal sample dataset was rarefied to 3000 reads/sample and the ingesta datasets to 8000 reads/sample. Rarefaction curves were inspected to verify that the level of rarefaction adequately captured the phylogenetic diversity.

To assess microbial diversity within each sample (i.e. alpha diversity), the number of observed OTUs and the Chao1, Shannon and inverse Simpson indices were calculated, with the latter two taking not only the absence/presence, but also the abundance of OTUs into account [34]. For faecal samples, the influence of day p.i., experimental group and their interaction on alpha diversity was assessed by generalized linear mixed models (GLMMs), with animal ID as a random factor due to repeated sampling of individuals. A Poisson error structure with identity link was used to model observed OTUs, while a gamma distribution with identity link proved more appropriate for Chao1 and Shannon indices, and a gamma distribution with log link for the inverse Simpson index. Full models were compared with null models containing only the random factor in a maximum likelihood test, and the distribution of model residuals as well as residuals against fitted values were inspected graphically to assess model fit. Alpha diversity of ingesta samples was compared between experimental groups for each intestinal segment and day p.i. using Kruskal–Wallis tests with false discovery rate (FDR) correction of *P*-values, followed by Dunn’s post hoc test as appropriate.

To assess differences in microbiota composition (i.e. beta diversity), Jensen–Shannon distances (JSD) were calculated and analysed by permutational analysis of variance (PERMANOVA, 10,000 permutations). For faecal samples, the initial PERMANOVA included the predictors experimental group, day p.i. and their interaction, and was stratified by animal ID to account for repeated sampling. Because all predictors were significant in the initial model, separate PERMANOVAs were subsequently calculated for each infection group versus the control group, and for each sampling day to investigate differences in detail. Likewise, separate PERMANOVAs were calculated for each infection group versus the control group and each day p.i. regarding the ingesta samples.

The relative abundance of microbial phyla and genera was compared among the three groups at each day p.i. using Kruskal–Wallis tests with FDR correction of *P*-values. Dunn’s post hoc tests were conducted if corrected *P*-values were  $\leq 0.05$ . To further analyse differences in the abundance of specific microbial taxa at the species level, DESeq2 differential expression analyses were conducted using the non-rarefied datasets, as DESeq2 performs an internal normalization [32].

### Microbial network analysis

Microbial association networks were constructed based on faecal samples, as the number of only six pigs slaughtered per group at each time point did not allow network construction for ingesta samples. Networks were constructed for days –3, 14, 21 and 35 p.i. via the R package “NetCoMi” [35] using the SPIEC-EASI (Sparse Inverse Covariance Estimation for Ecological Association Inference) association measure, which takes the compositional structure of amplicon-based datasets into account [36]. This was not possible for day 49 due to the smaller sample size per group at this time point. For network construction, OTUs present in less than 20% of all faecal samples were excluded, but no prior rarefaction was performed, as normalization is included in the “spiec.easi” function. Networks were calculated separately for each group and day p.i. The greedy modularity algorithm was used for network clustering [37]. Normalized centrality measures (degree, betweenness, closeness and eigenvector centrality) were calculated. “Hub” OTUs, i.e. influential OTUs in the networks, were defined as having an eigenvector centrality value above the 95% quantile of the empirical eigenvector centrality distribution [35]. Eigenvector centrality is a measure of influence in a network, which takes not only the number of connections of a node (i.e. the degree) into account, but also the degree of its neighbours.

## Results

All control animals remained *Ascaris*-seronegative throughout the study period. Of the 36 animals in the infection groups, five were excluded from the study (two from the single- and three from the trickle-infected group) as they did not seroconvert until slaughter and had no visible worms in the intestines. The average daily weight gain ranged from 0.28 to 0.5 kg for pigs slaughtered on day 21 p.i., from 0.30 to 0.47 kg for those slaughtered on day 35 p.i. and from 0.32 to 0.60 kg for those slaughtered on day 49 p.i. A significant difference in daily weight gain between the experimental groups was observed only on day 49 p.i. (Kruskal–Wallis  $\chi^2=7.5$ ,  $df=2$ ,  $P=0.024$ ), with significantly higher daily weight gain in the trickle-infected than in the single-infected and the control groups (Dunn's all-pairs test, Bonferroni–Holm-adjusted  $P=0.048$  for both comparisons).

### Faecal samples

#### Alpha and beta diversity

The quality-filtered dataset contained 6,879,264 sequences, with a mean of 19,882 sequences per sample (standard deviation [SD]:14,362). After rarefying to 3000 reads per sample, 340 of the original 346 faecal samples were retained, containing 591 unique OTUs (13 phyla, 27 families, 80 genera).

Analysis of alpha diversity indices by GLMMs indicated a significant interaction effect of day p.i. and the *A. suum* infection group regarding the inverse Simpson index, with a significant decline in the single-infected group versus the control group over the course of the infection. In contrast, trickle-infected pigs showed a consistently higher inverse Simpson index during the entire study period (Table 1, Fig. 2). No significant differences regarding the number of observed OTUs, the Chao1 index or the Shannon index were detected (Table 1).

Microbial community composition was significantly affected by day p.i., experimental group and their interaction (PERMANOVA of JSDs, Table 2). Therefore, separate PERMANOVAs were calculated for each infection group versus the control group, and for each sampling day. Results revealed that the overall microbiota composition of the single-infected group differed significantly from the control group on days 14 and 21, whereas the trickle-infected group consistently showed a significantly different microbiota composition from that of the control group, except on day 28 p.i. (Additional file 1).

#### Shifts in taxonomic composition

The dominant phyla in pig faeces were Firmicutes and Bacteroidetes, with mean relative abundance of 52.7% (SD: 15.4%) and 33.1% (SD: 12.1%) per sample, respectively. Regarding relative microbial abundance at the

phylum level, only a few differences were noted between the groups. On day 3 p.i., the trickle-infected group had significantly higher relative abundance of Fibrobacteres than the control group (Kruskal–Wallis  $\chi^2=18.9$ ,  $df=2$ ,  $P_{\text{adj}}=0.008$ /Dunn's test  $P=0.007$ ), while on day 14 p.i., Verrucomicrobia were significantly increased in the trickle-infected group as compared with control animals (Kruskal–Wallis  $\chi^2=17.8$ ,  $df=2$ ,  $P_{\text{adj}}=0.008$ /Dunn's test  $P=0.026$ ). Significant differences at the genus level were mainly observed in the trickle-infected group, with lower *Dorea* but higher *Peptococcus* relative abundance before infection. Thereafter, differences included increased relative abundance of *Bifidobacterium*, *Fibrobacter* and an unclassified Clostridia genus on day 3 p.i., an unclassified Prevotellaceae genus on day 7 p.i., an unclassified Eubacteriaceae and unclassified Verrucomicrobia genus on day 14 p.i. and *Campylobacter* on day 21 p.i. (Table 3). Decreased relative abundance was observed for *Bifidobacterium* on days 28 and 35 p.i. In the single-infected group, *Peptococcus* was increased before infection, and an increase in *Streptococcus* was observed on days 28 and 35 p.i. No significant differences were observed on days 42 and 49 p.i. at the genus level.

DESeq2 analysis indicated that in faecal samples of both infection groups, the number of differentially abundant bacterial species, as compared with the control group, gradually increased during the course of the experiment. At the species level, between 1 (day –3) and 24 (day 49) species were significantly altered in abundance (Benjamini–Hochberg adjusted  $P\leq 0.01$ ) in the single-infected group as compared with the control group, while this number ranged from 6 (day –3) to 27 (day 49) in the trickle-infected group (Fig. 3). In both groups, a prominent increase in an *Acetanaerobacterium* sp. was noted on day 7 p.i. A further similarity was an increase in a *Coproccoccus* sp. on days 21, 28 and 49 p.i. Notably, in the trickle-infected group, several species assigned to the genus *Bifidobacterium* decreased in abundance as from day 21 p.i., while members of the genera *Dorea*, *Coproccoccus*, *Ruminococcus*, *Lactobacillus*, *Paraprevotella* and *Parabacteroides* showed an increase towards the end of the observation period (Fig. 3).


#### Faecal microbial association networks

The filtered dataset used for network analyses contained 375 OTUs, which occurred in at least 20% of faecal samples. Due to the exclusion of some animals, networks were based on 15 to 18 (days –3, 14, 21) or 10 to 12 faecal samples (day 35). Network characteristics are summarized in Table 4, and partial networks (100 most central nodes only) are visualized in Fig. 4. Network density was similar in all three experimental groups from day –3 through day 21 p.i., but decreased in the control

**Table 1** Influence of *A. suum* infection group and day post-infection on faecal microbial alpha diversity

	Observed OTUs (Poisson error/identity link)			Chao1 index (gamma error/identity link)			Shannon index (gamma error/identity link)			Inverse Simpson index (gamma error/log link)						
	Est	SE	<i>z</i>	<i>P</i>	Est	SE	<i>t</i>	<i>P</i>	Est	SE	<i>t</i>	<i>p</i>				
Intercept	115.36	7.51	15.36	<b>&lt; 0.001</b>	122.32	10.05	12.17	<b>&lt; 0.001</b>	3.77	0.11	34.03	<b>&lt; 0.001</b>	3.15	0.10	31.95	<b>&lt; 0.001</b>
Day p.i.	0.10	0.07	1.55	0.122	0.00	0.23	0.01	0.993	-0.01	0.16	-0.08	0.934	0.00	0.00	0.16	0.876
Group: single vs control	1.43	10.96	0.13	0.897	-0.18	14.22	-0.01	0.990	0.29	0.17	1.79	0.073	0.04	0.14	0.27	0.791
Group: trickle vs control	20.47	11.10	1.84	0.065	17.49	14.70	1.19	0.234	0.00	0.00	-0.14	0.888	0.34	0.15	2.25	<b>0.024</b>
Day p.i.*Group (single vs control)	-0.16	0.09	-1.72	0.086	-0.03	0.31	-0.09	0.929	-0.01	0.00	-1.48	0.140	-0.01	0.00	-3.81	<b>&lt; 0.001</b>
Day p.i.*Group (trickle vs control)	0.22	0.11	1.70	0.089	0.35	0.41	0.844	0.399	0.00	0.00	-0.22	0.830	0.00	0.00	-0.84	0.400

GLMMs were carried out to analyse the influence of the *A. suum* infection group (single-infection, trickle-infection, non-infected control) and day post-infection (day p.i.) on different alpha diversity metrics. Significant *P*-values are printed in bold. Est.: estimate; OTUs: operational taxonomic units; p.i.: post-infection; SE: standard error



and single-infected group on day 35 p.i., while network modularity, clustering coefficient and the proportion of positive associations showed the opposite pattern. In the trickle-infected group, only the clustering coefficient and modularity showed an increase on day 35 p.i., while the remaining measures were comparable to the preceding time points. Thus, networks from the control and single-infected groups on day 35 p.i. showed a higher degree of partitioning into groups of well-connected (i.e. associated) OTUs than the trickle-infected group.

In each network, between 17 and 19 OTUs were identified as “hub nodes” based on eigenvector centrality, i.e. OTUs with a high degree of association to other OTUs, which were themselves highly connected. These OTUs can be regarded as influential in the network, affecting the overall OTU composition. Prior to infection (day -3), the set of hub nodes was rather similar among the three experimental groups, with two hub OTUs shared by the single-infected and the control group, four by the trickle-infected and the control group and two between the single- and trickle-infected groups (Additional file 2). Between-group similarity decreased on day 14 p.i., with only two hub nodes each shared between the single-infected and the

trickle-infected and the control group, respectively. On day 21 p.i., the single-infected group shared six hub OTUs with the control group, while the trickle-infected group shared only one hub OTU (assigned to *Olsenella scatoligenes*) with both other groups. On day 35 p.i., both infection groups shared one hub OTU each with the control group.

The temporal stability of hub nodes in the control group was low at the beginning of the study, with only one common hub node identified in the networks from days 14 and 21 p.i., but four hub nodes, mainly from the family Lachnospiraceae, remained consistent from day 21 to day 35 p.i. (Additional file 2). In the infection groups, temporal stability was rather low throughout the course of infection. In the single-infected group, one hub node each remained consistent from days 14 to 21 p.i. and days 21 to 35 p.i. In the trickle-infected group, two hub nodes remained consistent from day -3 to day 14 p.i. and one from days 14 to 21 p.i., but none from days 21 to 35 p.i.

In terms of hub OTU taxonomy, the most well-represented families in the control group and single-infected group were Lachnospiraceae and/or Prevotellaceae, except for day -3 and day 14 p.i., when the set of hub nodes in the single-infected group was taxonomically

**Table 2** PERMANOVA of porcine faecal microbiota composition

Term	df	SS	MS	F	R <sup>2</sup>	P
Group	2	2.39	1.19	8.00	0.04	<b>&lt;0.001</b>
Day p.i.	1	1.97	1.97	13.22	0.04	<b>&lt;0.001</b>
Group*Day p.i.	2	1.30	0.65	4.36	0.02	<b>&lt;0.001</b>
Residuals	334	49.76	0.15		0.90	
Total	339	55.41			1.00	

PERMANOVA was carried out to test the effect of the experimental *A. suum* group (single-infection, trickle-infection, non-infected control) and day post-infection (p.i.) on microbiota composition in porcine faecal samples, based on Jensen–Shannon distances. Significant *P*-values are printed in bold. *df* degrees of freedom; MS: mean squares; p.i.: post-infection; SS: sum of squares

**Table 3** Genus-level differences in the faecal microbiota between *A. suum* single-infected (SI), trickle-infected (TI) and non-infected (C) pigs

	Mean relative abundance ± SD			Kruskal–Wallis test			Dunn's test	
	SI	TI	C	X <sup>2</sup>	df	P <sub>FDR-adj</sub>	P (SI vs. C)	P (TI vs. C)
Day -3								
<i>Dorea</i> (Lachnospiraceae)	1.00 ± 0.67	0.20 ± 0.40	1.42 ± 1.41	16.65	2	0.033	0.901	<b>&lt;0.001</b>
<i>Peptococcus</i> (Peptococcaceae 1)	0.21 ± 0.18	0.34 ± 0.32	0.06 ± 0.12	15.37	2	0.033	<b>0.010</b>	<b>&lt;0.001</b>
Day 3 p.i.								
<i>Bifidobacterium</i> (Bifidobacteriaceae)	0.00 ± 0.00	1.29 ± 2.99	0.00 ± 0.00	15.10	2	0.033	1.000	<b>0.001</b>
<i>Fibrobacter</i> (Fibrobacteraceae)	0.01 ± 0.03	0.76 ± 1.31	0.21 ± 0.57	18.89	2	0.033	0.232	<b>0.007</b>
Unclassified Clostridia	0.06 ± 0.12	1.31 ± 0.25	0.04 ± 0.13	15.80	2	0.033	0.595	<b>&lt;0.001</b>
Day 7 p.i.								
Unclassified Prevotellaceae	0.25 ± 0.48	1.30 ± 1.02	0.31 ± 0.42	16.50	2	0.033	0.776	<b>0.002</b>
Day 14 p.i.								
Unclassified Eubacteriaceae	0.26 ± 0.43	0.98 ± 0.75	0.16 ± 0.19	15.6	2	0.033	0.999	<b>&lt;0.001</b>
Unclassified Verrucomicrobia	0.04 ± 0.12	1.81 ± 2.66	0.41 ± 0.80	16.20	2	0.033	0.169	<b>0.024</b>
Day 21 p.i.								
<i>Campylobacter</i> (Campylobacteraceae)	0.25 ± 0.79	1.24 ± 1.22	0.1 ± 0.24	15.6	2	0.033	0.963	<b>&lt;0.001</b>
Day 28 p.i.								
<i>Bifidobacterium</i> (Bifidobacteriaceae)	3.25 ± 7.49	0.00 ± 0.00	1.76 ± 2.00	14.7	2	0.035	0.503	<b>0.013</b>
<i>Streptococcus</i> (Streptococcaceae)	1.29 ± 1.54	0.06 ± 0.18	0.01 ± 0.02	16.3	2	0.033	<b>&lt;0.001</b>	0.995
Day 35 p.i.								
<i>Bifidobacterium</i> (Bifidobacteriaceae)	3.24 ± 4.31	0.00 ± 0.00	3.14 ± 4.04	15.8	2	0.033	0.588	<b>0.007</b>
<i>Streptococcus</i> (Streptococcaceae)	2.77 ± 4.15	0.27 ± 0.59	0.07 ± 0.22	15.0	2	0.033	<b>&lt;0.001</b>	0.831

Only genera that were significantly different between groups (*P*-values printed in bold) are shown. No significant differences were noted on days 42 and 49 p.i. C: control; *df*: degrees of freedom; FDR: false discovery rate; p.i.: post-infection; TI: trickle infection; SD: standard deviation; SI: single infection


rather diverse (Fig. 4, Additional file 2). Most hub nodes of the trickle-infected group were assigned to Prevotellaceae and Ruminococcaceae prior to infection, while on day 14 p.i., Prevotellaceae, Lachnospiraceae and Ruminococcaceae were almost equally represented. Prevotellaceae and Veillonellaceae predominated on day 21 p.i., followed by Lachnospiraceae and Ruminococcaceae on day 35 p.i. Therefore, the trickle-infected group displayed the lowest temporal stability in terms of hub node taxonomy at the family level.

## Intestinal content

### Alpha and beta diversity

Several ingesta samples, especially from the jejunum, had to be excluded due to insufficient DNA for sequencing or low sequencing output. Consequently, 106 of 138 small intestine samples and 96 of 98 large intestine samples were retained after quality filtering, yielding a total of 3,419,792 (mean 32,262 ± 15,440 per sample) and 2,852,943 (mean 29,718 ± 14,289) sequences, respectively. Rarefaction to 8000 reads/sample resulted in the removal of four further jejunum samples. In the small





intestine, the rarefied sequences were assigned to 255 unique OTUs (8 phyla, 44 families, 84 genera), whereas 469 unique OTUs (9 phyla, 32 families, 71 genera) were present in the large intestine samples.

A wide range in alpha diversity among ingesta samples from the jejunum and a general decrease in diversity from the proximal to the distal small intestine were noted (Fig. 5). However, comparability between groups was limited due to the low number of successfully sequenced jejunum samples, and no significant differences in alpha diversity measures were detected in the small intestine (Kruskal–Wallis tests, FDR-adjusted  $P > 0.05$ ). Regarding the large intestine, a significant reduction in observed OTUs, Shannon index and inverse Simpson index were noted on day 35 p.i. in the caecum of the single-infected group relative to the other groups (Kruskal–Wallis  $\chi^2 = 11.58/10.59/10.59$ ,  $df = 2$ , FDR-adjusted  $P = 0.003/0.005/0.005$ , Dunn’s test

$P = 0.002/0.012/0.012$ ). In fact, only 24.0 OTUs (SD: 6.4) were detected on average in the caecum samples of the single-infected group on day 35, as opposed to 252.2 (SD: 56.1) in the control and 196.2 (SD: 58.8) in the trickle-infected group. No significant differences in alpha diversity were detected regarding the colon.

Because of the low number of jejunum ingesta samples, beta diversity analyses were not carried out for this sample subset. Regarding the other intestinal compartments, initial PERMANOVA indicated a significant main effect of group for the ileum and caecum, as well as a significant effect of day p.i. for the caecum and colon and a significant interaction effect of group and day p.i. in the ileum (Table 5). Conducting PERMANOVAs for each of the infection groups versus the control group and for each day p.i. separately revealed significant differences in microbiota composition between the

**Table 4** Characteristics of microbial association networks determined from faecal samples of *A. suum*-infected and non-infected pigs

	Day -3				Day 14 p.i.				Day 21 p.i.				Day 35 p.i.			
	C	SI	TI	TI	C	SI	TI	TI	C	SI	TI	TI	C	SI	TI	TI
Number of nodes	363	359	370	366	369	350	366	366	365	359	364	364	346	324	350	
Edge density	0.022	0.026	0.027	0.023	0.028	0.023	0.023	0.023	0.025	0.021	0.024	0.024	0.017	0.016	0.021	
Positive edges (%)	82.48	76.193	75.73	77.228	76.326	80.675	77.228	77.228	80.427	82.619	77.997	77.997	86.944	89.260	76.471	
Clustering coefficient	0.089	0.123	0.119	0.104	0.093	0.121	0.104	0.104	0.113	0.118	0.109	0.109	0.184	0.230	0.184	
Modularity	0.38	0.355	0.340	0.378	0.326	0.394	0.378	0.378	0.368	0.411	0.364	0.364	0.527	0.568	0.442	
No. of modules (greedy algorithm)	10	11	8	14	8	11	14	14	10	11	9	9	14	16	15	
Mean normalized eigenvector centrality ( $\pm$ SD)	0.404 ( $\pm$ 0.213)	0.358 ( $\pm$ 0.219)	0.348 ( $\pm$ 0.227)	0.352 ( $\pm$ 0.215)	0.468 ( $\pm$ 0.216)	0.305 ( $\pm$ 0.215)	0.352 ( $\pm$ 0.219)	0.352 ( $\pm$ 0.219)	0.318 ( $\pm$ 0.210)	0.276 ( $\pm$ 0.220)	0.412 ( $\pm$ 0.227)	0.412 ( $\pm$ 0.227)	0.214 ( $\pm$ 0.196)	0.209 ( $\pm$ 0.210)	0.273 ( $\pm$ 0.198)	

C: control; SI: single infection; p.i.: post-infection; SD: standard deviation; TI: trickle infection

single-infected group and the control group on days 21 and 35 p.i. in the ileum and caecum, whereas no significant differences were found on day 49 p.i. Regarding the trickle-infected group, no significant differences in overall microbiota composition relative to the control group were found in any intestinal compartment (Fig. 6; Additional file 3).

#### Shifts in taxonomic composition

In the small intestine, Firmicutes was the dominant phylum, with mean relative abundance of 90.56% per sample (SD: 11.7%). In the large intestine, Firmicutes showed a mean relative abundance of 74.6% (SD: 12.3%), while on average 19.8% (SD: 10.9%) of the sequences were assigned to Bacteroidetes.

The 12 most abundant bacterial genera in the ileum, caecum and colon of pigs slaughtered at days 21 and 35 p.i. are shown in Fig. 7. The microbiota of the ileum was dominated by the genera *Lactobacillus*, *Clostridium* sensu stricto (s.s.) and *Terrisporobacter*, while the caecum and colon were dominated by *Lactobacillus*, *Prevotella* and *Clostridium* s.s., among others. At the phylum level, no statistically significant differences in relative abundance were detected among the ingesta samples of the different experimental groups. At the genus level, however, significant differences were apparent in caecum samples from days 21 and 35 p.i., but not day 49 p.i. On day 21 p.i., seven and eight genera showed a higher relative abundance in the single- and the trickle-infected groups, respectively, than in the control group, with an increase in *Megasphaera*, *Blautia*, *Fusicatenibacter* and *Holdemanella* in both infection groups (Table 6). Furthermore, *Romboutsia* was decreased in the single-infected group as compared with the control group. On day 35 p.i., 26 bacterial genera were significantly reduced in the caecum of the single-infected group (Table 6), in line with the significantly reduced alpha diversity. Four of these genera (*Fusicatenibacter*, *Ruminococcus* 2, *Collinsella* and *Holdemanella*) also showed significantly lower relative abundance in the trickle-infected group at this time point.

At the species level, DeSeq2 analysis indicated significant changes in bacterial abundance in the ileum, caecum and colon of both infection groups (Additional file 4). The prevailing pattern across all three intestinal compartments was that of a lower abundance of many taxa relative to the control group, except for the ileum of

the single-infected group on days 35 and 49 p.i. Several species in ileal ingesta showed increased abundance on day 35 p.i. in this group, including *Bifidobacterium thermophilum*, *Eubacterium coprostanoligenes*, *Lactococcus raffinolactis* and unknown species of *Oscillibacter*, *Veillonella*, *Streptococcus*, Ruminococcaceae, Porphyromonadaceae, Demequinaceae and *Lachnospiracea* incertae sedis. *Bifidobacterium thermophilum* and the unknown *Streptococcus* sp. were also elevated in the trickle-infected group on day 35 p.i. In the single-infected group, *Eubacterium coprostanoligenes* and the unknown species of *Veillonella*, *Oscillibacter*, Ruminococcaceae and Porphyromonadaceae remained elevated on day 49 p.i.

In the caecum of the single-infected group, only a few bacterial taxa were significantly altered in abundance on days 21 and 49 p.i., while 74 bacterial taxa were decreased in abundance and only two (*Uruburuella suis* and an unknown *Streptococcus* sp.) were increased on day 35 p.i. *Uruburuella suis* remained significantly increased in this group on day 49 p.i., while *Lactobacillus equicursoris* was the only taxon to remain decreased from day 35 to day 49 p.i. (Additional file 4).


In contrast, only minor changes were observed at the species level in the caecum of the trickle-infected group and in the colon of both infection groups. However, considering all studied intestinal compartments (ileum, caecum, colon) and all time points, the following parallels seem noteworthy: On day 35 p.i., an unknown *Streptococcus* sp. was elevated in all three compartments of both infection groups, except in the caecum of the trickle-infected group. In addition, a decrease in *L. equicursoris* was observed on day 49 p.i. in all compartments of both infection groups. The decrease in various *Bifidobacterium* spp. on day 49 p.i. was a common pattern in all compartments of the trickle-infected group, as well as in faecal samples (see section above).

#### Discussion


The present study examined faecal and ingesta samples from different intestinal segments to characterize changes in the intestinal microbiota of pigs during the first 7 weeks of *A. suum* infection. Unfortunately, the sequencing success of samples from the jejunum, where *A. suum* resides, was low, limiting the possibility of making meaningful comparisons between experimental groups for this intestinal compartment. Bacterial abundance is rather low in the small intestine [38], and low

(See figure on next page.)

**Fig. 4** Association networks of the faecal microbiota of *A. suum* single-infected and trickle-infected pigs on day -3 (prior to infection) and on days 14, 21 and 35 p.i., compared with a non-infected control group. Only the top 100 nodes (i.e. OTUs) with the largest eigenvector centrality and only positive associations are shown. Node colour indicates taxonomy at the family level, while node size is proportional to eigenvector centrality. Hub nodes (based on eigenvector centrality) are highlighted by a bold rim



**Fig. 4** (See legend on previous page.)




**Fig. 5** Microbial diversity (observed OTUs, Shannon index and inverse Simpson index) in the small intestine (A) and large intestine (B) of *A. suum*-infected and non-infected control pigs. Asterisks indicate statistically significant differences (Kruskal–Wallis test, FDR-corrected  $P < 0.05$ )

**Table 5** PERMANOVAs of microbiota composition in ingesta samples from the ileum, caecum and colon

Compartment	Term	df	SS	MS	F-value	R <sup>2</sup>	P-value
Ileum	Group	2	0.27	0.13	3.31	0.12	<b>0.037</b>
	Day p.i.	1	0.06	0.07	1.64	0.03	0.205
	Group*Day p.i.	2	0.24	0.12	2.99	0.11	<b>0.046</b>
	Residuals	40	1.60	0.04	-	0.74	-
	Total	45	2.17	-	-	1.00	-
Caecum	Group	2	0.30	0.15	2.01	0.08	<b>0.028</b>
	Day p.i.	1	0.37	0.37	4.97	0.09	<b>&lt;0.001</b>
	Group*Day p.i.	2	0.13	0.07	0.89	0.03	0.547
	Residuals	42	3.10	0.07	-	0.80	-
	Total	47	3.90	-	-	1.00	-
Colon	Group	2	0.17	0.08	1.67	0.07	0.078
	Day p.i.	1	0.19	0.19	3.84	0.08	<b>0.001</b>
	Group*Day p.i.	2	0.06	0.03	0.61	0.03	0.846
	Residuals	42	2.04	0.05	-	0.84	-
	Total	47	2.45	-	-	1.00	-

PERMANOVAs testing the effect of experimental group and day p.i. on the microbial composition were based on OTU-level Jensen–Shannon distance values. Significant  $P$ -values are printed in bold. df: degrees of freedom; MS: mean squares; p.i.: post-infection; SS: sum of squares




sequencing success has also been a problem in previous studies [e.g. 39, 40].

#### Alpha diversity was differentially affected by *A. suum* single and trickle infection

The current study used young pigs of a few weeks to approximately 3 months of age (to mimic infections in fattening farms) whose intestinal microbiota is still developing. Previous studies have reported both an increase [41] and a decrease [42] in microbial alpha diversity in healthy pigs during this time span. Here, a marginal increase in observed OTUs and inverse Simpson index was observed over time in the non-infected control group. In contrast, a significant decrease in alpha diversity, more precisely in the inverse Simpson index, which takes microbial abundance into account, occurred in faecal samples of pigs single-infected with 10,000 *A. suum* eggs. Further, a dramatic decrease in all examined alpha diversity measures was observed in the caecum of the single-infected group at day 35 p.i. A decrease in microbial diversity was also observed in the colon contents of pigs 54 days after infection with 300 *A. suum* eggs each on three consecutive days, independent of the intestinal

worm burden [19]. In the current study, however, colonic microbial diversity was not significantly affected. The apparent discrepancy between alpha diversity analyses of colon and faecal samples may be due to the smaller sample sizes regarding ingesta, reducing statistical power. Furthermore, alpha diversity in the caecum was restored by day 49 p.i. *Ascaris suum* excretory–secretory (ES) products include a suite of antimicrobial factors, possibly to actively manipulate its microbial environment. Interestingly, the spectrum of antimicrobial products changes throughout the parasite's life cycle, with a higher diversity of antimicrobial components detected in adult parasites compared with L4 [15]. Moulting of L4 occurs around days 21–29 p.i.; thus, the sharp drop in alpha diversity observed at day 35 p.i. in the caecum of single-infected pigs may be due to the recent development of L4 to the pre-adult stage. However, it remains speculative why this sharp reduction in alpha diversity was not observed in the other intestinal compartments. Most of the significantly reduced genera in the caecum belonged to the families Lachnospiraceae and Ruminococcaceae, which are generally more abundant in the porcine large intestine than in the small intestine [43, 44]. It is conceivable that the *A. suum* ES antigen preferentially affects these taxa, which would explain why no alpha diversity reduction was seen in the ileum. Furthermore, the amount of antimicrobial products reaching the colon may be too small or degenerated to cause comparable effects. In addition, it remains an open question which mechanisms lead to diversity restoration between days 35 and 49 p.i. The so-called self-cure reaction, which causes partial elimination of the worm burden, takes place earlier, approximately between days 17 and 21 following a single experimental infection [5], and is thus unlikely to be responsible for the restoration of alpha diversity in the caecum.

In contrast to the single-infected group, no significant alpha diversity changes were noted in the different intestinal compartments of the trickle-infected group, while alpha diversity in faecal samples was higher than in the control group throughout the course of the study, including before infection. Previous studies have reported mixed results regarding microbial alpha diversity during helminth infections. Consequently, a recent meta-analysis identified no significant differences between helminth-infected and non-infected hosts, apart from rodents, which showed decreased microbial alpha diversity upon helminth infection [45]. In contrast, increased alpha diversity has been reported in humans naturally infected with soil-transmitted helminths, including *Ascaris* spp., in different parts of the world [e.g. 46–48]. These ambiguous results may be driven by differential effects of experimental, high-dose infections versus



**Fig. 7** Genus-level taxonomic composition of the microbiota in the ileum, caecum and colon of *A. suum*-infected vs non-infected control pigs on days 21 and 35 p.i.

continuous trickle infections, aside from species-specific differences between host–parasite systems. In consequence, care should be taken when extrapolating results from experimental infection studies using a single infection dose to natural scenarios.

***Ascaris suum* predominantly affected the microbiota composition of the caecum**

The microbiota composition of the caecum was predominantly affected by *A. suum* infection, as indicated by significant differences in the relative abundance of

**Table 6** Genus-level differences in the caecal microbiota between *A. suum* single-infected (SI), trickle-infected (TI) and non-infected (C) pigs

	Mean relative abundance (%) ± SD			Kruskal–Wallis test			Dunn's test	
	SI	TI	C	$\chi^2$	df	$P_{\text{FDR-adj}}$	$P$ (SI vs C)	$P$ (TI vs C)
Day 21 p.i.								
<i>Gemmiger</i> (Ruminococcaceae)	4.65 ± 1.69	7.53 ± 1.3	0.57 ± 1.25	11.24	2	<b>0.038</b>	0.138	<b>0.002</b>
<i>Prevotella</i> (Prevotellaceae)	26.19 ± 14.44	10.66 ± 7.28	3.52 ± 5.58	9.24	2	<b>0.049</b>	<b>0.005</b>	0.229
<i>Megasphaera</i> (Veillonellaceae)	2.93 ± 2.22	2.24 ± 1.37	0.11 ± 0.28	10.52	2	<b>0.038</b>	<b>0.012</b>	<b>0.012</b>
<i>Faecalibacterium</i> (Ruminococcaceae)	3.02 ± 0.84	6.02 ± 1.68	0.57 ± 0.89	11.78	2	<b>0.038</b>	0.101	<b>0.001</b>
<i>Romboutsia</i> (Peptostreptococcaceae)	0.17 ± 0.11	0.41 ± 0.14	7.37 ± 10.76	11.13	2	<b>0.038</b>	<b>0.002</b>	0.132
<i>Blautia</i> (Lachnospiraceae)	2.39 ± 0.9	2.58 ± 0.78	0.3 ± 0.56	9.57	2	<b>0.047</b>	<b>0.026</b>	<b>0.012</b>
<i>Fusicatenibacter</i> (Lachnospiraceae)	0.27 ± 0.12	0.27 ± 0.14	0.03 ± 0.07	9.41	2	<b>0.048</b>	<b>0.017</b>	<b>0.020</b>
<i>Collinsella</i> (Coriobacteriaceae)	0.54 ± 0.38	0.2 ± 0.15	0.03 ± 0.06	10.37	2	<b>0.038</b>	<b>0.003</b>	0.129
<i>Holdemanella</i> (Erysipelotrichaceae)	0.22 ± 0.08	0.35 ± 0.23	0.02 ± 0.05	10.33	2	<b>0.038</b>	<b>0.035</b>	<b>0.006</b>
<i>Olsenella</i> (Coriobacteriaceae)	0.14 ± 0.08	0.23 ± 0.13	0.02 ± 0.04	9.40	2	<b>0.048</b>	0.069	<b>0.007</b>
<i>Peptococcus</i> (Peptococcaceae 1)	0.08 ± 0.11	0.07 ± 0.02	0.00 ± 0.00	9.38	2	<b>0.048</b>	0.074	<b>0.007</b>
<i>Acidaminococcus</i> (Acidaminococcaceae)	0.22 ± 0.31	0.00 ± 0.00	0 ± 0.01	11.89	2	<b>0.038</b>	<b>0.007</b>	0.893
Day 35 p.i.								
<i>Gemmiger</i> (Ruminococcaceae)	0.00 ± 0.00	2.60 ± 1.1	4.13 ± 1.57	11.95	2	<b>0.038</b>	<b>0.002</b>	0.575
<i>Prevotella</i> (Prevotellaceae)	0.06 ± 0.16	18.06 ± 4.57	9.69 ± 7.82	11.93	2	<b>0.038</b>	<b>0.041</b>	0.491
<i>Megasphaera</i> (Veillonellaceae)	0.00 ± 0.00	0.32 ± 0.58	3.90 ± 3.73	10.50	2	<b>0.038</b>	<b>0.003</b>	0.074
<i>Roseburia</i> (Lachnospiraceae)	0.01 ± 0.02	3.59 ± 1.51	3.02 ± 2.05	10.95	2	<b>0.038</b>	<b>0.013</b>	0.970
Unknown Lachnospiraceae	0.00 ± 0.00	4.25 ± 2.26	5.41 ± 1.29	11.28	2	<b>0.038</b>	<b>0.005</b>	0.918
<i>Butyricicoccus</i> (Clostridiaceae)	0.00 ± 0.00	0.30 ± 0.36	2.77 ± 1.26	12.57	2	<b>0.038</b>	<b>&lt;0.001</b>	0.051
<i>Faecalibacterium</i> (Ruminococcaceae)	0.00 ± 0.00	0.63 ± 0.75	3.21 ± 1.61	12.60	2	<b>0.038</b>	<b>&lt;0.001</b>	0.127
<i>Lachnospiraceae</i> inc. sed. (Lachnospiraceae)	0.01 ± 0.02	1.80 ± 0.61	2.07 ± 1.53	10.95	2	<b>0.038</b>	<b>0.013</b>	0.970
<i>Phascolarctobacterium</i> (Acidaminococcaceae)	0.00 ± 0.00	1.36 ± 0.92	0.94 ± 0.78	11.28	2	<b>0.038</b>	<b>0.014</b>	0.918
<i>Blautia</i> (Lachnospiraceae)	0.00 ± 0.00	1.01 ± 0.62	1.76 ± 0.44	11.71	2	<b>0.038</b>	<b>0.001</b>	0.217
<i>Coprococcus</i> (Lachnospiraceae)	0.00 ± 0.00	1.03 ± 0.63	1.88 ± 1.22	10.58	2	<b>0.038</b>	<b>0.003</b>	0.430
<i>Dorea</i> (Lachnospiraceae)	0.00 ± 0.00	0.13 ± 0.1	0.56 ± 0.07	13.16	2	<b>0.038</b>	<b>&lt;0.001</b>	0.093
Unknown Erysipelotrichales	0.00 ± 0.00	1.16 ± 1.46	0.6 ± 0.51	13.14	2	<b>0.038</b>	<b>&lt;0.001</b>	0.094
Unknown Bacteroidia	0.00 ± 0.00	1.16 ± 1.46	0.60 ± 0.51	9.35	2	<b>0.048</b>	<b>0.017</b>	1.000
Unknown Clostridiales	0.03 ± 0.06	1.70 ± 2.1	1.68 ± 0.92	9.72	2	<b>0.046</b>	<b>0.007</b>	0.769
<i>Oribacterium</i> (Lachnospiraceae)	0.00 ± 0.00	0.14 ± 0.19	0.5 ± 0.34	9.92	2	<b>0.043</b>	<b>0.003</b>	0.102
<i>Fusicatenibacter</i> (Lachnospiraceae)	0.00 ± 0.00	0.05 ± 0.11	0.26 ± 0.22	9.95	2	<b>0.043</b>	<b>0.004</b>	<b>0.049</b>
<i>Ruminococcus</i> 2 (Ruminococcaceae)	0.00 ± 0.00	0.04 ± 0.08	0.38 ± 0.25	12.19	2	<b>0.038</b>	<b>0.002</b>	<b>0.014</b>
<i>Collinsella</i> (Coriobacteriaceae)	0.00 ± 0.00	0.01 ± 0.02	0.29 ± 0.16	13.03	2	<b>0.038</b>	<b>0.001</b>	<b>0.009</b>
Unknown Prevotellaceae	0.00 ± 0.00	0.69 ± 0.5	0.37 ± 0.24	11.48	2	<b>0.038</b>	<b>0.019</b>	0.805
<i>Holdemanella</i> (Erysipelotrichaceae)	0.00 ± 0.00	0.00 ± 0.00	0.28 ± 0.15	14.41	2	<b>0.038</b>	<b>0.001</b>	<b>0.002</b>
Unknown Firmicutes	0.00 ± 0.00	0.08 ± 0.07	0.26 ± 0.22	10.58	2	<b>0.038</b>	<b>0.003</b>	0.430
<i>Selenomonas</i> (Selenomonadaceae)	0.00 ± 0.00	0.41 ± 0.55	0.23 ± 0.18	11.30	2	<b>0.038</b>	<b>0.014</b>	0.918
<i>Anaerovibrio</i> (Veillonellaceae)	0.00 ± 0.00	0.48 ± 0.28	0.22 ± 0.29	11.95	2	<b>0.038</b>	<b>0.032</b>	0.575
<i>Paraeggerthella</i> (Coriobacteriaceae)	0.00 ± 0.00	0.03 ± 0.04	0.11 ± 0.05	10.82	2	<b>0.038</b>	<b>0.002</b>	0.062
Unknown Coriobacteriaceae	0.00 ± 0.00	0.13 ± 0.13	0.28 ± 0.18	11.10	2	<b>0.038</b>	<b>0.002</b>	0.312

Only genera that were significantly different between groups ( $P$ -values printed in bold) are listed. No significant differences were noted on day 49 p.i. C: control;  $df$ : degrees of freedom; FDR, false discovery rate; p.i., post-infection; TI: trickle infection; SD: standard deviation; SI: single infection

various microbial genera. On day 21 p.i., an increase in several short-chain fatty acid (SCFA)-producing genera was noted in both infection groups in this intestinal compartment. The opposite was observed on day

35 p.i., with a significantly reduced abundance of several SCFA-producing genera in both infection groups. Specifically, *Blautia*, *Fusicatenibacter*, *Holdemanella* and *Megasphaera* were significantly increased in both



infection groups on day 21 p.i., while the trickle-infected group additionally displayed an increase in *Faecalibacterium*. *Fusicatenibacter*, *Ruminococcus* 2, *Collinsella* and *Holdemanella* were reduced in both groups on day 35 p.i. Increased production of SCFAs such as butyrate and propionate has been observed in a variety of intestinal helminth infections and is regarded as beneficial for both the parasites and the host, attenuating inflammation and serving as an energy source for intestinal epithelial cells [2, 49, 50]. The shift from an increase to a decrease in SCFA-producing taxa may again be related to the moulting of L4 to adult parasites around days 21–29 p.i., and the accompanying shift in the spectrum of antimicrobial products [15]. SCFA production may be especially beneficial for the parasite during its L4 stage, when it is vulnerable to the “self-cure reaction”, which is characterized by an increase in mucosal eosinophils, macrophages, T-cells and, consequently, smooth muscle contractility [6]. SCFAs are known to modulate T-cell function [51], and may attenuate the self-cure reaction at a critical time point for the parasite, while older parasites are able to counteract the increased peristaltic movements due to their large size and strong musculature [6]. In addition, it has been suggested that adult *A. suum* themselves may induce a strong downregulation of inflammatory pathways in the porcine intestinal mucosa during later stages of infection [52].

In line with the reduction in alpha diversity on day 35 p.i. in the caecum of the single-infected group discussed above, various other taxa were affected. Overall, 26 bacterial genera showed a significantly reduced abundance, including for example *Prevotella*, *Roseburia* and *Gemmiger*, in addition to the above-mentioned genera. *Prevotella* is considered a keystone taxon in the large intestine of pigs, with a profound influence on the community structure and functioning of the microbiota [53]. Indeed, OTUs assigned to *Prevotella* were frequently identified as “hub nodes” in the faecal microbiota networks of the present study. In the caecum on day 35 p.i., the relative abundance of this genus amounted to 9.7% and 18.1% on average in the control group and the trickle-infected group, respectively, but only 0.1% in the single-infected group. The genus *Prevotella* has been positively associated with production parameters such as feed intake, feed conversion efficiency and weight gain, possibly due to their ability to ferment complex carbohydrates [54]. Thus, an *A. suum*-related depletion of *Prevotella* in the caecum might contribute to the lower feed intake and weight gain observed in some studies on *A. suum*-infected pigs [e.g. 55, 56]. However, in a previous study on *A. suum*-infected pigs, Wang et al. [19]

rather observed an increase in *Prevotella* in colon contents at day 54 p.i., while the significant genus-level reduction in *Prevotella* in the current study occurred only transiently in the caecum, and not in the colon or the faeces, and was limited to the single-infected group. In terms of weight gain, we did not observe significant differences between the single-infected and the control group at any time point, whereas higher weight gain was indeed noted in the trickle-infected pigs, in line with the rather high *Prevotella* abundance in this group on day 35 p.i. However, the difference in weight gain was statistically significant only on day 49 p.i., when no significant differences in *Prevotella* abundance were evident, possibly because of the transient nature of the microbial alterations.

#### ***Ascaris suum* infection modulated bacterial taxa with probiotic properties**

As a result of the significant reduction in various taxa, the caecal microbiota of single-infected pigs was mainly characterized by members of the genus *Lactobacillus* on day 35 p.i., although the increase in relative abundance of *Lactobacillus* was not statistically significant. Since next-generation sequencing data only allow the assessment of relative and not absolute abundance, it remains unclear whether *Lactobacillus* populations had indeed expanded, or whether they remained constant and overall absolute microbial abundance was reduced. Interestingly, Andersen-Civil et al. [18] observed a similar expansion of the genus *Lactobacillus* in the jejunum and colon of *A. suum*-infected pigs at day 14 p.i. Since a decrease in lactase activity has been observed during *A. suum* infection [57], and lactose intolerance has also been reported in *A. lumbricoides*-infected children [58], higher lactose levels in the intestinal contents of infected pigs might boost lactic acid bacteria such as *Lactobacillus*. Alternatively, it has been suggested that increased mucus secretion during helminth infection may favour lactobacilli [18]. An increase in *Lactobacillus* has also been repeatedly reported in the gut microbiota of helminth-infected rodents [45] and in *T. suis*-infected pigs [59]. Further, it has been suggested that *Lactobacillus* spp. might confer a survival advantage to *Heligmosomoides polygyrus* in mice [60] and *Trichuris* spp. in pigs [59]. Like SCFAs, lactate may downregulate proinflammatory responses in the gut [61]. However, on the species level, no clear picture regarding *Lactobacillus* was observed, as an increase in *L. salivarius* and *L. pontis* occurred in faecal samples at some time points as from day 21 p.i., whereas in ingesta samples rather a decrease in certain *Lactobacillus* spp. was observed. Specifically, *L. equicursoris* was decreased in the ileum, caecum and colon on day 35 p.i. (single-infected group only) and day 49 p.i. (both infection

groups), and *L. mucosae* in the ileum on days 35 and 49 p.i. (trickle-infected group only).

One striking alteration observed in the trickle-infected group was the reduction in several *Bifidobacterium* spp., including *B. boum* and *B. thermophilum*, in faecal samples (day 28 p.i. onwards) and in ingesta samples from the ileum, caecum and colon (day 49 p.i.). In a recent meta-analysis, a reduction in the genus *Bifidobacterium* was identified as a common feature across different helminth–host systems [45]. Like *Lactobacillus* spp., members of the genus *Bifidobacterium* are considered probiotic, beneficial bacteria [62]. Recently, *Bifidobacterium animalis* ssp. *lactis* was shown to exert a protective effect during *A. suum* infection, attenuating the parasite's effect on glucose transport as measured in Ussing chambers and modulating the immune response towards increased anti-*A. suum* antibody production [12]. Along these lines, feeding of a fructan-rich chicory-based diet to piglets decreased *A. suum* burden, while increasing the abundance of *Bifidobacterium* spp. in the ileum [63]. Therefore, reducing *Bifidobacterium* populations might be considered a survival strategy for the parasite.

#### ***Ascaris suum* affected temporal stability of faecal microbial association networks**

Faecal microbial association networks showed a low degree of temporal stability in the first 3 weeks of the study in all three experimental groups, in line with the dynamic nature of the gut microbiota in growing pigs [41]. However, in the control group and the single-infected group, a tendency towards stabilization of the networks was observed between days 21 and 35 p.i., indicated by a decrease in network density and a higher degree of partitioning into groups of well-connected OTUs. Furthermore, the identity of several influential OTUs (hub nodes) remained consistent in the control group from day 21 to 35. The control group and the single-infected group also exhibited a higher degree of similarity in terms of hub node identity (shared hub nodes) and family membership of hub nodes, especially on days 14 and 21 p.i. In contrast, the trickle-infected group shared fewer hub OTUs with the control group and displayed little temporal stability in terms of hub node taxonomy over time. In general, most hub nodes in the microbial networks from the current study were members of Lachnospiraceae, Prevotellaceae and Ruminococcaceae across groups and time points, in line with previous findings in young pigs [43]. However, Lachnospiraceae and Prevotellaceae clearly dominated in the control and single-infected group, especially from day 21 p.i. onwards, while in the trickle-infected group, the

pattern was more variable over time, with Ruminococcaceae or Veillonellaceae sometimes dominating over either of the aforementioned families, and a larger proportion of varying other families contributing to the set of hub nodes. Therefore, trickle infection with *A. suum* seemed to prevent the establishment of stable microbial interaction networks until day 35 p.i. Due to sample size considerations, networks were not calculated for days 42 and 49 p.i. or for the different intestinal compartments. Therefore, further studies are required to assess whether stabilization occurs later during infection, or whether *A. suum* continues to perturb the porcine intestinal microbiota during its patent phase after trickle infection.

#### **Conclusions**

The present study revealed marked differences regarding changes in the porcine intestinal microbiota induced by a single versus trickle infection with *A. suum*. The single infection caused a remarkable, but transient, decrease in microbial diversity in the caecum. Although this was not observed following trickle infection, an increase in SCFA-producing genera in the caecum on day 21 p.i., which shifted to a decrease on day 35 p.i., was a common finding in both groups, possibly related to changes in ES products following the parasite's final moult. Moreover, trickle infection, but not single infection, prevented the stabilization of microbial interaction networks. The differences observed between the groups may be due to a more heterogeneous worm population in the trickle-infected group, as larvae continued to arrive in the small intestine over a longer time span. Furthermore, an inverse relationship between infection dose and successful establishment of worms in the intestine has been reported [64], but this view was later challenged [5]. The high degree of overdispersion observed regarding *A. suum* burden in experimental and natural infections may rather be due to an individual predisposition [65]. It was not possible to assess the effect of individual worm burden on the results in the current study, as the intestines were subjected to measurement of electrogenic nutrient transport immediately after slaughter, precluding intestinal washing to recover all parasites [13]. However, the study of Wang et al. [19] showed that the effect of *A. suum* on microbial diversity in the colon was independent of worm burden. Nevertheless, further studies should take this factor into account, especially when examining chronically infected pigs. The current study filled the gap between previous observations made on pigs infected for 14 days and 54 days with *A. suum* [17, 19]; however, studies on chronically infected pigs beyond this time point are still lacking.

## Abbreviations

FDR: False discovery rate; GLMM: Generalized linear mixed model; JSD: Jensen–Shannon distance; OTU: Operational taxonomic unit; PERMANOVA: Permutational analysis of variance; p.i.: Post-infection; SCFA: Short-chain fatty acid; SD: Standard deviation; SE: Standard error.

## Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s13071-022-05535-w>.

**Additional file 1.** Results of PERMANOVA for each day p.i. testing the effect of the *A. suum* experimental group (single-infection, trickle-infection, non-infected control) on microbiota composition in porcine faecal samples, based on Jensen–Shannon distances. Significant *P*-values are printed in bold.

**Additional file 2.** Hub nodes identified in faecal microbial association networks from *Ascaris suum* single-infected and trickle-infected pigs and a non-infected control group based on eigenvector centrality. Hubs shared between experimental groups on the same day are highlighted in yellow, while hubs occurring in subsequent networks of the same group are highlighted in green.

**Additional file 3.** Results of PERMANOVAs testing the effect of each *A. suum* experimental group vs the control group on the microbiota composition in ingesta samples from the ileum, caecum and colon on days 21, 35 and 49 p.i. Analyses were based on OTU-level Jensen–Shannon distance values.

**Additional file 4.** Heatmap of differentially abundant species in the (A) ileum and (B) caecum and colon of *A. suum*-infected pigs compared with a non-infected control group, as determined by DESeq2 analysis. Only differences with Benjamini–Hochberg-adjusted *P*-values  $\leq 0.01$  are shown as coloured tiles.

## Acknowledgements

The authors would like to thank Ulla Kuettler for excellent technical assistance and Sebastian Sprengel of Microsynth SeqLab for technical advice.

## Author contributions

AS and CS designed the study. CS acquired the funding and supervised the project. AS, SKo and SKI performed the animal experiments and collected the samples. AS analysed and interpreted the data and drafted the manuscript. LW, DB and CS contributed to data interpretation. LW, SKo, SKI, GB, DB and CS reviewed and edited the manuscript. All authors read and approved the final manuscript.

## Funding

Open Access funding enabled and organized by Projekt DEAL. This work was financially supported by the Deutsche Forschungsgemeinschaft (German Research Foundation; Grant Number STR 1171/16-1).

## Availability of data and materials

The datasets generated during the current study are available in the National Center for Biotechnology Information Sequence Read Archive (SRA) under the SRA accession: PRJNA790705.

## Declarations

### Ethics approval and consent to participate

Animal experiments were conducted in accordance with the German Animal Welfare act as well as the national and international guidelines for animal welfare and were approved by the ethics commission of the Animal Care and Use Committee of the Lower Saxony State Office for Consumer Protection and Food Safety (*Niedersächsisches Landesamt für Verbraucherschutz und Lebensmittelsicherheit*) under reference number 33.19.42502-04-16/2272.

### Consent for publication

Not applicable.

## Competing interests

The authors declare that there are no competing interests.

## Author details

<sup>1</sup>Institute for Parasitology, Centre for Infection Medicine, University of Veterinary Medicine Hannover, Buenteweg 17, 30559 Hanover, Germany. <sup>2</sup>Department of Safety and Quality of Meat, Max Rubner-Institut, Federal Research Institute of Nutrition and Food, E.-C.-Baumann-Straße 20, 95326 Kulmbach, Germany. <sup>3</sup>Institute for Physiology and Cell Biology, University of Veterinary Medicine Hannover, Bischofsholer Damm 15, 30173 Hanover, Germany.

Received: 28 July 2022 Accepted: 10 October 2022

Published online: 17 November 2022

## References

- Sommer F, Backhed F. The gut microbiota—masters of host development and physiology. *Nat Rev Microbiol.* 2013;11:227–38. <https://doi.org/10.1038/nrmicro2974>.
- Midha A, Ebner F, Schlosser-Brandenburg J, Rausch S, Hartmann S. Trilateral relationship: *Ascaris*, microbiota, and host cells. *Trends Parasitol.* 2021;37:251–62. <https://doi.org/10.1016/j.pt.2020.09.002>.
- Zhou C, Chen J, Niu H, Ouyang S, Wu X. Study on the population evolution of *Ascaris lumbricoides* and *Ascaris suum* based on whole genome resequencing. *Vet Parasitol.* 2020;279:109062. <https://doi.org/10.1016/j.vetpar.2020.109062>.
- Herricks JR, Hotez PJ, Wanga V, Coffeng LE, Haagsma JA, Basáñez M-G, et al. The global burden of disease study 2013: what does it mean for the NTDs? *PLOS Negl Trop Dis.* 2017;11:e0005424. <https://doi.org/10.1371/journal.pntd.0005424>.
- Roepstorff A, Eriksen L, Slotved HC, Nansen P. Experimental *Ascaris suum* infection in the pig: worm population kinetics following single inoculations with three doses of infective eggs. *Parasitology.* 1997;115:443–52. <https://doi.org/10.1017/S0031182097001480>.
- Masure D, Wang T, Vlaminc J, Claerhoudt S, Chiers K, Van den Broeck W, et al. The intestinal expulsion of the roundworm *Ascaris suum* is associated with eosinophils, intra-epithelial T cells and decreased intestinal transit time. *PLOS Negl Trop Dis.* 2013;7:e2588. <https://doi.org/10.1371/journal.pntd.0002588>.
- Pilitt PA, Lichtenfels JR, Tromba FG, Madden PA. Differentiation of late fourth and early fifth stages of *Ascaris suum* Goeze, 1782 (Nematoda: Ascaridoidea) in swine. *Proc Helminthol Soc Wash.* 1981;48:1–7.
- Nansen P, Roepstorff A. Parasitic helminths of the pig: factors influencing transmission and infection levels. *Int J Parasitol.* 1999;29:877–91. [https://doi.org/10.1016/S0020-7519\(99\)00048-X](https://doi.org/10.1016/S0020-7519(99)00048-X).
- Thamsborg SM, Nejsum P, Mejer H. Chapter 14—impact of *Ascaris suum* in Livestock. In: Holland C, editor. *Ascaris: the neglected parasite.* Amsterdam: Elsevier; 2013. p. 363–81.
- Clark P, Bilkei G. Production losses due to ascarid infestation of outdoor maintained pigs. *Tierarztl Umsch.* 2003;58:425–31.
- Dawson HD, Beshah E, Nishi S, Solano-Aguilar G, Morimoto M, Zhao A, et al. Localized multigene expression patterns support an evolving Th1/Th2-like paradigm in response to infections with *Toxoplasma gondii* and *Ascaris suum*. *Infect Immun.* 2005;73:1116–28. <https://doi.org/10.1128/iai.73.2.1116-1128.2005>.
- Solano-Aguilar G, Shea-Donohue T, Madden KB, Quinoñes A, Beshah E, Lakshman S, et al. *Bifidobacterium animalis* subspecies *lactis* modulates the local immune response and glucose uptake in the small intestine of juvenile pigs infected with the parasitic nematode *Ascaris suum*. *Gut Microbes.* 2018;9:422–36. <https://doi.org/10.1080/19490976.2018.1460014>.
- Koehler S, Springer A, Issel N, Klinger S, Strube C, Breves G. Changes in porcine nutrient transport physiology in response to *Ascaris suum* infection. *Parasit Vectors.* 2021;14:533. <https://doi.org/10.1186/s13071-021-05029-1>.
- Koehler S, Springer A, Issel N, Klinger S, Wendt M, Breves G, et al. *Ascaris suum* nutrient uptake and metabolic release, and modulation of host intestinal nutrient transport by excretory-secretory and cuticle antigens in vitro. *Pathogens.* 2021;10:1419. <https://doi.org/10.3390/pathogens10111419>.

15. Midha A, Janek K, Niewianda A, Henklein P, Guenther S, Serra DO, et al. The intestinal roundworm *Ascaris suum* releases antimicrobial factors which interfere with bacterial growth and biofilm formation. *Front Cell Infect Microbiol*. 2018. <https://doi.org/10.3389/fcimb.2018.00271>.
16. Li RW, Wu S, Li W, Navarro K, Couch RD, Hill D, et al. Alterations in the porcine colon microbiota induced by the gastrointestinal nematode *Trichuris suis*. *Infect Immun*. 2012;80:2150–2157. <https://doi.org/10.1128/iai.00141-12>.
17. Williams AR, Krych L, Fauzan Ahmad H, Nejsup P, Skovgaard K, Nielsen DS, et al. A polyphenol-enriched diet and *Ascaris suum* infection modulate mucosal immune responses and gut microbiota composition in pigs. *PLoS ONE*. 2017;12:e0186546–e0186546. <https://doi.org/10.1371/journal.pone.0186546>.
18. Andersen-Civil AIS, Myhill LJ, Büdeyri Gökçöz N, Engström MT, Mejer H, Zhu L, et al. Dietary proanthocyanidins promote localized antioxidant responses in porcine pulmonary and gastrointestinal tissues during *Ascaris suum*-induced type 2 inflammation. *FASEB J*. 2022;36:e22256. <https://doi.org/10.1096/fj.2021101603RR>.
19. Wang Y, Liu F, Urban JF, Paerewijck O, Geldhof P, Li RW. *Ascaris suum* infection was associated with a worm-independent reduction in microbial diversity and altered metabolic potential in the porcine gut microbiome. *Int J Parasitol*. 2019;49:247–56. <https://doi.org/10.1016/j.ijpara.2018.10.007>.
20. Mejer H, Roepstorff A. *Ascaris suum* infections in pigs born and raised on contaminated paddocks. *Parasitology*. 2006;133:305–12. <https://doi.org/10.1017/S0031182006000394>.
21. Davis A, Kohler C, Alsallaq R, Hayden R, Maron G, Margolis E. Improved yield and accuracy for DNA extraction in microbiome studies with variation in microbial biomass. *Biotechniques*. 2019;66:285–9. <https://doi.org/10.2144/btn-2019-0016>.
22. Herlemann DPR, Labrenz M, Jürgens K, Bertilsson S, Waniek JJ, Andersson AF. Transitions in bacterial communities along the 2000 km salinity gradient of the Baltic Sea. *ISME J*. 2011;5:1571–9. <https://doi.org/10.1038/ismej.2011.41>.
23. Martin M. Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnetjournal*. 2011;17:10–2. <https://doi.org/10.14806/ej.17.1.200>.
24. Edgar RC. Search and clustering orders of magnitude faster than BLAST. *Bioinformatics*. 2010;26:2460–1. <https://doi.org/10.1093/bioinformatics/btq461>.
25. Edgar RC, Flyvbjerg H. Error filtering, pair assembly and error correction for next-generation sequencing reads. *Bioinformatics*. 2015;31:3476–82. <https://doi.org/10.1093/bioinformatics/btv401>.
26. Edgar RC. UNCRSS: filtering of high-frequency cross-talk in 16S amplicon reads. *bioRxiv*. 2016. <https://doi.org/10.1101/088666>.
27. Cole JR, Wang Q, Fish JA, Chai B, McGarrell DM, Sun Y, et al. Ribosomal Database Project: data and tools for high throughput rRNA analysis. *Nucleic Acids Res*. 2014;42:D633–42. <https://doi.org/10.1093/nar/gkt1244>.
28. Edgar RC. SINTAX: a simple non-Bayesian taxonomy classifier for 16S and ITS sequences. *bioRxiv*. 2016. <https://doi.org/10.1101/074161>.
29. R Core Team: R: A language and environment for statistical computing. 4.1.0 edn. Vienna, Austria: R Foundation for Statistical Computing; 2021.
30. McMurdie PJ, Holmes S. phyloseq: an R package for reproducible interactive analysis and graphics of microbiome census data. *PLoS ONE*. 2013;8:e61217. <https://doi.org/10.1371/journal.pone.0061217>.
31. Oksanen J, Blanchet FG, Kindt R, Legendre P, Minchin PR, O'Hara R, et al. vegan: Community ecology package. R package version 2.5-6. <https://CRAN.R-project.org/package=vegan>. 2019. Accessed 25 Mar 2020.
32. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol*. 2014;15:550. <https://doi.org/10.1186/s13059-014-0550-8>.
33. Kuznetsova A, Brockhoff PB, Christensen RHB. lmerTest package: tests in linear mixed effects models. *J Stat Softw*. 2017;82:1–26. <https://doi.org/10.18637/jss.v082.i13>.
34. Morris EK, Caruso T, Buscot F, Fischer M, Hancock C, Maier TS, et al. Choosing and using diversity indices: insights for ecological applications from the German Biodiversity Exploratories. *Ecol Evol*. 2014;4:3514–24. <https://doi.org/10.1002/ece3.1155>.
35. Peschel S, Müller CL, von Mutius E, Boulesteix A-L, Depner M. NetCoMi: network construction and comparison for microbiome data in R. *Brief Bioinform*. 2020;22:bbaa290. <https://doi.org/10.1093/bib/bbaa290>.
36. Kurtz ZD, Müller CL, Miraldi ER, Littman DR, Blaser MJ, Bonneau RA. Sparse and compositionally robust inference of microbial ecological networks. *PLOS Comput Biol*. 2015;11:e1004226. <https://doi.org/10.1371/journal.pcbi.1004226>.
37. Clauset A, Newman ME, Moore C. Finding community structure in very large networks. *Phys Rev E Stat Nonlin Soft Matter Phys*. 2004;70:066111. <https://doi.org/10.1103/PhysRevE.70.066111>.
38. Richards JD, Gong J, Lange CFM. The gastrointestinal microbiota and its role in monogastric nutrition and health with an emphasis on pigs: current understanding, possible modulations, and new technologies for ecological studies. *Can J Anim Sci*. 2005;85:421–35. <https://doi.org/10.4141/a05-049>.
39. De Rodas B, Youmans BP, Danzeisen JL, Tran H, Johnson TJ. Microbiome profiling of commercial pigs from farrow to finish. *J Anim Sci*. 2018;96:1778–94. <https://doi.org/10.1093/jas/sky109>.
40. Chong CYL, Vatanen T, Oliver M, Bloomfield FH, O'Sullivan JM. The microbial biogeography of the gastrointestinal tract of preterm and term lambs. *Sci Rep*. 2020;10:9113. <https://doi.org/10.1038/s41598-020-66056-z>.
41. Ke S, Fang S, He M, Huang X, Yang H, Yang B, et al. Age-based dynamic changes of phylogenetic composition and interaction networks of health pig gut microbiome feeding in a uniformed condition. *BMC Vet Res*. 2019;15:172. <https://doi.org/10.1186/s12917-019-1918-5>.
42. Han GG, Lee J-Y, Jin G-D, Park J, Choi YH, Kang S-K, et al. Tracing of the fecal microbiota of commercial pigs at five growth stages from birth to shipment. *Sci Rep*. 2018;8:6012. <https://doi.org/10.1038/s41598-018-24508-7>.
43. Zhang L, Wu W, Lee Y-K, Xie J, Zhang H. Spatial heterogeneity and co-occurrence of mucosal and luminal microbiome across swine intestinal tract. *Front Microbiol*. 2018. <https://doi.org/10.3389/fmicb.2018.00048>.
44. Kelly J, Daly K, Moran AW, Ryan S, Bravo D, Shirazi-Beechey SP. Composition and diversity of mucosa-associated microbiota along the entire length of the pig gastrointestinal tract; dietary influences. *Environ Microbiol*. 2017;19:1425–38. <https://doi.org/10.1111/1462-2920.13619>.
45. Scotti R, Southern S, Boinett C, Jenkins TP, Cortés A, Cantacessi C. MICHELINdb: a web-based tool for mining of helminth-microbiota interaction datasets, and a meta-analysis of current research. *Microbiome*. 2020;8:10. <https://doi.org/10.1186/s40168-019-0782-7>.
46. Rosa BA, Supali T, Gankpala L, Djuardi Y, Sartono E, Zhou Y, et al. Differential human gut microbiome assemblages during soil-transmitted helminth infections in Indonesia and Liberia. *Microbiome*. 2018;6:33. <https://doi.org/10.1186/s40168-018-0416-5>.
47. Rubel MA, Abbas A, Taylor LJ, Connell A, Tanes C, Bittinger K, et al. Lifestyle and the presence of helminths is associated with gut microbiome composition in Cameroonians. *Genome Biol*. 2020;21:122. <https://doi.org/10.1186/s13059-020-02020-4>.
48. Gordon CA, Krause L, McManus DP, Morrison M, Weerakoon KG, Connor MC, et al. Helminths, polyparasitism, and the gut microbiome in the Philippines. *Int J Parasitol*. 2020;50:217–25.
49. Koh A, De Vadder F, Kovatcheva-Datchary P, Bäckhed F. From dietary fiber to host physiology: short-chain fatty acids as key bacterial metabolites. *Cell*. 2016;165:1332–45. <https://doi.org/10.1016/j.cell.2016.05.041>.
50. Zaiss Mario M, Rapin A, Lebon L, Dubey Lalit K, Mosconi I, Sarter K, et al. The intestinal microbiota contributes to the ability of helminths to modulate allergic inflammation. *Immunity*. 2015;43:998–1010. <https://doi.org/10.1016/j.immuni.2015.09.012>.
51. Kim CH, Park J, Kim M. Gut microbiota-derived short-chain fatty acids, T cells, and inflammation. *Immune Netw*. 2014;14:277–88. <https://doi.org/10.4110/in.2014.14.6.277>.
52. Midttun HLE, Acevedo N, Skallerup P, Almeida S, Skovgaard K, Andresen L, et al. *Ascaris suum* infection downregulates inflammatory pathways in the pig intestine in vivo and in human dendritic cells in vitro. *J Infect Dis*. 2017;217:310–9. <https://doi.org/10.1093/infdis/jix585>.
53. Amat S, Lantz H, Munyaka PM, Willing BP. *Prevotella* in pigs: The positive and negative associations with production and health. *Microorganisms*. 2020;8:1584. <https://doi.org/10.3390/microorganisms8101584>.
54. Mach N, Berri M, Estellé J, Levenez F, Lemonnier G, Denis C, et al. Early-life establishment of the swine gut microbiome and impact on host phenotypes. *Environ Microbiol Rep*. 2015;7:554–69. <https://doi.org/10.1111/1758-2229.12285>.

55. Hale OM, Stewart TB, Marti OG. Influence of an experimental infection of *Ascaris suum* on performance of pigs. *J Anim Sci*. 1985;60:220–5. <https://doi.org/10.2527/jas1985.601220x>.
56. Vlaminck J, Düsseldorf S, Heres L, Geldhof P. Serological examination of fattening pigs reveals associations between *Ascaris suum*, lung pathogens and technical performance parameters. *Vet Parasitol*. 2015;210:151–8. <https://doi.org/10.1016/j.vetpar.2015.04.012>.
57. Forsum E, Nesheim MC, Crompton DWT. Nutritional aspects of *Ascaris* infection in young protein-deficient pigs. *Parasitology*. 1981;83:497–512. <https://doi.org/10.1017/S0031182000080483>.
58. Carrera E, Nesheim MC, Crompton DW. Lactose maldigestion in *Ascaris*-infected preschool children. *Am J Clin Nutr*. 1984;39:255–64. <https://doi.org/10.1093/ajcn/39.2.255>.
59. Stolzenbach S, Myhill LJ, Andersen LOB, Krych L, Mejer H, Williams AR, et al. Dietary inulin and *Trichuris suis* infection promote beneficial bacteria throughout the porcine gut. *Front Microbiol*. 2020;11:312. <https://doi.org/10.3389/fmicb.2020.00312>.
60. Reynolds LA, Smith KA, Filbey KJ, Harcus Y, Hewitson JP, Redpath SA, et al. Commensal-pathogen interactions in the intestinal tract. *Gut Microbes*. 2014;5:522–32. <https://doi.org/10.4161/gmic.32155>.
61. Iraporda C, Errea A, Romanin DE, Cayet D, Pereyra E, Pignataro O, et al. Lactate and short chain fatty acids produced by microbial fermentation downregulate proinflammatory responses in intestinal epithelial cells and myeloid cells. *Immunobiology*. 2015;220:1161–9. <https://doi.org/10.1016/j.imbio.2015.06.004>.
62. Turróni F, Ventura M, Buttó LF, Duranti S, O'Toole PW, Motherway MOC, et al. Molecular dialogue between the human gut microbiota and the host: a *Lactobacillus* and *Bifidobacterium* perspective. *Cell Mol Life Sci*. 2014;71:183–203. <https://doi.org/10.1007/s00018-013-1318-0>.
63. Jensen AN, Mejer H, Mølbak L, Langkjær M, Jensen TK, Angen Ø, et al. The effect of a diet with fructan-rich chicory roots on intestinal helminths and microbiota with special focus on *Bifidobacteria* and *Campylobacter* in piglets around weaning. *Animal*. 2011;5:851–60. <https://doi.org/10.1017/S175173111000251X>.
64. Jørgensen RJ, Nansen P, Nielsen K, Eriksen L, Andersen S. Experimental *Ascaris suum* infection in the pig. Population kinetics following low and high levels of primary infection in piglets. *Vet Parasitol*. 1975;1:151–7. [https://doi.org/10.1016/0304-4017\(75\)90017-5](https://doi.org/10.1016/0304-4017(75)90017-5).
65. Boes J, Medley GF, Eriksen L, Roepstorff A, Nansen P. Distribution of *Ascaris suum* in experimentally and naturally infected pigs and comparison with *Ascaris lumbricoides* infections in humans. *Parasitology*. 1998;117:589–96. <https://doi.org/10.1017/S0031182098003382>.

## Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more [biomedcentral.com/submissions](https://biomedcentral.com/submissions)

