



Review article


A systematic scoping review of microbial pathogens in ruminants with or without a history of abortions in Nigeria

Kabiru O. Akinyemi^{1*}, Samuel O. Ajoseh¹, Abdul-Azeez Anjorin¹, Wasiu O. Salami¹, Aminat O. Lawal¹, Marwa Bassiouny², Heinrich Neubauer² and Gamal Wareth^{2,3}

¹ Department of Microbiology, Faculty of Science, Lagos State University, Ojo, Lagos, Nigeria

² Friedrich-Loeffler-Institut, Institute of Bacterial Infections and Zoonoses (IBIZ), 96a, D-07743 Jena, Germany

³ Institute of Infectious Diseases and Infection Control, Jena University Hospital, Am Klinikum 1, 07747 Jena, Germany



Abstract

Abortifacient pathogens such as bacterial [*Brucella* spp., *Listeria* spp., *Leptospira interrogans* ser., *Coxiella burnetii*, *Campylobacter* spp., *Anaplasma* spp., *Chlamydia* spp.], mycotic [*Aspergillus* spp. and *Candida* spp.], protozoan [*Toxoplasma gondii*, *Neospora* spp.], and viral [Blue-tongue virus (BTV), Schmallenberg virus (SBV), Bovine viral diarrhea virus (BVDV), and Peste des petits ruminants virus (PPRV)] pathogens are challenges for the productive and reproductive performance of ruminants (cattle, sheep, and goats) globally. No comprehensive report on epidemiology, associated risk factors, or economic burden of these infectious pathogens is available for Nigeria. This review estimated the distribution and burden of abortive pathogens in ruminants in Nigeria for the last twenty-two years (2000-2022). Research articles reporting the detection of any of the above-mentioned abortive pathogens in ready-to-slaughter ruminants (RTSR), sick ruminants (SR), and ruminants with abortive history (RAWH) in Nigeria were accessed using different repositories, including Google Scholar, Proquest, PubMed, ResearchGate and Scopus to determine the prevalence, spatial distribution, and associated risk factors. From a total of 140 articles selected for this review, eight bacterial, four viral, two parasitic, and two mycotic infectious agents were reported for Nigeria. This study reveals a prevalence of 28.2% viral agents, 14.43% bacterial pathogens, 14.24% protozoans, and 28.1% fungal agents in the reported tested samples. Brucellosis was the most often reported among bacterial diseases, followed by leptospirosis and listeriosis. PPRV infection was the most common viral disease, followed by BTV. Additionally, two parasitic diseases, neosporosis and toxoplasmosis, and two mycotic diseases, aspergillosis and candidiasis, were reported. In this study, stillbirth and abortion were recorded in 49.2% of sheep with PPRV, 58.95% in goats with *Chlamydomytila abortus* and PPRV, and 6.4% in cattle with *Brucella abortus* and *Histophilus somni* infections. Lack of vaccines, open markets, and extensive husbandry systems were among the risk factors associated with different abortive pathogens. This study is a useful tool for researchers and government officers in risk assessment and management of livestock to improve livestock production in Nigeria.

Keywords: Abortive diseases, Brucellosis, Morbidity, Ruminants, Stillbirth

Citation: Akinyemi, K. O., Ajoseh, S. O., Anjorin, A., Salami, W. S., Lawal, A. O., Bassiouny, M., Neubauer, H. and Wareth, G. A systematic scoping review of microbial pathogens in ruminants with or without a history of abortions in Nigeria. Ger. J. Vet. Res. 3 (3): 34-51. <https://doi.org/10.51585/gjvr.2023.3.0062>

Introduction

The term "abortion" refers to the termination of pregnancy between the 42nd and 28th day of gestation for cattle, the 20th and 150th for goats, and the 21st and 152nd for sheep (Peter, 2000). Farmers and veterinarians often face difficulties in diagnosing the cause of abortions. Herds have experienced a precipitous and severe increase in abortions over a prolonged period, with huge and inevitable financial losses in the livestock sector (De Vries, 2006; Hossein-Zadeh, 2013). Infectious and non-infectious causes contribute to ruminant abortion. In a few studies, non-infectious variables like hereditary and non-genetic illnesses have been reported (Peter, 2000; Hansen, 2002; Okoli, 2003; Sani and Amanloo, 2007; Sen et al., 2010; Dereje et al., 2018).

Heat stress, production stress, seasonal effects, and seasonal fluctuations are the most significant non-genetic risks for abortion (Hansen, 2002; Sani and Amanloo, 2007). Genetic disorders, such as chromosomal abnormalities and single genes, can lead to abortion in cows, sheep, goats, and calves' sterility (Okoli, 2003). However, infectious reproductive diseases, including bacterial, viral, fungal, and parasitic agents, are responsible for massive

fetal deaths in ruminants (Okoli, 2003; Sen et al., 2010). Depending on the stage of pregnancy, different syndromes are noticed. In cases where the placenta experiences a rapid bacterial infection from the early stages of gestation through the other stages, foetal death can occur due to sepsis (Dereje et al., 2018). The deceased foetus is often ejected within five days, and autolytic alterations cover the small gross lesions induced by the pathogens (Quinn et al., 2011).

The retention of the foetal membranes is a typical side effect of metritis. The foetus may be born normally if the infection happens at a late stage of gestation, but it may not survive (Quinn et al., 2001). From an economic point of view, more than ten diseases are responsible for stillbirth and abortion in both large and small ruminants (Quinn et al., 2001), with brucellosis and peste des petits ruminant (PPR) infections being the most important of these diseases due to the large economic loss caused particularly in endemic areas (El-Yuguda et al., 2013; Akinyemi et al., 2022). Other significant infectious abortive diseases in ruminants include anaplasmosis, leptospirosis, listeriosis, bovine viral diarrhea (BVD), campylobacteriosis, chlamyphilosis, my-




Figure 1: Flowchart of a systematic review of abortion diseases in ruminants for selection of eligible articles. Ready-to-slaughter ruminants (RTSR), sick ruminants (SR), or ruminants with a history of abortion (RWAH).

cotic abortion, neosporosis, *Arcanobacterium pyogenic* disease, Q-fever, bluetongue viral disease (BTV), Schmallenberg virus (SBV) infection, *Tritrichomonas foetus* disease, epizootic bovine abortion, and toxoplasmosis (Dereje et al., 2018). Although the precise percentage of cases caused by infectious agents is unknown, 90% of cases in which an aetiological diagnosis is made have an infectious origin (Parthiban et al., 2015). In Nigeria, ruminants constitute a substantial proportion of the livestock sub-sector, and about 80% of the estimated ruminant population lives in the Northern part of Nigeria (Lawal et al., 2012; El-Yuguda et al., 2013). The estimated cattle population is 34.5 million, with 22.1 million sheep and 13.9 million goats. In the recent past, livestock contributed an estimated 23.3% to the Nigerian National Gross Domestic Product (GDP) and 9% to the agro-economy (Neils et al., 2011). To meet the food needs of a growing population (red meat and dairy products), farmers and the government paid attention to improving the breed, pasture, and animal health to boost livestock productivity and address the high rate of urbanization. Several factors have hampered these efforts, of which infectious reproductive health complications are the main impact. In ruminants, abortive diseases have been of paramount interest since these diseases are connected with significant financial losses to the livestock industry (Neils et al., 2011). However, there is no comprehensive review of microbial pathogens in ruminants with or without a history of abortions and their associated risk factors. Thus, this study addresses these gaps based on the review of published articles in the last two decades in Nigeria.

Materials and methods

Area of study

This study was conducted in six geopolitical zones of Nigeria including 36 States: North-East (NE), 6 States: North-West (NW), 7 States: North-Central (NC), 6 States: South-East (SE), 5 States: South-West (SW), 6 States and South-South (SS), 6 States including a federal capital territory (Abuja). The North-West zone comprises 242,425 km², or 25.75%, of the nation's total land area. There are around 272,451 km² in the North-East zone. The North-Central zone spans over 193,088 km² of the state. In the Southern zone of Nigeria, however, 79,665 km² of land is occupied by the South-West zone, 79,525 km² by the South-South zone and roughly 84,587 km² by the South-East zone.

Search strategy and data acquisition

The systematic review protocols (PRISMA-P) 2015 (Moher et al., 2015) checklist was followed for the review of meta-analysis of original research studies that reported twenty microbial pathogens in cattle and small ruminants with or without a history of abortion using public scientific databases such as Google Scholar, Scopus, PubMed, Proquest, Academia, and ResearchGate, from January 1, 2000, to December 1, 2022. The objective was to identify original articles that reported on prevalence, serotypes, and antibiotic resistance in all six geopolitical zones of Nigeria, with a specific focus on culture, serology, and molecular approaches.

Additionally, the following search strings/terms were used: Peste des petits ruminants virus, *Brucella*, *Ureaplasma*, *Anaplasma* spp., *Anaplasma marginale*, *Leptospira interrogans* ser, *Listeria monocytogenes*, viral diarrhea virus, *Campylobacter* spp., *Campylobacter fetus*, *Chlamydomphila* spp., *Arch-nobacterium pyogen*, *Aspergillus* spp., *Candida* spp., *Mortellella wolfii*, *Neospora caninum*, *Coxiella burnetii*, BTV, SBV, *Tritrichomonas foetus*, *Histophilus somni*, *Toxoplasma* spp., *Toxoplasmas gondii*; or brucellosis, undulant fever, Mediterranean fever, gastric remittent fever, Malta fever, ureaplasmosis, anaplasmosis, leptospirosis, listeriosis, bovine viral diarrhoea (BVD), campylobacteriosis, chlamydomphilitis mycotic abortion, neosporosis, *Arch-nobacterium pyogen* disease, Q-fever, bluetongue viral disease, schmallenberg viral infection, *Tritrichomonas foetus* disease, *Histophilus somni* infection, epizootic bovine abortion, toxoplasmosis, bovine herpes virus Pestis des petits, animals, cattle, small ruminants, febrile, abortion, abortive, incidence, prevalence, sensitivity testing, antibiotic susceptibility, antimicrobial activity, or treatment.

Selection criteria and inclusion criteria

Online publications (full text/abstracts) were individually evaluated. Following this study's inclusion criteria, relevant papers were retrieved and examined, including their reference sections. Publications including full-text articles, abstracts, dissertations, thesis, short communication, and conference proceedings in the English language on cohort, case, and cross-sectional studies were included if samples were taken from a named farm, herd, community, slaughterhouse/abattoir, state, or geopolitical zone, or if the study recruited animals (cattle, sheep, and goats) originated from Nigeria.




Figure 2: Categories of abortive diseases and numbers of publications in these statuses from 2000-2022. Abbreviations: ready-to-slaughter ruminants (RTSR), sick ruminants (SR), or ruminants with a history of abortion (RWHA), bovine viral diarrhoea (BVD), bluetongue viral disease (BTV), Schmallenberg virus (SBV), and peste des petits ruminant virus (PPRV).

The following abortive pathogens were of concern: *Brucella*, PPRV, *Ureaplasma* spp., *Anaplasma* spp., *Anaplasma marginale*, *Leptospira interrogans*, *Listeria monocytogenes*, BVD, *Campylobacter* spp., *Campylobacter fetus*, *Chlamydia* spp., *Archaeobacterium pyogenes*, *Aspergillus* spp., *Candida* spp., *Mucor* spp., *Mortellella wolfii*, *Neospora caninum*, *C. burnetii*, BTV, SBV, *Trichomonas foetus*, *H. somni*, agents of epizootic bovine abortion, *Toxoplasma* spp., and *Toxoplasma gondii*. Ready-to-slaughter ruminants (RTSR), sick ruminants (SR), or ruminants with a history of abortion (RWHA) were included.

Exclusion criteria

Studies were disregarded if the reported abortive pathogens were not supported by serology, culture, or molecular approach results and if the abortive pathogens in the inclusion criteria were not isolated or detected from RTSR, SR, or RWHA samples. This study did not include reports based on books, book chapters, systematic reviews, experimental studies, ecological associations, or clinical diagnoses. Two de-duplication tools, including the Mendeley citation management and Ovid multi-file search, were used to identify and eliminate duplicate articles.

Data extraction

From each of the chosen articles, descriptive and quantitative data on the inclusion criteria established were evaluated and a standard template was made using a Microsoft Excel 2013 file. Double data extraction and entry were done to assure accuracy. State, geopolitical zone, year of sampling/study, first author's name, publication date, date of sample collection, source/type of sample (blood, body fluid, secretion), sample size, number of ruminants, types of animals, testing method, number of positive samples from serology, culture, and molecular methods, the number of isolates, and species of the abortive pathogen with biotypes were among the data extracted from the articles.

Results

Data analysis

A descriptive analysis of the data obtained was carried out. Between 1st January 2000 and 1st December 2022, 9543 papers were retrieved from the searched online databases. A total of 189 articles were determined to be qualified, and 140 articles

were retained after duplicated papers had been removed. Specifically, 125 articles reported a single abortive pathogen, and 12 articles reported two abortive pathogens, while three reported more than two abortive pathogens in RTSR, SR, and RWHA (Figure 1). Sixteen possible abortifacient pathogens were identified in this investigation, and the prevalence was estimated as the proportion of any abortifacient pathogen isolated/detected from positive samples in the total samples recorded.

Methods of detection of abortifacient pathogens from reviewed publications

The publications that were reviewed reported sixteen different detection techniques. Seven included in Table 1-Table 3 were the most popular or widely used. Among them are polymerase chain reaction (PCR), culture (CTL), serum agglutination test (SAT), and the serum agglutination test with EDTA, Rose Bengal plate test (RBPT), competitive enzyme-linked immunosorbent assay (c-ELISA), indirect enzyme-linked immunosorbent assay (i-ELISA), and immunoglobulin G/immunoglobulin M-enzyme-linked immunosorbent assay (IgG/IgM-ELISA). Other detection techniques include the use of Warthin Starry Silver (WSS), immunohistochemistry and impregnation (IH), microscopic agglutination test (MAT), and microhaematocrit test (MICRO/HCT). Out of the seven extensively reported methods of detection, the RBPT test is most frequently used to identify antibodies to *Brucella* species in 42 articles, followed by c-ELISA (n=33), CTL (n=29), SAT/SAT-EDTA (n=18), PCR (n=10), IgG/IgM-ELISA (n=9), while the least frequent test used was the i-ELISA (n=7). Intriguingly, it should be noted that of the 140 publications examined, 105 (75%) conducted screening on RTSR (Table 1), 23 (16.4%) on SR (Table 2), and 12 (8.6%) on RWHA (Table 3).

Categories of abortive diseases in Nigeria

A total of 140 research articles covering 31 states (Benue, Plateau, Kwara, Niger, Nasarawa, Borno, Yobe, Bauchi, Adamawa, Taraba, Gombe, Kano, Kaduna, Katsina, Sokoto, Jigawa, Zamfara, Enugu, Ebonyi, Imo, Anambra, Abia, Cross River, Akwa-Ibom, Rivers, Bayelsa, Oyo, Osun, Ondo, Lagos, and Ogun) and the federal capital territory (Abuja) were identified. Suspected abortive diseases were reported in six




Figure 3: Geo-political distribution of pathogens suspected to likely cause abortion in ruminants.

geo-political zones in Nigeria. Four categories of these diseases were found, including bacterial, viral, parasitic, and fungal abortive. Eight bacterial genera (*Brucella*, *Leptospira*, *Listeria*, *Anaplasma*, *Chlamydia*, *Campylobacter*, *Histophilus*, and *Coxiella*), four viral agents (BVDV, PPRV, BTV, and SBV), two protozoans (*Neospora* and *Toxoplasma*) and two fungi (*Aspergillus* and *Candida*) were reported from three animal statuses (RTSR, SR, and RWAH). One hundred and one articles on bacterial diseases, i.e., brucellosis (n=53), leptospirosis (n=9), listeriosis (n=5), Q-fever (n=4), campylobacteriosis (n=3), anaplasmosis (n=25), chlamydiosis (n=1) and *H. somni* associated pneumonia (n=1). Also, eight articles on parasitic abortive diseases were reviewed, including neosporosis (n=3) and toxoplasmosis (n=5). Furthermore, 30 articles on viral abortive diseases were reviewed, with PPRV having the highest number of publications (n=24), followed by BTV (n=3), BVDV (n=2), and SBV (n=1), as well as two articles on mycotic infections. Detailed data can be found in Figure 2.

Geographical distribution of suspected pathogens likely to cause abortion in ruminants

According to the literature reviews for this study, the suspected abortive pathogens have been reported in several geopolitical regions of Nigeria. Eight agents were reported in less than four publications and only in a few geopolitical areas. They are *H. somni*, *Chlamydia* spp., *Neospora* spp., BVDV, BTV, SBV, and fungal pathogens (*Aspergillus* spp. and *Candida albicans*). Eight groups of important pathogens were documented in two or more geopolitical zones and published in at least four articles. These include *Leptospira interrogans* ser., *Anaplasma* spp., *Campylobacter* spp., PPRV, *C. burnetii*, *Listeria* spp., *Toxoplasma* spp., and *Brucella* spp. (Figure 3). Notably, *Brucella* spp. and PPRV were reported in all six geopolitical regions of Nigeria (NE, NC, NW, SS, SE, and SW), followed by *Anaplasma*

spp. in five regions (SW, SS, NC, NW, and NE), *Leptospira interrogans* ser. in four regions (NE, NW, SW, and SS), *C. burnetii* in three regions (SW, NW, and NE), and *Listeria* spp. in two regions (NW and NE). Within the six geopolitical zones, there are 36 states, and of these, 31 states had at least one case of one of the sixteen infections known to induce abortion in small ruminants. Only five states (Kebbi-NW, Kogi-NC, Ekiti-SW, Edo-SS, and Delta-SS) had no cases of suspected infections likely to cause abortion.

Prevalence of suspected pathogens likely to cause abortion in ruminants

In this study, the estimated cumulative prevalence of suspected pathogens likely to result in abortion among the pathogens considered is 19.36% (21835/112804). The pooled prevalence s from cattle, sheep, and goats of different statuses of the abortive bacterial pathogen is shown in Table 4, while the pooled prevalence of viral, protozoal, and fungal pathogens is shown in Table 5. A prevalence of 28.2% (11534/40883) for viruses, 28.1% (47/167) for fungi, 14.24% (403/2831) for protozoans, and 14.43% (9851/68252) for bacterial pathogens was noted. The four viral agents that make up 28.2% prevalence are BVDV (57%), BTV (39%), SBV (29%), and PPRV (27.0%). Of the 14.43% bacterial pathogens documented, *H. somni* has the highest (100%) prevalence of the eight bacterial pathogens reported from reviewed literature on RTSR, SR, and RWAH of cattle, goats, and sheep. *Leptospira interrogans* ser prevalence was 27.2%, followed by *Brucella* species at 14%, *Campylobacter* species at 13%, *Anaplasma* species at 23%, *C. burnetii* at 9%, *Chlamydia* spp. at 4%, and *Listeria* spp. at 3.2%. Additionally, the prevalence for the *Neospora* and *Toxoplasma* spp. were 10% and 15.0%, respectively. Interestingly, 28.1% of mycotic infections are caused by two fungal genera (*Aspergillus* spp. and *Candida albicans*) were reported from only two publications in this study.

Table 1: Methods of detecting pathogens likely to cause abortion in RTSR based on sample sources and the number of articles.

No. of articles	Sample type (n)	Pathogen type (n)	Detection methods*								References	
			PCR	CTL	SAT/SAT-EDTA	BPT	c-ELISA	i-ELISA	IgG/IgM-ELISA	Other		
24	Blood (24)	<i>Anaplasma marginale</i> (20); <i>A. centrale</i> (1); <i>A. bigemina</i> (1); <i>A. ovis</i> (5)	18	2	-	-	-	-	-	2	3	Akande et al. (2010); Kamani et al. (2010a); Pukuma et al. (2011); Ademola and Onyiche (2013); Pam et al. (2013); Abdullahi et al. (2014); Olatunde et al. (2014); Anyanwu et al. (2016); Elelu et al. (2016); Lorusso et al. (2016); Adua and Idahor (2017); Deme et al. (2017); David et al. (2018); Adejoh et al. (2019); Atsuwe et al. (2019); Happi et al. (2020a,b); Olorunshola et al. (2020); Gboeloh and Araka (2022); Kamani et al. (2022b) Kamani et al. (2022a); Onyiche et al. (2022)
8	Blood (8); kidney samples (1)	<i>Leptospira interrogans</i> serovar <i>hardjo</i> (4); <i>L. interrogans</i> (2); <i>Leptospira</i> spp. (1); <i>L. pomona</i> (1); <i>L. icterohaemorrhagiae</i> (1)	-	2	-	-	-	-	-	3	3	Agunloye et al. (2000); Junaidu et al. (2011); Ngbede et al. (2012); Bashiru et al. (2013); Ngbede et al. (2013); Oruene and Bekwele (2020) Ajayi et al. (2021); Stephen et al. (2022)
2	Raw milk (2)	<i>Listeria monocytogenes</i> (2); <i>L. innocua</i> (1), <i>L. seelighi</i> (1), <i>L. welshine</i> (1)	-	2	-	-	-	-	-	-	-	Yakubu et al. (2012); Uwanibe et al. (2014)
4	Blood (4)	<i>Coxiella burnetii</i> (4)	-	-	-	-	-	-	4	-	-	Adamu et al. (2018); Cadmus et al. (2020); Adamu et al. (2022)
44	Sera (40); milk (5)	<i>Brucella abortus</i> (39); <i>Brucella</i> spp. (20); <i>Brucella melitensis</i> (3)	1	-	15	35	19	-	-	-	2	Bale et al. (2003); Junaidu and Garba (2006); Junaidu et al. (2006); Cadmus et al. (2008); Bertu et al. (2010); Cadmus et al. (2010); Junaidu et al. (2011); Mbuk et al. (2011); Mohammed et al. (2011); Wungak et al. (2011); Lawal et al. (2012); Mai et al. (2012); Alhaji and Wungak (2013); Cadmus et al. (2013) Cadmus et al. (2021); Kaltungo et al. (2013); Maurice (2013); Ogugua et al. (2014); Zubairu et al. (2014); Dauda et al. (2015); Kaltungo et al. (2015); Adamu et al. (2016); Akinseye et al. (2016) Alhaji et al. (2016); Buhari et al. (2016); Dawang et al. (2016); Jajere et al. (2016); Agada et al. (2017); Aworh et al. (2017); Ayinmode et al. (2017); Farouk et al. (2017); Adamu et al. (2018); Agada et al. (2018); Ekere et al. (2018); Ogugua et al. (2018); Olufemi et al. (2018); Shu'aibu et al. (2018); Kaltungo et al. (2019); Yusuf and Abdulrasheed (2019); Buhari et al. (2020); Cadmus et al. (2020); Ibrahim et al. (2020); Mohammed et al. (2020); Ukwueze et al. (2020)
2	Rectal swab (1); Preputial washing 1)	<i>Campylobacter jejuni</i> (1), <i>C. coli</i> (1), <i>C. lari</i> (1), <i>C. laridis</i> (1), <i>C. fetus</i> (3)	-	2	-	-	-	-	-	-	-	Salihu et al. (2009); Mai et al. (2013)
5	Blood; serum	<i>Toxoplasma gondii</i> (5)	-	-	-	-	-	-	3	1	1	Joshua and Akinwumi (2003); Kamani et al. (2010b); Onyiche and Ademola (2015); Ayinmode and Abiola (2016); Ayinmode et al. (2017) Ayinmode (2013); Ayinmode et al. (2017)
2	Blood (2)	<i>Neospora caninum</i> (2)	-	-	-	-	-	-	-	2	-	Ayinmode (2013); Ayinmode et al. (2017)
3	Blood (4)	Blue tongue virus(1)	2	-	-	-	-	2	-	-	-	Oluwayelu et al. (2011); Oladimeji and Kelechi (2019); Zanna et al. (2021)
1	Blood/serum (1)	Schmallenberg virus (1)	-	-	-	-	-	1	-	-	-	Oluwayelu et al. (2015)
2	Sera (2)	Bovine viral diarrhea virus (2)	-	-	-	-	-	-	-	2	-	Peter et al. (2016); Sanusi et al. (2016)
8	Sera (47); lymph nodes (1); nasal swab (1)	Peste des petits ruminants virus (8)	1	5	-	-	-	-	-	-	3	El-Yuguda et al. (2009a); Lawal et al. (2011); El-Yuguda et al. (2013); Nwobodo et al. (2013); Victor et al. (2017); Bello et al. (2018); Chukwudi et al. (2020); Bukar et al. (2020)
Total number of articles= 105			22	11	15	35	22	7	10	12		

*Keys. Other: (LA: Latex agglutination; LM: Light Microscopy; MAT: Microscopic Agglutination Test; IH: Impregnation and Immunohistochemistry; Micro/HCT: Microhaematocrit); PCR: Polymerase Chain Reaction; SAT: Serum Agglutination Test/SAT-EDTA; c-ELISA: Competitive ELISA; i-ELISA: Indirect-ELISA; CTL: Culture.

Table 2: Methods of detecting pathogens likely to cause abortion in diseased Animals based on sample sources and the number of articles.

No. of articles	Sample type (n)	Pathogen type (n)	Detection Methods*								References	
			PCR	CTL	SAT/SAT-EDTA	BPT	c-ELISA	i-ELISA	IgG/IgM-ELISA	Other		
2	Blood (2)	<i>Listeria species</i> (1); <i>L. monocytogenes</i> (1)	-	2	-	-	-	-	-	-	-	Peter (2015); Idris Ibrahim (2021)
4	Blood/serum (4)	<i>Brucella spp.</i> (4)	-	-	1	4	-	-	-	-	-	Dauda et al. (2015); Bamidele et al. (2020); Cadmus et al. (2020); Ukwueze et al. (2020)
1	Lungs (1); spleen (1); liver (1)	<i>H. somni</i> (1)	-	1	-	-	-	-	-	-	-	Odugbo et al. (2009)
1	Blood (1)	<i>Anaplasma marginale</i> (1); <i>A. centrale</i> (1)	-	-	-	-	-	-	-	-	1	Osue et al. (2022)
1	Milk (1)	<i>Aspergillus niger</i> (1); <i>A. filmigatus</i> (1); <i>Fusarium chlamydosporium</i> (1); <i>A. nidule</i> (1); <i>A. terreus</i> (1); <i>Candida albicans</i> (1)	-	1	-	-	-	-	-	-	-	Adelowo (2020)
14	Sera (10); nasal swab (2); lungs (1); lymph (1); ocular swab (1); liver (1); intestine (1)	Peste des petits ruminants virus (14)	6	1	-	-	6	-	-	-	4	Okoli (2003); Luther et al. (2005); Owai (2007); El-Yuguda et al. (2009b); Lawal et al. (2011); Luka et al. (2011); Ularamu et al. (2012); Woma et al. (2015); Jarikre and Emikpe (2017); Adedeji et al. (2019); Chukwudi et al. (2020, 2021); Idris Ibrahim (2021); Mantip et al. (2022)
Total number of articles= 23			6	5	1	4	6	0	0	5		

*Keys. Other: (LA: Latex agglutination; LM: Light Microscopy; MAT: Microscopic Agglutination Test; IH: Impregnation and Immunohistochemistry; Micro/HCT: Microhaematocrit); PCR: Polymerase Chain Reaction; SAT: Serum Agglutination Test/SAT-EDTA; c-ELISA: Competitive ELISA; i-ELISA: Indirect-ELISA; CTL: Culture.

Table 3: Methods of detecting pathogens likely to cause abortion in RWAH based on sample sources and the number of articles.

No. of articles	Sample type (n)	Pathogen type (n)	Detection methods*								References	
			PCR	CTL	SAT/SAT-EDTA	BPT	c-ELISA	i-ELISA	IgG/IgM-ELISA	Other		
1	Preputial washing (1); cervical-vagina mucus swab (1)	<i>Campylobacter fetus</i> (1)	-	1	-	-	-	-	-	-	-	Mshelia et al. (2012)
1	Blood (1)	<i>Listeria monocytogenes</i> (1)	-	1	-	-	-	-	-	-	-	Peter (2015)
1	Blood	<i>Chlamydia abortus</i> (1)	-	1	-	-	-	-	-	-	-	Abubakar (2015)
1	Blood (1)	<i>Leptospira interrogans ser hardjo</i> (1)	-	-	-	-	-	-	-	1	1	Stephen et al. (2022)
5	Blood/sera (5); milk (6); Vagina Swab (6); hygroma fluid (8)	<i>Brucella abortus</i> (9); <i>Brucella spp.</i> (1)	-	9	2	3	-	-	-	-	-	Ocholi et al. (2004, 2005); Onoja et al. (2008); Mbuk et al. (2011); Bertu et al. (2020)
2	Serum (6)	Peste des petits ruminants virus (6)	-	1	-	-	5	-	-	-	1	Lawal et al. (2011); Wachida et al. (2018)
1	Blood (1)	<i>Aspergillus spp.</i> (1); <i>Candida spp.</i> (1)	-	1	-	-	-	-	-	-	-	Biobaku et al. (2016)
Total number of articles= 12			0	14	2	3	5	0	1	2		

*Keys. Other: (LA: Latex agglutination; LM: Light Microscopy; MAT: Microscopic Agglutination Test; IH: Impregnation and Immunohistochemistry; Micro/HCT: Microhaematocrit); PCR: Polymerase Chain Reaction; SAT: Serum Agglutination Test/SAT-EDTA; c-ELISA: Competitive ELISA; i-ELISA: Indirect-ELISA; CTL: Culture.

Risk factors associated with suspected abortive pathogens

Eight key risk factors related to infections likely to induce abortion in ruminants were documented. Age, season, and breed were the most often mentioned risk factors (Eniolorunda et al., 2008; Kamani et al., 2010a; Pukuma et al., 2011; Salihu et al., 2009; Ngbede et al., 2013; Sanusi et al., 2016; Victor et al., 2017; Ibrahim et al., 2020; Cadmus et al., 2020; Ajayi et al., 2021; Stephen et al., 2022). Seasonally, ruminants are more susceptible to *Leptospira*, *Toxoplasma*, and *Anaplasma* spp. transmission. (Eniolorunda et al., 2008; Kamani et al., 2010a; Pukuma et al., 2011). Additionally, due to the high wind speeds associated with the dry season, ruminants are reportedly more susceptible to PPRV and *Brucella* spp. during that time (Salihu et al., 2009; Victor et al., 2017).

Open grazing, pastoral systems, and semi-intensive husbandry are also among the most frequently mentioned risk factors for *Brucella* spp., *Toxoplasma* spp., PPRV, and BVDV (Kamani et al., 2010a; Sanusi et al., 2016; Victor et al., 2017; Ibrahim et al., 2020). Interestingly, ruminants without a history of immunization are more susceptible to PPRV than animals with a history of vaccination (Victor et al., 2017).

It should be highlighted that the source of the animals, such as the open market for purchase, is another significant risk factor for infection with *Brucella* spp. (Ibrahim et al., 2020), *C. burnetii* (Cadmus et al., 2020) and *Leptospira interrogans ser* (Pukuma et al., 2011). It was discovered that certain animal breeds, such as West Africa Dwarf (WAD) breeds of goats, Yankasa varieties of sheep, and Sokoto Gudali (SG) breeds of cattle, were more susceptible to contracting PPRV, *C. burnetii*, and *Brucella* spp. (Salihu et al., 2009; Victor et al., 2017; Cadmus et al., 2020). Sex (Pukuma et al., 2011; Sanusi et al., 2016; Victor et al., 2017; Stephen et al., 2022), history of abortion, and other reproductive diseases (Ibrahim et al., 2020) are additional reported risk factors as shown in Table 6.

Records of stillbirth and abortion in ruminants infected with microbial pathogens

This study recorded alarming stillbirth and abortion of 49.2% (369/782) in sick sheep with PPRV, 58.95% (619/1050) in sick goats with *Chlamydomphila abortus* and PPRV, and 6.4% (166/2606) in sick cattle with *Brucella abortus* and *H. somni* infections. Interestingly, 79.1% (291/368) of stillbirth and abortion were recorded in sick goats due to *Chlamydia* infection, 50% (95/184) in sick cattle due to histophilosis, 49.7% (319/642) in sick goats due to PPRV infection, and 22.5% (9/40) in sick goat due to aspergillosis and candidiasis. Furthermore, 13.0% (36/270) of abortions in bovine due to campylobacteriosis and 2.0% (35/2152) due to brucellosis were recorded.

Discussion

Infections in ruminants pose a significant danger to veterinary public health and food security due to financial implications (Herrero et al., 2013). Therefore, this study systematically reviews the literature reporting possible abortifacient pathogens in Nigerian ruminants. To our knowledge, this is the first comprehensive analysis of the diversity of possible abortifacient pathogens in ruminants in Nigeria. It was observed that 31 out of the 36 states reported abortive pathogens in ruminants documented by 140 eligible published articles. The lack of information from the remaining five states (Edo, Kebbi, Kogi, Ekiti, and Delta) is an indication of gaps in the published articles, which may be due to a lack of definitive diagnostic tools, inadequate infrastructure for assays for definitive detection of abortive pathogens, and a lack of researchers' interest in studying and publishing reports on these pathogens. In this study, the overall prevalence of the sixteen documented abortive pathogens is proven. The sixteen possible abortifacient pathogens are grouped into four categories according to etiology, i.e., bacterial, viral, protozoan, and fungal disease. A prevalence of 28.2% for viruses, 14.43% for bacterial pathogens, 14.24% for protozoans, and 28.1% for fungi were recorded.

Viral abortifacient diseases in Nigeria

An overall seroprevalence of 28.2% for viral abortive pathogens was documented, such as BVDV (57%), BTV (39%), SBV (29%), and PPRV (27%) prevalence was recorded. Notably, a 57% prevalence of BVDV was reported only in RTSR with BTV and SBV. Although they are not yet confirmed zoonotic, the potential to cause zoonoses may not be ruled out in the future. Similar studies around the world had reported different prevalent rates, such as 46.3% in western China (Gao et al., 2013), 70% in South Africa (Njiro et al., 2011), between 50-90% recorded in Iran (Sani and Amanloo, 2007), Canada (Ahmad et al., 2011) and South Vietnam (Duong et al., 2008). However, a lower prevalence of BVDV was reported in Yucatan cattle in Mexico (Solis-Calderon et al., 2005).

The variation of the prevalent rate recorded at different locations across the globe has been attributed to different factors such as the management method, animal populations, husbandry system, diagnostic techniques, and biosecurity measures (Stand Alenius, 2012). In Nigeria, the first case of BVDV was reported in 1977 in Northern Nigeria (Taylor et al., 1977). After that, in 1994, the pathogen was detected in sick sheep and goats in Maiduguri (Baba et al., 1994) and in 2016 in Sokoto state (Sanusi et al., 2016).

The seroprevalence of BTV in this study was 39%. This result was higher than the 12.2% recorded in China (Gong et al., 2021) and 25.7% in Saudi Arabia (Yousef et al., 2012). The disparity could be attributed to differences in animal populations, study scope, and circulating strains. Interestingly, all Nigerian BTV strains number NIG 1982/10 found from the reviewed articles in this study were similar to the Australian BTV-16 strain through phylogenetic analysis, representing a distinct virus lineage within the major eastern topotype (Mertens et al., 2013). Although SBV has been reported in other countries such as Ethiopia (Sibhat et al., 2018), the Netherlands (Elbers et al., 2012), Iraq (Al-Baroodi, 2021), and Germany (Wernike et al., 2014) but only one article reported a seroprevalence of 29% for SBV in Nigerian cattle from the year 2000 till date (Oluwayelu et al., 2015). An indication that this virus is not common in Nigerian ruminants.

PPRV was the most frequently reported possible abortifacient viral agent with or without a history of abortion, with a pooled seroprevalence of 27.0% from 21 of the 36 states across the six geopolitical zones. This result indicates that this virus is endemic and continually circulating in Nigeria despite the availability of PPRV vaccine intervention (Zhao et al., 2021). The high prevalence rates of PPRV recorded in this study were in consonance with studies documented in other countries like Sudan (42.0%) (Ali et al., 2019), Saudi Arabia (24.1%) (Hemida et al., 2020), Rwanda (18.8%) (Shyaka et al., 2021), and the north-eastern region of India (14.5%) (Balamurugan et al., 2020). In Nigeria, the coverage of the PPRV vaccine is poor (Fadiga et al., 2013; Zhao et al., 2021). It is important to note that among the four abortive viral agents identified in this study, only PPRV was reported in RTSR cattle, sheep, goats, diseased sheep and goats, and RWAH cattle and goats.

Several African countries, Middle Eastern states, the Arabian Peninsula, and south India have been affected by PPRV, causing significant economic hardships (Balamurugan et al., 2010; Muse et al., 2012; Balamurugan et al., 2020). PPRV is considered a major obstacle to the successful rearing of small ruminants and has a profound impact on the financial situation of rural farmers. However, there is a lack of sufficient epidemiological data on PPRV in Nigeria, which is crucial for developing effective preventative and control strategies (Banyard et al., 2010; Zahur et al., 2014). Serological assays such as c-ELISA, agar diffusion, and PCR have been widely used for the detection of other viruses, including BTV, BVDV, and SBV. However, unlike PPRV, detailed information regarding the origin, transmission dynamics, vaccination history, mortality, and origin of BVDV, BTV, and SBV is currently unavailable.

Bacterial abortifacient diseases in Nigeria

Notably, a total of 14.43% bacterial abortive pathogens pooled prevalence was found. Of this, 100% prevalence for *H. somni*, 27% for *Leptospira interrogans ser*, 14% for *Brucella* spp., 13.0%

Table 4: The pooled prevalence of abortive bacterial pathogens from cattle, sheep, and goats of different statuses.

Pathogens detected	Samples	No. of samples	Positive samples	Prevalence (%)	Animal status (no.)*								
					RTSR (no.)			Sick/diseased animal (no.)			RWAH (no.)		
					Goat	Sheep	Cattle	Goat	Sheep	Cattle	Goat	Sheep	Cattle
<i>Leptospira interrogans ser</i>	Blood, kidney samples	1963	534	27	113	20	392	0	0	0	6	0	3
<i>Listeria</i> spp.	Milk, Blood	4989	174	3	0	0	99	10	58	0	5	2	0
<i>Chlamydia</i> spp.	Sera	368	14	4	0	0	0	0	0	0	12	2	0
<i>Campylobacter</i> spp.	Preputial washing, cervical-vagina mucus swab	3284	429	13	0	39	380	0	0	0	0	0	10
<i>Coxiella burnetii</i>	Blood, sera	1683	157	9	46	42	69	0	0	0	0	0	0
<i>Histophilus</i> spp.	Lungs, spleen, and liver samples	184	184	100	0	0	184	0	0	0	0	0	0
<i>Brucella</i> spp.	Blood, milk, aborted fetus, vs., hygroma fluid, Sera	47228	6397	14	640	524	4104	29	17	57	10	376	640
<i>Anaplasma</i> spp.	Blood	8553	1962	23	34	105	1810	0	0	13	0	0	0
Total		68252	9851	13.20	833	730	7038	39	75	70	33	380	653

*RTSR: Ready to slaughter ruminants; RWAH: Ruminants with abortive history; HF: hygroma fluid; no: number..

Table 5: The pooled prevalence of viral, protozoal, and fungal pathogens from cattle, sheep, and goats of different statuses.

Pathogens detected	Samples	No. of samples	Positive samples	Prevalence (%)	Animal status (no.)*								
					RTSR (no.)			Sick/diseased animal (no.)			RWAH (no.)		
					Goat	Sheep	Cattle	Goat	Sheep	Cattle	Goat	Sheep	Cattle
Peste des petits ruminants virus	Lungs, lymph, ocular, sera, large intestine, liver	38796	10602	27	1236	982	32	7180	1172	0	0	0	0
Blue tongue virus	Blood, sera	1262	492	39	26	265	201	0	0	0	0	0	0
Schmallenberg virus	Serum	120	35	29	0	0	35	0	0	0	0	0	0
Bovine viral diarrhoea virus	Sera	705	405	57	40	44	321	0	0	0	0	0	0
Total viral agents		40883	11534	28.2	1302	1291	589	7180	1172	0	0	0	0
<i>Toxoplasma</i> spp.	Blood, serum	2597	379	15	22	30	327	0	0	0	0	0	0
<i>Neospora</i> spp.	Blood, sera	234	24	10	0	0	24	0	0	0	0	0	0
Total Hemo-protozoan		2831	403	14.24	22	30	348	0	0	0	0	0	0
<i>Aspergillus</i> spp.	Blood, sera	167	46	27.54	0	0	0	38	0	0	9	0	0
<i>Candida albicans</i>	Blood, sera	167	1	0.59	0	0	0	38	0	0	9	0	0
Total fungi		167	47	28.1	0	0	0	38	0	0	9	0	0

*RTSR: Ready to slaughter ruminants; RWAH: Ruminants with abortive history; HF: Hygroma fluid; no: number.

for *Campylobacter* spp., 23.3% for *Anaplasma*, 9% for *C. burnetii*, 3.8% for *Chlamydia* spp. and 3.2% for *Listeria* spp. was recorded. Notably, *C. burnetii* and *Neospora* spp. were only reported in RTSR by four and two publications, respectively. These findings imply the risk of zoonotic transmission, posing a significant threat to public health. Of the total prevalence of *Anaplasma* (23.3%) recorded, 23.08% prevalence was found in RTSR cattle, 13.3% in RTSR goats, 7.68% in RTSR sheep, and 18.57% in SR cattle with no record on RWAH.

Six *Anaplasma* spp. were reported, i.e., *A. marginale*, mostly (85%) reported, *A. centrale*, *A. ovis*, *A. platys*, and *A. bigemina*. It should be noted that only *A. marginale* and *A. centrale* were reported in the SR cattle (Osue et al., 2022). The most common detection methods reported include the microscopic hematocrits method used by 22/30 articles, followed by PCR in 7/30 articles, and the cultural method considered by 2/30 articles (Pam et al., 2013; Okwelum et al., 2021). *A. marginale* is widely distributed in Nigeria, covering 14/36 states within 5/6 geo-political zones (Figure 3).

The protozoan *A. marginale* is the cause of anaplasmosis, a contagious, hemolytic, and transmissible disease of cattle, sheep, and goats (Underwood et al., 2015). The genus *Anaplasma* belongs to the family *Anaplasmatocae* within the phylum *Rick-*

ettsiales (Underwood et al., 2015). *A. ovis* and *A. centrale* were equally implicated in anaplasmosis in Nigeria. Anaplasmosis has been reported to be endemic in North African and Middle Eastern countries and represents a threat not only to the economies of these countries but also to public health (Parvizi et al., 2020). Studies conducted in other countries with a high prevalence of anaplasmosis have reported similar findings. For example, in Northern Pakistan, *Anaplasma* spp. was found in 21.7% of ruminants (Niaz et al., 2021), and a prevalence of 27.27% of *A. marginale*, *A. ovis*, and *A. centrale* was reported in Khartoum State, Sudan (Eisawi et al., 2020).

The prevalence of *C. burnetii*, another abortifacient bacterial pathogen investigated, was found to be 9%, which is higher than the 6.1% prevalence recorded in Bangladesh (Chakrabarty et al., 2016) and the 4.0% prevalence reported in Senegal (Magadu and Thompson, 2023). This study demonstrates unequivocally that ruminants are a substantial source of *C. burnetii* and present a serious threat to public health by spreading the disease to people in places where they are raised. Interestingly, histophilosis was reported in diseased ruminants with a 100% prevalence rate. Anorexia, coughing, nasal discharge, dyspnea, diarrhea, neck distension, lethargy, lameness of the limbs, recumbency, ocular discharge, and death (50% case-fatality) were

Table 6: Risk factors associated with abortive pathogens.

Pathogens	Animal	Climate	Gender	Age	Breed	HS	Source	Abortion/ other rep. dis.	VH	Vector	Risk factors <i>p-value</i>	Reference
<i>Anaplasma</i> spp.	Cattle	M-S (rainy)	M	Adult	NIB	-	-	-	-	Tick	<0.05	Pukuma et al. (2011)
<i>Leptospira interrogans</i>	Cattle	-	F	Adult	NIB	-	-	-	-	-	<0.05	Stephen et al. (2022)
<i>L. interrogans</i>	Goat	-	-	< 2	-	-	-	-	-	-	<0.05	Junaidu et al. (2011)
<i>L. interrogans</i>	Cattle	-	-	-	Exotic	-	-	-	-	-	<0.05	Ngbede et al. (2013)
<i>L. interrogans</i>	Cattle, sheep, and goat	M-S (rainy)	-	-	-	-	-	-	-	-	<0.05	Eniolorunda et al. (2008)
BVDV	Cattle	M-S (rainy)	M-F	-	-	-	-	-	-	-	<0.05	Sanusi et al. (2016)
PPRV	Goat	HWS (Dry)	F	Adult	WAD	OG, PS	-	-	WTVH	-	<0.05	Victor et al. (2017)
PPRV	Sheep	HWS (Dry)	F	Adult	Yankasa	OG, PS	-	-	WTVH	-	<0.05	Victor et al. (2017)
<i>Coxiella burnetii</i>	Cattle	-	-	-	SG, CB	-	OM	-	-	-	<0.05	Cadmus et al. (2020)
<i>Toxoplasma gondii</i>	Sheep	Rainy	-	Adult	-	Semi-intensive	-	-	-	-	<0.05	Kamani et al. (2010a)
<i>Toxoplasma gondii</i>	Goat	Rainy	-	Adult	-	Semi-intensive	-	-	-	-	<0.05	Kamani et al. (2010a)
<i>Brucella</i> spp.	Cattle	-	-	-	-	PS	OM	Yes	-	-	<0.05	Ibrahim et al. (2020)
<i>Brucella</i> spp.	Cattle	Dry	F	Adult	SG	-	-	-	-	-	<0.05	Salihu et al. (2009)

*Key. M: male, F: female, M-S (Rainy): May-September (Rainy), HWS: high wind speed in dry season, NIB: non-indigenous breeds, BVDV: bovine viral diarrhoea virus, PPRV: peste des petits ruminants virus, SG: Sokoto Gudani, CB: crossbreed, WAD: West African Dwarf, OM: open market, OG: open grazing, PS: pastoral system, WTVH: without vaccination history. The *p-value* was estimated from the combination of two or more risk factors, Seasons: in tropical regions have only two seasons; rainy and dry, based on rainfall. This differs from temperate regions, where all four seasons are determined by temperature differences.

among the clinical symptoms reported on histophillosis (Odugbo et al., 2009). Similar studies have also been reported from the USA, Canada, Japan, New Zealand, Argentina, Scandinavia, Australia, and Eastern and Western Europe (Siddaramppa and Inzana, 2004).

H. somni is still an emerging disease in Nigeria and Africa. The detection of *H. somni* in Nigerian cattle herds is problematic because, except for *H. influenzae*, this organism is unmatched in the family *Pasteurellaceae* for its versatility as a commensal organism and for the wide range of clinical manifestations that can result from septicemia. Without mentioning the digestive system, *H. somni* infections of numerous bovine organ systems have been documented in all sexes, ages, and feedlots or dairy products in other countries of the world (Siddaramppa and Inzana, 2004).

Notably, both *Listeria* and *Brucella* were the only bacterial abortifacient pathogens reported in the three groups of ruminant animal status. For instance, five publications reported listeriosis, with two from RTSR, two from SR, and one from RWAH. However, of the 53 articles that reported *Brucella* spp., 44 were on RTSR, four on SR, and five on RWAH. Of the 3.2% *Listeria* spp. prevalence recorded, 44.6% was found in RTSR cattle, 0.5% in SR goats, 3.3% in SR sheep, 1.4% in RWAH goats, and 4.2% in RWAH sheep. *Listeria* spp. such as *L. innocua*, *L. ivanovii*, *L. monocytogenes*, *L. welshimeri*, and *L. seeligeri* were identified mainly by cultural methods. The overall *Listeria* spp. prevalence (3.2%) is lower than that of 9.5% (19/200) reported in Zambia (Mpundu et al., 2022).

Based on the findings of Dereje et al. (2018), listeriosis can present in three primary clinical forms: meningoencephalitis, abortion, and septicemia. Most clinical cases are associated with infection caused by *L. monocytogenes* and *L. ivanovii* (Radostits and Done, 2007). Experimental studies have demonstrated that within 24 hours of the onset of bacteremia, *L. monocytogenes* invades the reproductive organs and fetus following ingestion or injection into the bloodstream (Dereje et al., 2018). Abortion results within 5–10 days (Radostits and Done, 2007). Abortions and *Listeria* infections typically appear in tropical

countries in the late winter, early spring, or rainy season (Radostits and Done, 2007). The last trimester of pregnancy is when abortions are most frequently noticed, and when all animals of the herd consume the same batch of tainted silage at the same time, abortion storms might happen (Yaeger, 2007). However, the prevalence of listeriosis in Nigeria in ruminant animals is sparsely reported. Thus, surveillance is necessary for making informed decisions and policies for diagnostics, therapeutics, prevention, and control.

In this study, 14% *Brucella* spp. prevalence was recorded, 11.3% prevalence was found in RTSR cattle, 7.2% in RTSR goats, 8.6% in RTSR sheep, 4.6% in SR cattle, 6.13% in SR goats, 4.4% in SR sheep, 16.7% in RWAH sheep, and 7.7% in RWAH. Notably, a 2.0% (35/2152) mortality/stillbirth rate was estimated. Interestingly, of the 53 articles on *Brucella* spp., 44, 4, and 5 articles were published on RTSR, SR, and RWAH, respectively. The species of *Brucella* identified include *B. abortus*, most often reported, and *B. melitensis*. These species were mostly detected by five methods: RBPT, PCR, culture, c-ELISA, and SAT/SAT-EDTA tests. Specifically, 80% of the literature reviewed on brucellosis used the RBPT method, 45% c-ELISA, 40% SAT/SAT-ELISA, 1% PCR (Yusuf and Abdulrasheed, 2019), and 3% isolation through the conventional method (Bale et al., 2003; Ocholi et al., 2004, 2005; Onoja et al., 2008).

Various studies on animal brucellosis with different prevalence rates have been published worldwide. For example, in Jordan, a prevalence of 6.5% was reported in cattle (Al-Majali et al., 2009), while in Southwestern Ethiopia, the prevalence was found to be 3.1% in cattle (Ibrahim et al., 2010). In Punjab, India, a prevalence of 18.6% was observed in cattle (Aulakh et al., 2008). In Bangladesh, the prevalence rates were 2.66% in cattle, 3.15% in goats, and 2.31% in sheep (Rahman et al., 2011). The high prevalence and coverage in 19/36 states within the six geopolitical zones indicate the high endemicity of animal brucellosis that may be associated with the lack of vaccination in Nigeria and other associated risk factors (Akinyemi et al., 2022).

Animal brucellosis is a major threat to food security. In Nigeria, neither the Federal nor State Ministries of Health pay

brucellosis the attention needed, even though the World Health Organization (WHO) now lists it as one of the top neglected zoonoses (Akinyemi et al., 2022). Infertility, late abortion, retained foetal membranes, and decreased productivity are the main reproductive symptoms of brucellosis in livestock (El-Diasty et al., 2021). Numerous species of animals are susceptible to brucellosis, but particularly susceptible are cattle, sheep, goats, and pigs due to the preferential infection of *Brucella* strains of certain animal species (El-Diasty et al., 2021). The disease is still widely spread in Africa, especially in regions with significant animal populations (McDermott and Arimi, 2002). Usually, brucellosis screenings are not conducted on livestock slaughtered for human consumption at Nigeria's several abattoirs, resulting in underreporting of brucellosis incidence in Nigeria (Akinyemi et al., 2022).

Leptospira interrogans serovar and *Campylobacter* spp. are significant bacterial abortive pathogens reported only in RTSR and RWAH animal statuses. An overall 27% prevalence of *Leptospira* was recorded in this study. Of this, 1.6% prevalence was recorded in RWAH goats, 0.8% in RWAH cattle, 15.5% in RTSR goats, 7.1% in RTSR sheep, and 30.3% in RTSR cattle. *L. interrogans* serovars detected included the following serovars: *hardjo*, *interrogans*, *pomona*, and *icterohaemorrhagiae*. They were diagnosed by IgG/IgM-ELISA, culture, and MAT. Most of the time, a diagnosis depends on identifying antibodies from the diseased animal. Globally, sheep, goats, and cow herds are infected with *Leptospira* serovar *hardjo*.

Like in Nigeria, significant seroprevalence has been reported in cattle (37%) and goats (29%) in India (Vijayachari P et al., 2014). Another study conducted in India using ELISA revealed a lower seroprevalence of 9.11% *Leptospira* serovar *hardjo* in cattle (Pandian et al., 2015). In Iran, a microscopic agglutination test revealed a prevalence of 4.3% in cattle (Haji Hajikolaei et al., 2007). Despite being native to Africa, leptospirosis epidemiology is poorly understood in sub-Saharan Africa (de Vries et al., 2014). In this study, *Leptospira interrogans* serovar was reported by nine reviewed articles in 16.67% (6/36) states of Nigeria, including Sokoto (NW), Kaduna (NW), Rivers state (SS), Borno (NE), Oyo (SW) and Ogun state (SW), on eight animals of RTSR and one of RWAH status (Stephen et al., 2022). This indicates that leptospirosis is an emerging and rapidly spreading disease in Nigeria. Leptospirosis has become recognized as an emerging zoonosis due to the rising incidence of cases worldwide (Vijayachari et al., 2008).

The prevalence of leptospirosis has been reported to significantly reduce reproductive effectiveness, decrease milk production, cause miscarriage, early neonatal death and stillbirth, and retention of foetal membranes in cattle and other ruminants, which result in significant financial losses for the livestock industry (Faine et al., 1999; Walker, 2005; Abiyi et al., 2015). Due to the African free trade policy, numerous animals are currently imported from diverse sources without being tested for leptospirosis. As a result, several pathogenic *Leptospira* serovars are frequently introduced to supplement existing ones (Stephen et al., 2022). Herd abortion and other symptoms of leptospirosis are poorly understood in Nigeria but are typically linked to illnesses such as brucellosis, campylobacteriosis, and listeriosis (Ajayi et al., 2021). In cases where the reason is not immediately apparent, viral agents are frequently blamed. Clinical indications of leptospirosis in ruminants are commonly missed because leptospirosis-causing bacteria are frequently challenging to isolate from diseased animals (Jubril et al., 2011).

The overall prevalence of *Campylobacter* spp. was 13.0% in this study; of this, 19.9% was estimated in RTSR cattle, 3.5% in RTSR sheep (Salihu et al., 2009; Mai et al., 2013), and 13.0% in RWAH cattle (Mshelia et al., 2012), covering only five of the 36 states in Nigeria, all within the northern Nigeria region including Lake Chad basin (NE), Sokoto (NW), Kano (NW), Kaduna (NW) and Adamawa (NE). Interestingly, a high prevalence of *Campylobacter* spp. was reported in both RTSR and RWAH cattle. The detection of campylobacteriosis within two of the northern regions only connotes the endemic nature of this pathogen in the region with the tendency to spread across other regions of the country due to the trans-regional mobility of the nomadic pastoralists in Nigeria.

Various *Campylobacter* species, namely *C. fetus*, *C. coli*,

C. jejuni, *C. laridis*, and *C. lari*, were identified through cultural methods by the studies investigated in this review (Salihu et al., 2009; Mshelia et al., 2012; Mai et al., 2013). This highlighted the significant burden of campylobacteriosis in the country and the subsequent economic losses associated with it. Similarly, 31.1% of *Campylobacter* spp. prevalence was reported from RTSR bovine in Finland (Hakkinen et al., 2007), 51.2% in the US (Englen et al., 2007), and 16.5% in France (Chatre et al., 2010). The detection of *Leptospira interrogans* serovar and *Campylobacter* species in RTSR and RWAH poses a public health threat and is connected with significant economic loss to livestock production (Ezeh et al., 1987; El-Yuguda et al., 2013).

Interestingly, Chlamydia species with an overall prevalence of 3.8% were reported only from one state (Taraba state) (Abubakar, 2015) in North-Eastern Nigeria. It should be noted that it was the only eligible article that conducted a study on RWAH doe and ewe aborted fetuses. Of the overall prevalence, 2.6% were reported in RWAH ewes, and 4.1% in RWAH does. Abubakar (2015) was the first in-depth investigation into *Chlamydomphila abortus* infection in small ruminants on a larger scale in Nigeria. *Chlamydia* was first discovered when an outbreak of sheep abortions at a research farm in Vom, Plateau State, was reported by Okoh (Okoh, 1986). Later, lambs and goats in the city of Maiduguri were shown to have antibodies against *Chlamydomphila abortus*, with seroprevalences of 3.3% in sheep and 3.6% in goats (Amin and Silsmore, 1993). Several studies conducted in different countries have reported varying prevalence rates of *Chlamydomphila abortus*.

For example, in Turkey, a seroprevalence of 5.38% was documented in sheep (Otlu et al., 2007), while in Saudi Arabia, the rates were 7.52% in sheep and 34.50% in goats (Aljumaah, 2012). In Brazil, the prevalence in goats was reported as 50.0% (Santos et al., 2012), and in Lithuania, it was 54.5% in sheep (Bagdonas et al., 2007). In Jordan, prevalence rates of 21.8% in sheep and 11.4% in goats were recorded (Al-Qudah et al., 2004), while in Mexico, the prevalence was 31.1% in healthy sheep and 21.3% in sheep with a history of clinical abortion (Jiménez-Estrada et al., 2008). Comparatively, the prevalence in Nigeria was somewhat similar to the rate of 2.88% reported by Zhao (2012) in China.

The low prevalence of *Chlamydomphila abortus* in Nigeria might be due to the extensive husbandry system of small ruminant production, which is also the most popular practice in northern parts of Nigeria. Furthermore, the lack of data from other regions in Nigeria might be connected to missing research interest. It is known that intensively maintained flocks experience more chlamydial abortions than flocks managed via extensive management systems (Longbottom and Coulter, 2003; Maley et al., 2009; Stuen and Longbottom, 2011). Due to frequent usage of the same lambing/kidding field or pen, whether outdoors or inside, intensive management increases the likelihood of severe environmental contamination by infectious chlamydial elementary bodies. A key factor in the disease's spread is subjecting vulnerable females to such intense illness (Longbottom and Coulter, 2003; Maley et al., 2009; Stuen and Longbottom, 2011). It should be noted that none of the bacterial abortive pathogens have been studied using known molecular epidemiological surveillance tools for possible diagnostic, therapeutic, and prophylactic interventions in this reviewed study.

Hemoprotozoal abortifacient diseases in Nigeria

19.8% of protozoans' abortive prevalence was calculated from data from 8 articles. Of this, 10% of *Neospora* spp. and 14.59% of *T. gondii* prevalence was estimated from 3 and 5 publications, respectively. The prevalence of 10% *Neospora* spp. found in this study is generally lower than that recorded for Iran (32%) (Youssefi et al., 2009), Western Romania (27.7%) (Imre et al., 2012), and Senegal (17.9%) (Kamga-Waladjo et al., 2010). Notably, the prevalence of *N. caninum* in 10% of the herds indicates that neosporosis is endemic in Nigeria and that farmers must be educated. A key strategy for controlling neosporosis is to stop dogs (the definitive host) from contaminating pastures and feeding with waste (Hernandez et al., 2001; Dubey and Schares, 2011).

Interestingly, the 14.59% *T. gondii* prevalence was only recorded from RTSR by five articles (Joshua and Akinwumi,

2003; Kamani et al., 2010a; Onyiche and Ademola, 2015; Ayinmode and Abiola, 2016; Ayinmode et al., 2017). Of this, 14.21% prevalence was estimated in RTSR cattle and 2.55% in RTSR sheep. It threatens the consumer of undercooked, fresh meat and abattoirs (Dubey et al., 2020). Although toxoplasmosis was only recorded in 2/36 states in Nigeria [Oyo state (SW) and Borno state (NE)], there is a possibility of the widespread to other states (Dubey et al., 2020). A lower prevalence of 11.8% (64/541) was recorded in Spain (Mainar et al., 1996). However, a high prevalence of 41.7% in goats and 59.3% in sheep of *T. gondii* were detected in Northern Italy (Giannouli et al., 2010).

Other European countries, including the Netherlands, Portugal, France, Switzerland, Romania, Greece, and Spain, also reported high seroprevalence rates with large differences in *T. gondii* prevalence ranging from 18.5% to 52.8% in goats and from 27.8% to 89% observed in sheep. This disparity in prevalence may be connected to study locations, animal race, husbandry system, and detection methods (Gazzonis et al., 2015).

Mycotic abortifacient diseases in Nigeria

In this study, an overall 28.74% prevalence of abortive mycotic disease was found in SR goats and RWAH goats in Ogun state (SW) and Sokoto state (NW) (Biobaku et al., 2016). However, a lower prevalence of 11.7% was reported in Tanzania (Karimuribo et al., 2006). For clinical mastitis of goats in Nigeria prevalence ranging from 5.7% to 10.7% was recorded (Ameh et al., 1993; Ameh and Tari, 1999; Danmallam et al., 2018; Danmallam and Pimenov, 2019). It should be noted that aspergillosis is one of the most (27.54%) reported abortive diseases in ruminants, particularly in pregnant animals, resulting in significant economic and public health complications (Pal, 2015). Aspergillosis in the tropics might be connected to the indiscriminate feeding habits of ruminants. Therefore, it is suggested that animals raised in the tropics, particularly those in Sub-Saharan Africa, should be closely tended to or managed intensively to prevent mycotoxicosis. The best diet for ruminants is dry, high-quality feed that is not contaminated with moisture, thereby reducing the prevalence of mold development and feed contamination (Pal, 2015).

Risk factors contributing to abortion in Nigeria

Eight significant risk factors were reported to be connected to abortive pathogens in ruminants in this study (Table 6). The most often cited risk factors were age, season, and breed. Ruminants are more prone to contracting *Leptospira*, *Toxoplasma*, and *Anaplasma* species during the rainy season between April and October in Nigeria (Eniolorunda et al., 2008; Kamani et al., 2010a; Pukuma et al., 2011). Ruminants are allegedly more vulnerable to PPRV and *Brucella* species during the dry season between November and March in Nigeria as a result of the high wind speeds associated with this period (Salihu et al., 2009; Victor et al., 2017). The most often cited risk factors for *Brucella*, *Toxoplasma*, PPRV, and BVDV are open grazing, pastoral systems, and semi-intensive husbandry (Kamani et al., 2010a; Sanusi et al., 2016; Victor et al., 2017; Ibrahim et al., 2020).

Non-vaccinated ruminants are more vulnerable to PPRV than vaccinated animals (Victor et al., 2017). It should be noted that the source of the animals for purchase, such as the open markets, is another significant risk factor for *Brucella* (Ibrahim et al., 2020), *C. burnetii* (Cadmus et al., 2020), and *Leptospira interrogans* serovars infection (Pukuma et al., 2011). Certain animal breeds, such as West Africa Dwarf (WAD) breeds of goats, Yankasa varieties of sheep, and Sokoto Gudali (SG) breeds of cattle, were more susceptible to contracting PPRV, *C. burnetii*, and *Brucella* species (Salihu et al., 2009; Victor et al., 2017; Cadmus et al., 2020). Sex (Pukuma et al., 2011; Salihu et al., 2009; Sanusi et al., 2016; Victor et al., 2017; Stephen et al., 2022), history of abortion, and other reproductive diseases (Ibrahim et al., 2020) are additional reported risk factors.

Stillbirth, economic and public health implications of abortive pathogens among ruminants

This study, therefore, confirmed the increasing prevalence of pathogens likely to cause abortion in RTSR and RWAH animals that had no vaccination except PPRV. Furthermore, alarming fetal loss is recorded for diseased goats, sheep, and cattle. An overall 58.95% (619/1050) stillbirth and abortion in sick goats,

49.2% (369/782) in sick sheep, and 6.4% (166/2606) in sick cattle are connected to six abortive diseases, i.e., chlamydiosis, aspergillosis, PPRV infection, brucellosis, campylobacteriosis, and histophillosis.

Notably, 79.1% (291/368) stillbirth and abortion in sick goats was reported due to *Chlamydia* infection, 49.7% (319/642) stillbirth and abortion in sick goats due to PPRV infection, 55.63% (95/184) in sick cattle due to histophillosis, 22.5% (9/40) stillbirth and abortion in sick goat due to aspergillosis and candidiasis. A 13.0% (36/270) abortions and stillbirths were recorded in bovine due to *Campylobacter* infection and 2.0% (35/2152) in cattle due to *Brucella* infection. These diseases are associated with significant economic loss. For instance, around 80% of Nigeria's population relies on 34.5 million cattle, 22.1 million sheep, and 13.9 million goats that produce meat, milk, and other products (Lawal-Adebawale, 2012).

The annual value of cattle production in Nigeria exceeds USD 300 million. A beef farmer in the country spends approximately USD 200 for a typical beef cow (Bourn et al., 1994). Hence, over 1 million cattle in Nigeria may be affected by different abortive pathogens with direct costs of over USD 10 million due to abortion and other reproductive complications caused by the seven pathogens implicated in RWAH (Mshelia et al., 2012), as observed in this current study. Also, 12 of the 16 abortive pathogens recorded are zoonotic agents, while the remaining four viral agents are epizootic. Hence, conducting continuous surveillance of these pathogens to ascertain the true burden of these pathogens in veterinary and human medicine and their economic impacts is important.

Conclusion

This study confirmed the high prevalence of 16 abortive pathogens in ruminants in Nigeria, with brucellosis, PPRV, and anaplasmosis being widely distributed in over 18 of the 36 states of Nigeria. Notably, 12 abortive pathogens (*Anaplasma*, *Leptospira interrogans* ser, *Toxoplasma*, *Listeria*, *C. burnetii*, *Neospora*, *Brucella*, *Campylobacter*, BTV, SBV, BVDV, and PPRV) were recorded from RTSR, 7 (*Listeria*, *Brucella*, *H. somni*, *Aspergillus*, *Candida*, *Anaplasma*, and PPRV) from SR and 7 (*Campylobacter*, *Listeria*, *Chlamydia*, *Leptospira interrogans* ser, *Brucella*, *Aspergillus*, and PPRV) from RWAH.

Specifically, 105, 23, and 12 publications reported abortive pathogens from RTSR, SR, and RWAH, respectively. Several risk factors were identified, such as lack of vaccine for all the abortive pathogens except PPRV, open market systems of procuring animals, husbandry systems, and non-indigenous breeds. All the pathogens except viral abortive agents reported are considered zoonotic public health threats. However, the high prevalence of these pathogens is of great economic significance as they are linked to mortality, stillbirth, infertility, and low productivity of ruminants in Nigeria. Therefore, there is a need for molecular epidemiology, constant surveillance of abortive pathogens, improved husbandry systems, vaccine production, massive administration, and intensive education of pastoralists to improve livestock production in Nigeria.

Article Information

Funding. This research received no external funding.

Conflict of Interest. The authors declare no conflict of interest.

Authors Contributions. Conceptualization, KOA; methodology, SOA and KOA; software, SOA; validation, KOA, SOA, AA and GW, HN; formal analysis, SOA; investigation, SOA.; resources, KOA, SOA, MB, GW, HN; data curation, KOA, SOA; writing—original draft preparation, SOA; writing review and editing, KOA, MB, GW, HN; visualization, KOA, GW, AA, MB, WOS, AOL, and SOA; supervision, KOA. Co-supervision AA; project administration. All authors have read and agreed to the published version of the manuscript.

Publisher's Note. The claims and data contained in this manuscript are solely those of the author(s) and do not represent those of the GMPC publisher, editors, or reviewers. GMPC publisher and the editors disclaim the responsibility for any injury to people or property resulting from the contents of this article.

References

- Abdullahi, S., Idris, Y., Alhaji, M., Kabir, I., Abubakar, S., 2014. Prevalence of haemoparasitic infections in dairy cattle (Friesian breeds) at Nagari integrated dairy farms, Gauta-Nike village, Keffi local government area, Nasarawa state, North Central of Nigeria. *Scientific Journal of Zoology* 3, 17–23. [10.14196/sjz.v3i2.1234](https://doi.org/10.14196/sjz.v3i2.1234).
- Abiayi, E., Inabo, H., Jatau, E., Makinde, A., Sar, T., 2015. Seroprevalence of *Leptospira* antibodies in cattle slaughtered for sale in some North Central States of Nigeria. *Research Journal of Veterinary Sciences* 8, 21–28.
- Abubakar, M., 2015. Detection of *Chlamydophila abortus* antibodies in aborted sheep and goats in Northern and Central parts of Taraba State.
- Adamu, G., Kia, G., Saidu, S., Ejeh, F., 2022. Serosurvey and risk factors of *Coxiella burnetii* in sheep and goats in three agricultural zones of Borno State, Nigeria. *Authorea Preprints*.
- Adamu, S.G., Atsanda, N.N., Tijjani, A.O., Usur, A.M., Sule, A.G., Gulani, I.A., 2016. Epidemiological study of bovine brucellosis in three senatorial zones of Bauchi State, Nigeria. *Veterinary World* 9, 48–52. [10.14202/vetworld.2016.48-52](https://doi.org/10.14202/vetworld.2016.48-52).
- Adamu, S.G., Kabir, J., Umoh, J.U., Raji, M.A., 2018. Seroprevalence of brucellosis and Q fever (Coxiellosis) in cattle herds in Maigana and Birnin Gwari agro-ecological zone of Kaduna State, Nigeria. *Tropical Animal Health and Production* 50, 1583–1589. [10.1007/s11250-018-1598-3](https://doi.org/10.1007/s11250-018-1598-3).
- Adedeji, A.J., Dashe, Y., Akanbi, O.B., Woma, T.Y., Jambol, A.R., Adole, J.A., Bolajoko, M.B., Chima, N., Asala, O., Tekki, I.S., 2019. Co-infection of peste des petits ruminants and goatpox in a mixed flock of sheep and goats in Kanam, North Central Nigeria. *Veterinary Medicine and Science* 5, 412–418.
- Adejoh, V., Pam, V., Uzoigwe, N., Naphtali, R., Yohanna, J., Pam, R., Ombugadu, A., 2019. A survey of ticks and tick-borne parasites in commercial cattle at Lafia, Nasarawa State, Nigeria. *Nigerian Journal Parasitology* 40, 240–244.
- Adelowo, M., 2020. Clinical mastitis in red sokoto goats: Prevalence, presentation, associated fungal agents and their antimycotic susceptibility patterns in a semi-arid region of Nigeria. *Bulletin of Animal Health and Production in Africa* 68, 263–273.
- Ademola, I., Onyiche, T., 2013. Haemoparasites and haematological parameters of slaughtered ruminants and pigs at Bodija abattoir, Ibadan, Nigeria. *African Journal of Biomedical Research* 16, 101–105. URL: <https://www.ajol.info/index.php/ajbr/article/view/109060>.
- Adua, M., Idahor, K., 2017. Haematological evaluation of haemoparasites in cattle and goats slaughtered at Lafia abattoir, Nigeria. *Asian Journal of Biology* 4, 1–5.
- Agada, C., Goden, C., Ogugua, J., 2017. Prevalence of bovine brucellosis and analysis of risk factors in resident cattle herds of Kanke Local Government Area, Plateau State, Nigeria. *Nigerian Veterinary Journal* 38, 104–116. URL: <https://www.ajol.info/index.php/nvj/article/view/164116>.
- Agada, C., Ogugua, A., Anzaku, E., 2018. Occurrence of brucellosis in small ruminants slaughtered in lafia central abattoir, Nasarawa State, Nigeria. *Sokoto Journal of Veterinary Sciences* 16, 16–23. [10.4314/sokjvs.v16i1.3](https://doi.org/10.4314/sokjvs.v16i1.3).
- Agunloye, C., Adeniyi, A., Aremu, O., Oladeji, J., Ojo, M., Ogundipe, G., 2000. An evaluation of an IgG ELISA for the diagnosis of bovine leptospirosis. *Bulletin of Animal Health and Production in Africa* 48, 45–48.
- Ahmad, A., Rabbani, M., Muhammad, K., Shabbir, M.Z., Yaqub, T., Munir, K., Akhter, F., Cepica, A., 2011. Prevalence of bovine viral diarrhoea virus persistency in 12 Holstein cattle dairy herds of Charlottetown, Canada. *Pakistan Journal of Zoology* 43, 255–261.
- Ajayi, O.L., Antia, E.R., Esther, O.O., Oladipo, T.M., Olaniyi, O.M., Awoyomi, O.J., 2021. Detection of pathogenic spp. in goats from slaughterhouses in Southwestern Nigeria using isolation, histochemistry and immunohistochemistry. *Macedonian Veterinary Review* 44, 5–16.
- Akande, F., Takeet, M., Makanju, O., 2010. Haemoparasites of cattle in Abeokuta, South West Nigeria. *Science world journal* 5, 19–21. URL: <https://www.scienceworldjournal.org/article/view/8591/6064>.
- Akinseye, V.O., Adesokan, H.K., Ogugua, A.J., Adedoyin, F.J., Otu, P.I., Kolawole, N.O., Awosanya, E.A., Cadmus, E.O., Cadmus, S.I., Kwaghe, A.V., 2016. Sero-epidemiological survey and risk factors associated with bovine brucellosis among slaughtered cattle in Nigeria. *Onderstepoort Journal of Veterinary Research* 83, 1–7. [10.4102/ojvr.v83i1.1002](https://doi.org/10.4102/ojvr.v83i1.1002).
- Akinyemi, K.O., Fakorede, C.O., Amisu, K.O., Wareth, G., 2022. Human and animal brucellosis in Nigeria: A systemic review and meta-analysis in the last twenty-one years (2001-2021). *Veterinary Sciences* 9. [10.3390/vetsci9080384](https://doi.org/10.3390/vetsci9080384).
- Al-Baroodi, S.Y., 2021. Seroprevalence of Schmallenberg virus infection as emerging disease in cattle in Iraq. *Iraqi Journal of Veterinary Sciences* 35, 495–499.
- Al-Majali, A.M., Talafha, A.Q., Ababneh, M.M., Ababneh, M.M., 2009. Seroprevalence and risk factors for bovine brucellosis in Jordan. *Journal of Veterinary Science* 10, 61–65.
- Al-Qudah, K., Sharif, L., Raouf, R., Hailat, N., Al-Domy, F., 2004. Seroprevalence of antibodies to *Chlamydophila abortus* shown in Awassi sheep and local goats in Jordan. *Veterinarni Medicina* 49, 460.
- Alhaji, N., Wungak, Y., 2013. Epizootiological survey of bovine brucellosis in nomadic pastoral camps in Niger state, Nigeria. *Nigerian Veterinary Journal* 34.
- Alhaji, N.B., Wungak, Y.S., Bertu, W.J., 2016. Serological survey of bovine brucellosis in Fulani nomadic cattle breeds (*Bos indicus*) of North-central Nigeria: Potential risk factors and zoonotic implications. *Acta Tropica* 153, 28–35. [10.1016/j.actatropica.2015.10.003](https://doi.org/10.1016/j.actatropica.2015.10.003).
- Ali, W.H., Osman, N.A., Asil, R.M., Mohamed, B.A., Abdelgadir, S.O., Mutwakil, S.M., Mohamed, N.E.B., 2019. Serological investigations of peste des petits ruminants among cattle in the Sudan. *Tropical Animal Health and Production* 51, 655–659. [10.1007/s11250-018-1740-2](https://doi.org/10.1007/s11250-018-1740-2).
- Aljumaah, R.S., 2012. Serological prevalence of ovine and caprine chlamydophilosis in Riyadh region, Saudi Arabia. *African Journal of Microbiology Research* 6, 2654–2658. [10.5897/AJMR11.1056](https://doi.org/10.5897/AJMR11.1056).
- Ameh, J., Addo, P., Adekeye, J., Gyang, E., 1993. Prevalence of clinical mastitis and of intramammary infections in Nigerian goats. *Preventive Veterinary Medicine* 17, 41–46. [10.1016/0167-5877\(93\)90053-v](https://doi.org/10.1016/0167-5877(93)90053-v).
- Ameh, J., Tari, I., 1999. Observations on the prevalence of caprine mastitis in relation to predisposing factors in Maiduguri. *Small Ruminant Research* 35, 1–5.
- Amin, J., Silsmore, A., 1993. A serological survey of some abortifacient diseases of sheep and goats in the Maiduguri area of Nigeria. *Bulletin of Animal Health and Production in Africa* 41, 123–128.
- Anyanwu, N., Iheanacho, C., Adogo, L., 2016. Parasitological screening of haemo-parasites of small ruminants in karu local government area of Nasarawa State, Nigeria. *British Microbiology Research Journal* 11, 1–8. [10.9734/BMRJ/2016/22358](https://doi.org/10.9734/BMRJ/2016/22358).
- Atsuwe, T., Obisike, V., Azua, E., Imandeh, G., 2019. Prevalence of tick borne haemoparasites in some breeds of cattle and goats slaughtered in some abattoirs within Makurdi, Nigeria. *Journal of Applied Life Sciences International* 21, 1–5. [10.9734/jalsi/2019/v21i430110](https://doi.org/10.9734/jalsi/2019/v21i430110).
- Aulakh, H.K., Patil, P.K., Sharma, S., Kumar, H., Mahajan, V., Sandhu, K.S., 2008. A study on the epidemiology of bovine brucellosis in Punjab (India) using milk-ELISA. *Acta Veterinaria Brno* 77, 393–399.
- Aworh, M.K., Okolocha, E.C., Awosanya, E.J., Fasina, F.O., 2017. Sero-prevalence and intrinsic factors associated with *Brucella* infection in food animals slaughtered at abattoirs in Abuja, Nigeria. *BMC Research Notes* 10, 1–7. [10.1186/s13104-017-2827-y](https://doi.org/10.1186/s13104-017-2827-y).
- Ayinmode, A., 2013. *Neospora caninum* infection in Nigeria: Evidence of occurrence in imported and native breeds of cattle. *Nigerian Veterinary Journal* 34. URL: <https://www.ajol.info/index.php/nvj/article/view/98886/88145>.
- Ayinmode, A., Abiola, J., 2016. Investigating potential sources of toxoplasmosis for humans from slaughtered food animals in Ibadan, Nigeria. *Folia Veterinaria* 60, 34–40. [10.1515/fv-2016-0016](https://doi.org/10.1515/fv-2016-0016).

- Ayinmode, A., Akinseye, V., Schares, G., Cadmus, S., 2017. Serological survey of toxoplasmosis, neosporosis and brucellosis among cattle herds in Oyo State, south-western Nigeria. *African Journal of Infectious Diseases* 11, 95–101. [10.21010/ajid.v11i2.13](https://doi.org/10.21010/ajid.v11i2.13).
- Baba, S., Bobbo, A., Akoma, M., Osiyemi, T., 1994. Slaughterhouse survey for antibodies against selected viruses in ruminant sera in Maiduguri, Borno State, Nigeria. *Der Tropenlandwirt-Journal of Agriculture in the Tropics and Subtropics* 95, 55–62.
- Bagdonas, J., Petkevicius, S., Russo, P., Pepin, M., Salomskas, A., 2007. Prevalence and epidemiological features of ovine enzootic abortion in Lithuania. *Polish Journal of Veterinary Sciences* 10, 239–244. URL: <https://www.ncbi.nlm.nih.gov/pubmed/18198539>.
- Balamurugan, V., Sen, A., Venkatesan, G., Yadav, V., Bhanuprakash, V., Singh, R., 2010. Isolation and identification of virulent peste des petits ruminants viruses from PPR outbreaks in India. *Tropical Animal Health and Production* 42, 1043–1046. [10.1007/s11250-010-9527-0](https://doi.org/10.1007/s11250-010-9527-0).
- Balamurugan, V., Varghese, B., Muthuchelvan, D., SowjanyaKumari, S., Kumar, K.V., Dheeraj, R., Govindaraj, G., Suresh, K., Hemadri, D., Roy, P., 2020. Seroprevalence of peste des petits ruminants in small ruminants in the north eastern region of India. *Veterinaria Italiana* 56, 55–60. [10.12834/VetIt.1985.10631.2](https://doi.org/10.12834/VetIt.1985.10631.2).
- Bale, J., Nuru, S., Addo, P., Adeyinka, I., 2003. Bacteriological investigation of sheep and goats milk for brucellosis in government farms in Northern Nigeria. *Nigerian Journal of Animal Production* 30, 107–116. [10.51791/njap.v30i1.1845](https://doi.org/10.51791/njap.v30i1.1845).
- Bamidele, F., Gidado, S., Edukugho, A., Cadmus, S., 2020. Seroprevalence of brucellosis in abattoir workers and slaughtered cattle in Ilorin metropolis Kwara State Nigeria. *International Journal of Infectious Diseases* 101, 532–533. [10.1016/j.ijid.2020.09.1379](https://doi.org/10.1016/j.ijid.2020.09.1379).
- Banyard, A.C., Parida, S., Batten, C., Oura, C., Kwiatek, O., Libeau, G., 2010. Global distribution of peste des petits ruminants virus and prospects for improved diagnosis and control. *The Journal of General Virology* 91, 2885–2897. [10.1099/vir.0.025841-0](https://doi.org/10.1099/vir.0.025841-0).
- Bashiru, G., Faleke, O.O., Salihu, Garba, H.S., Suleiman, N., Bashir, S.U., Aliyu, M.A., 2013. Seroprevalence of leptospirae in sheep slaughtered at sokoto metropolitan abattoir. *Journal of Veterinary Advances* 2, 26–29. URL: <https://api.semanticscholar.org/CorpusID:54835410>.
- Bello, M.B., Kazeem, H.M., Oladele, S.B., Fatihu, M.Y., Tambuwai, F.M., Jibril, A.H., 2018. Seroprevalence of peste des petits ruminants among unvaccinated small ruminants in Sokoto State, northwestern Nigeria. *Comparative Clinical Pathology* 27, 1–6. [10.1007/s00580-018-2711-8](https://doi.org/10.1007/s00580-018-2711-8).
- Bertu, W., Ajogi, I., Bale, J., Kwaga, J., Ocholi, R., 2010. Seroprevalence of brucellosis in small ruminants in Plateau State, Nigeria. *African Journal of Microbiology Research* 4, 1935–1938.
- Bertu, W., Ocholi, R., Gusi, A., Ngulukun, S., Ducrottoy, M., Moriyon, I., 2020. Isolation of *Brucella* strains in cattle from sedentary and nomadic communities and its public health implication. *International Journal of Infectious Diseases* 101, 533. [10.1016/j.ijid.2020.09.1380](https://doi.org/10.1016/j.ijid.2020.09.1380).
- Biobaku, K.T., Odetokun, I.A., Raji, L.O., Olurode, S.O., Ameen, S.A., 2016. A case report of abortion induced by aspergillus mycotoxicosis in Sokoto Red goat. *Alexandria Journal of Veterinary Sciences* 49, 91–94. [10.5455/ajvs.202495](https://doi.org/10.5455/ajvs.202495).
- Bourn, D., Wint, W., Blench, R., Woolley, E., 1994. Nigerian livestock resources survey. *World Animal Review* 78, 49–58.
- Buhari, H., Saidu, S., Kudi, C., Okolocha, E., Kaltungo, B., 2020. Seroprevalence of *Brucella* infection in small ruminants from two institutional farms and a slaughter slab in Zaria, Nigeria. *Sokoto Journal of Veterinary Sciences* 18, 91–99. [10.4314/sokjvs.v18i2.5](https://doi.org/10.4314/sokjvs.v18i2.5).
- Buhari, H., Sn, A., Mohammed, G., Raji, M., 2016. Serological evaluation of bovine brucellosis in the North Senatorial District of Kaduna State, Nigeria. *International Journal of Veterinary Science* 5, 24–28. URL: <https://www.ijvets.com/pdf-files/Volume-5-no-1-2016/24-28.pdf>.
- Bukar, B.A., El-Yuguda, A.D., Said, L., 2020. The first seroprevalence investigation of peste des petits ruminants virus among Sahel goat in Yobe State, Nigeria. *Asian Journal of Medicine and Health* , 33–3810. [9734/ajmah/2020/v18i430197](https://doi.org/10.9734/ajmah/2020/v18i430197).
- Cadmus, S., Adesokan, H., Adedokun, B., Stack, J., 2010. Seroprevalence of bovine brucellosis in trade cattle slaughtered in Ibadan, Nigeria, from 2004–2006. *Journal of the South African Veterinary Association* 81, 50–53. [10.4102/jsava.v81i1.96](https://doi.org/10.4102/jsava.v81i1.96).
- Cadmus, S., Osikoya, I., Adesokan, H., 2008. Brucellosis in trade cattle in lagos state: An investigation of two abattoirs. *Nigerian Veterinary Journal* 29. URL: <https://www.ajol.info/index.php/nvj/article/view/65151/52847>.
- Cadmus, S., Salam, S.P., Adesokan, H.K., Akporube, K., Ola-Daniel, F., Awosanya, E.J., 2021. Seroprevalence of brucellosis and Q fever infections amongst pastoralists and their cattle herds in Sokoto State, Nigeria. *Plos One* 16, e0254530. [10.1371/journal.pone.0254530](https://doi.org/10.1371/journal.pone.0254530).
- Cadmus, S.I., Akporube, K.A., Ola-Daniel, F., Adelakun, O.D., Akinseye, V.O., 2020. Seroprevalence and associated factors of brucellosis and Q-fever in cattle from Ibarapa area, Oyo State, South-western Nigeria. *The Pan African medical journal* 36, 370. [10.11604/pamj.2020.36.370.24925](https://doi.org/10.11604/pamj.2020.36.370.24925).
- Cadmus, S.I., Alabi, P.I., Adesokan, H.K., Dale, E.J., Stack, J.A., 2013. Serological investigation of bovine brucellosis in three cattle production systems in Yewa Division, south-western Nigeria. *Journal of the South African Veterinary Association* 84, 1–6. [10.4102/jsava.v84i1.217](https://doi.org/10.4102/jsava.v84i1.217).
- Chakrabartty, A., Bhattacharjee, P.K., Sarker, R.R., Rahman, A.K.M.A., Henning, K., Neubauer, H., Rahman, M.S., 2016. Prevalence of *coxiella burnetii* infection in cattle, black Bengal goats and ticks in Bangladesh. *Bangladesh Journal of Veterinary Medicine* 14, 65. [10.3329/bjvm.v14i1.28827](https://doi.org/10.3329/bjvm.v14i1.28827).
- Chatre, P., Haenni, M., Meunier, D., Botrel, M.A., Calavas, D., Madec, J.Y., 2010. Prevalence and antimicrobial resistance of campylobacter jejuni and campylobacter coli isolated from cattle between 2002 and 2006 in France. *Journal of food protection* 73, 825–831. [10.4315/0362-028x-73.5.825](https://doi.org/10.4315/0362-028x-73.5.825).
- Chukwudi, I.C., Mira, F., Ogbu, K.I., Malesa, R.P., Purpari, G., Luka, P.D., Ugochukwu, E.I., Heath, L.E., Guercio, A., Chah, K.F., 2021. Peste des petits ruminants in Nigeria: Identification and variations of lineage II and lineage IV in sheep and goats. *Veterinaria Italiana* 57. [10.12834/VetIt.2390.14842.1](https://doi.org/10.12834/VetIt.2390.14842.1).
- Chukwudi, I.C., Ogbu, K.I., Nwabueze, A.L., Olaolu, O.S., Ugochukwu, E.I., Chah, K.F., 2020. Update on peste des petits ruminants status in south east Nigeria: Serological and farmers' awareness investigation, and potential risk factors. *Tropical Animal Health and Production* 52, 3285–3291. [10.1007/s11250-020-02359-7](https://doi.org/10.1007/s11250-020-02359-7).
- Danmallam, F., Pimenov, N., Ngulukun, S.S., Mwankon, S., 2018. Prevalence and bacterial etiology of mastitis in small ruminants in toro local government area, Bauchi State, Nigeria. *Russian Journal of Agricultural and Socio-Economic Sciences* 79, 341–345. [10.18551/rjoas.2018-07.38](https://doi.org/10.18551/rjoas.2018-07.38).
- Danmallam, F.A., Pimenov, N.V., 2019. Study on prevalence, clinical presentation, and associated bacterial pathogens of goat mastitis in Bauchi, Plateau, and Edo states, Nigeria. *Veterinary World* 12, 638. [10.14202/vetworld.2019.638-645](https://doi.org/10.14202/vetworld.2019.638-645).
- Dauda, G.B., Cheryl, M., Folorunso, O.F., Ighodalo, I., 2015. Abattoir characteristics and seroprevalence of bovine brucellosis in cattle slaughtered at Bodija Municipal Abattoir, Ibadan, Nigeria. *Journal of Veterinary Medicine and Animal Health* 7, 164–168. [10.5897/JVMH2015.0370](https://doi.org/10.5897/JVMH2015.0370).
- David, O.F.S., Goria, K.P., Abraham, D.G.A., 2018. Haemoparasite fauna of domestic animals in plateau state, north central Nigeria. *Bayero Journal of Pure and Applied Sciences* 11, 156–161. [10.4314/bajopas.v11i2.19](https://doi.org/10.4314/bajopas.v11i2.19).
- Dawang, N., Danahap, L., Nash, K., Oluseye, E., Kalejaye, O., 2016. Occurrence of *Brucella* spp in slaughtered sheep and goats in Jos Main Abattior Jos, Plateau State, Nigeria. *African Journal Of Natural Sciences* 17.
- De Vries, A., 2006. Economic value of pregnancy in dairy cattle. *Journal of Dairy Science* 89, 3876–3885. [10.3168/jds.S0022-0302\(06\)72430-4](https://doi.org/10.3168/jds.S0022-0302(06)72430-4).
- Deme, G., Malann, Y., Olanrewaju, C., Lumi, E., 2017. Ticks and tick-borne infections in some livestock slaughtered at Gwagwalada abattoir, federal capital territory, Abuja, Nigeria. *Nigerian Journal of Parasitology* 38, 258. [10.4314/njpar.v38i2.23](https://doi.org/10.4314/njpar.v38i2.23).

- Dereje, T., Benti, D., Feyisa, B., Abiy, G., 2018. Review of common causes of abortion in dairy cattle in Ethiopia. *Journal of Veterinary Medicine and Animal Health* 10, 1–13. [10.5897/JVMAH2017.0639](https://doi.org/10.5897/JVMAH2017.0639).
- Dubey, J.P., Murata, F.H.A., Cerqueira-Cézar, C.K., Kwok, O.C.H., Su, C., 2020. Economic and public health importance of *Toxoplasma gondii* infections in sheep: 2009–2020. *Veterinary Parasitology* 286, 109195. [10.1016/j.vetpar.2020.109195](https://doi.org/10.1016/j.vetpar.2020.109195).
- Dubey, J.P., Schares, G., 2011. Neosporosis in animals- The last five years. *Veterinary Parasitology* 180, 90–108. [10.1016/j.vetpar.2011.05.031](https://doi.org/10.1016/j.vetpar.2011.05.031).
- Duong, M.C., Alenius, S., Huong, L.T.T., Björkman, C., 2008. Prevalence of *Neospora caninum* and bovine viral diarrhoea virus in dairy cows in Southern Vietnam. *The Veterinary Journal* 175, 390–394. [10.1016/j.tvjl.2006.01.016](https://doi.org/10.1016/j.tvjl.2006.01.016).
- Eisawi, N.M., El Hussein, A.R.M., Hassan, D.A., Musa, A.B., Hussien, M.O., Enan, K.A., Bakheit, M.A., 2020. A molecular prevalence survey on *Anaplasma* infection among domestic ruminants in Khartoum State, Sudan. *Tropical Animal Health and Production* 52, 1845–1852. [10.1007/s11250-019-02176-7](https://doi.org/10.1007/s11250-019-02176-7).
- Ekere, S.O., Njoga, E.O., Onunkwo, J.I., Njoga, U.J., 2018. Serosurveillance of *Brucella* antibody in food animals and role of slaughterhouse workers in spread of *Brucella* infection in Southeast Nigeria. *Veterinary World* 11, 1171–1178. [10.14202/vetworld.2018.1171-1178](https://doi.org/10.14202/vetworld.2018.1171-1178).
- El-Diasty, M., El-Said, R., Abdelkhalik, A., 2021. Seroprevalence and molecular diagnosis of sheep brucellosis in Dakahlia governorate, Egypt. *German Journal of Veterinary Research* 1, 34–39. [10.51585/gjvr.2021.0006](https://doi.org/10.51585/gjvr.2021.0006).
- El-Yuguda, A., Abubakar, M., Nabi, A., Andrew, A., Baba, S., 2009a. Outbreak of peste des petits ruminants in an unvaccinated sahel goat farm in Maiduguri, Nigeria. *African Journal of Biomedical Research* 12, 83–87. URL: <https://www.bioline.org.br/pdf?md09015>.
- El-Yuguda, A., Chabiri, L., Adamu, F., Baba, S.S., 2009b. Peste des petits ruminants virus (PPRV) infection among small ruminants slaughtered at the central abattoir, Maiduguri, Nigeria. *Sahel Journal of Veterinary Sciences* 8. URL: <https://www.ajol.info/index.php/sjvs/article/view/53683/0>.
- El-Yuguda, A.D., Saka, S.B., Ambali, A.G., Egwu, G.O., 2013. Seroprevalence of peste des petits ruminants among domestic small and large ruminants in the semi-arid region of north-eastern Nigeria. *Veterinary World* 6, 807. [10.14202/vetworld.2013.807-811](https://doi.org/10.14202/vetworld.2013.807-811).
- Elbers, A.R.W., Loeffen, W.L.A., Quak, S., de Boer-Luijtz, E., van der Spek, A.N., Bouwstra, R., Maas, R., Spierenburg, M.A.H., de Kluijver, E.P., van Schaik, G., van der Poel, W.H.M., 2012. Seroprevalence of Schmallenberg virus antibodies among dairy cattle, the Netherlands, winter 2011–2012. *Emerging Infectious Diseases* 18, 1065–1071. [10.3201/eid1807.120323](https://doi.org/10.3201/eid1807.120323).
- Elelu, N., Ferrolho, J., Couto, J., Domingos, A., Eisler, M.C., 2016. Molecular diagnosis of the tick-borne pathogen *Anaplasma marginale* in cattle blood samples from Nigeria using qPCR. *Experimental & Applied Acarology* 70, 501–510. [10.1007/s10493-016-0081-y](https://doi.org/10.1007/s10493-016-0081-y).
- Englen, M.D., Hill, A.E., Dargatz, D.A., Ladely, S.R., Fedorka-Cray, P.J., 2007. Prevalence and antimicrobial resistance of *Campylobacter* in US dairy cattle. *Journal of Applied Microbiology* 102, 1570–1577. [10.1111/j.1365-2672.2006.03189.x](https://doi.org/10.1111/j.1365-2672.2006.03189.x).
- Eniolorunda, O., Ajado, O., Oyedele, S., 2008. The prevalence and seasonality of ruminant animal diseases in Ijebu-Ode, Ogun State, Nigeria. *Nigerian Journal of Animal Production* 35, 286–292. [10.51791/njap.v35i2.2481](https://doi.org/10.51791/njap.v35i2.2481).
- Ezeh, A., Addo, P., Adesiyun, A., Lawande, R., 1987. Leptospiral antibody responses in four cattle herds in plateau state of Nigeria: A preliminary report. *Bulletin of Animal Health and Production in Africa* 35, 263–265.
- Fadiga, M., Jost, C., Ihedioha, J., 2013. Financial costs of disease burden, morbidity and mortality from priority livestock diseases in Nigeria: Disease burden and cost-benefit analysis of targeted interventions. *International Livestock Research Institute (ILRI) Research Report*.
- Faine, S., Adler, B., Bolin, C., Perolat, P., 1999. "Leptospira" and leptospirosis. *MediSci*. URL: <https://research.monash.edu/en/publications/leptospira-and-leptospirosis>.
- Farouk, U., Bale, J., Kwaga, J., Abdullahi, U., Ibrahim, S., Madobi, I., 2017. Preliminary study on brucellosis in cattle in Jigawa state Nigeria, in: *Proceedings of the 54th Annual Congress of the Nigerian Veterinary Medical Association*. University Press Limited Zaria, Kaduna State, Nigeria, pp. 66–71.
- Gao, S., Luo, J., Du, J., Lang, Y., Cong, G., Shao, J., Lin, T., Zhao, F., Belák, S., Liu, L., 2013. Serological and molecular evidence for natural infection of bactrian camels with multiple subgenotypes of bovine viral diarrhoea virus in western china. *Veterinary Microbiology* 163, 172–176. [10.1016/j.vetmic.2012.12.015](https://doi.org/10.1016/j.vetmic.2012.12.015).
- Gazzonis, A.L., Veronesi, F., Di Cerbo, A.R., Zanzani, S.A., Molineri, G., Moretta, I., Moretti, A., Piergili Fioretti, D., Invernizzi, A., Manfredi, M.T., 2015. *Toxoplasma gondii* in small ruminants in Northern Italy - Prevalence and risk factors. *Annals of Agricultural and Environmental Medicine* 22, 62–68. [10.5604/12321966.1141370](https://doi.org/10.5604/12321966.1141370).
- Gboeloh, L.B., Araka, A.D., 2022. Prevalence of tick-borne haemoparasitic infections in zebu cattle slaughtered in Yenagoa, Bayelsa State. *International Journal of Biological and Pharmaceutical Sciences Archive* 3, 024–030. [10.53771/ijbpsa.2022.3.1.0023](https://doi.org/10.53771/ijbpsa.2022.3.1.0023).
- Giannouli, M., Cuccurullo, S., Crivaro, V., Di Popolo, A., Bernardo, M., Tomasone, F., Amato, G., Brisse, S., Triassi, M., Utili, R., Zarrilli, R., 2010. Molecular epidemiology of multidrug-resistant *Acinetobacter baumannii* in a tertiary care hospital in Naples, Italy, shows the emergence of a novel epidemic clone. *Journal of Clinical Microbiology* 48, 1223–1230. [10.1128/JCM.02263-09](https://doi.org/10.1128/JCM.02263-09).
- Gong, Q.L., Wang, Q., Yang, X.Y., Li, D.L., Zhao, B., Ge, G.Y., Zong, Y., Li, J.M., Leng, X., Shi, K., 2021. Seroprevalence and risk factors of the bluetongue virus in cattle in China from 1988 to 2019: A comprehensive literature review and meta-analysis. *Frontiers in Veterinary Science* 7, 550381. [10.3389/fvets.2020.550381](https://doi.org/10.3389/fvets.2020.550381).
- Haji Hajikolaei, M.R., Ghorbanpour, M., Gharibi, D., Abdollahpour, G., 2007. Serologic study on leptospirosis infection in sheep in Ahvaz, southwestern Iran. *Iranian Journal of Veterinary Research* 8, 333–336. [10.22099/IJVR.2007.10](https://doi.org/10.22099/IJVR.2007.10).
- Hakkinen, M., Heiska, H., Hänninen, M.L., 2007. Prevalence of *Campylobacter* spp. in cattle in finland and antimicrobial susceptibilities of bovine campylobacter jejuni strains. *Applied and Environmental Microbiology* 73, 3232–3238. [10.1128/AEM.02579-06](https://doi.org/10.1128/AEM.02579-06).
- Hansen, P., 2002. Embryonic mortality in cattle from the embryo's perspective. *Journal of Animal Science* 80, E33–E44. [10.2527/animalsci2002.80E-Suppl_2E33x](https://doi.org/10.2527/animalsci2002.80E-Suppl_2E33x).
- Happi, A.N., Buba, D.M., Oluniyi, P.E., Akano, K., 2020a. Detection and identification of blood-borne infections in goats in Nigeria using light microscopy and polymerase chain reaction. *International Journal of Veterinary Science and Research* 6, 090–103. [10.17352/ijvsr](https://doi.org/10.17352/ijvsr).
- Happi, A.N., Osifade, O., Oluniyi, P.E., Ogunro, B.N., 2020b. Comparison of light microscopy and polymerase chain reaction for the detection of haemoparasites in cattle in Nigeria. *Acta Parasitologica* 65, 44–56. [10.2478/s11686-019-00123-y](https://doi.org/10.2478/s11686-019-00123-y).
- Hemida, M.G., Alghadeer, H.M., Alhammedi, M., Ali, S., 2020. Prevalence and molecular characterization of some circulating strains of the peste-des-petits-ruminants virus in Saudi Arabia between 2014–2016. *PeerJ* 8, e9035. [10.7717/peerj.9035](https://doi.org/10.7717/peerj.9035).
- Hernandez, J., Risco, C., Donovan, A., 2001. Association between exposure to *Neospora caninum* and milk production in dairy cows. *Journal of the American Veterinary Medical Association* 219, 632–635. [10.2460/javma.2001.219.632](https://doi.org/10.2460/javma.2001.219.632).
- Herrero, M., Grace, D., Njuki, J., Johnson, N., Enahoro, D., Silvestri, S., Rufino, M.C., 2013. The roles of livestock in developing countries. *Animal* 7 Suppl 1, 3–18. [10.1017/S1751731112001954](https://doi.org/10.1017/S1751731112001954).
- Hossein-Zadeh, N.G., 2013. Effects of main reproductive and health problems on the performance of dairy cows: A review. *Spanish Journal of Agricultural Research* , 718–735. [10.5424/sjar/2013113-4140](https://doi.org/10.5424/sjar/2013113-4140).
- Ibrahim, N., Belihu, K., Lobago, F., Bekana, M., 2010. Seroprevalence of bovine brucellosis and its risk factors in Jimma zone of Oromia Region, South-western Ethiopia. *Tropical Animal Health and Production* 42, 35–40. [10.1007/s11250-009-9382-z](https://doi.org/10.1007/s11250-009-9382-z).
- Ibrahim, Y., Mai, H.M., Kalla, D.J.U., Kabir, J., Nathaniel, J., 2020. Seroprevalence and potential risk factors of bovine brucellosis at the livestock-wildlife interface area of Yankari game

- reserve, Bauchi State, Nigeria. Nigerian Journal of Animal Production 46, 33–44. [10.51791/njap.v46i4.292](https://doi.org/10.51791/njap.v46i4.292).
- Idris Ibrahim, F., 2021. An appraisal of common infectious diseases of small ruminants encountered at zone II veterinary clinic, Potiskum, Yobe State, Nigeria: Economic and public health implications. URL: <http://w.huveta.hu/handle/10832/3103>.
- Imre, K., Morariu, S., Ilie, M.S., Imre, M., Ferrari, N., Genchi, C., Dărăbuș, G., 2012. Serological survey of *Neospora caninum* infection in cattle herds from Western Romania. The Journal of Parasitology 98, 683–685. [10.1645/GE-3023.1](https://doi.org/10.1645/GE-3023.1).
- Jajere, S.M., Atsanda, N.N., Bitrus, A.A., Hamisu, T.M., Ayo, A.O., 2016. Seroprevalence of brucellosis among cattle slaughtered in three municipal abattoirs of nigombe state, northeastern geria. Veterinary World 9, 1082–1086. [10.14202/vetworld.2016.1082-1086](https://doi.org/10.14202/vetworld.2016.1082-1086).
- Jarikre, T.A., Emikpe, B.O., 2017. First report of immunohistochemical detection of peste des petit ruminants, parainfluenza 3 and respiratory syncytial viral antigens in lungs of Nigerian goats. Journal of Immunoassay and Immunochemistry 38, 555–568. [10.1080/15321819.2017.1349669](https://doi.org/10.1080/15321819.2017.1349669).
- Jiménez-Estrada, J.M., Escobedo-Guerra, M.R., Arteaga-Troncoso, G., López-Hurtado, M., De Haro-Cruz, M.D.J., Jiménez, R.M.D.O., Guerra-Infante, F.M., 2008. Detection of *Chlamydia abortus* in sheep (*Ovis aries*) in Mexico. American Journal of Animal and Veterinary Sciences , 91–95URL: <https://thescpub.com/pdf/ajavsp.2008.91.95.pdf>.
- Joshua, R., Akinwumi, K., 2003. Prevalence of antibodies to *Toxoplasma gondii* in four breeds of cattle at Ibadan, Nigeria. Tropical Veterinarian 21, 134–137. [10.4314/tv.v21i3.4533](https://doi.org/10.4314/tv.v21i3.4533).
- Jubril, A.J., Ajayi, O., Ilugbo, M., Olugasa, B., Schobesberger, H., 2011. Isolation and prevalence of pathogenic *Leptospira interrogans* in slaughtered cattle in two abattoirs in southwestern Nigeria, in: Proceedings of the XVth International Congress of the International Society for Animal Hygiene, International Society for Animal Hygiene, Vienna, Austria. pp. 235–237.
- Junaidu, A., Garba, H., 2006. Application of competitive ELISA (CompELISA) rose bengal plate test (RBPT) and serum agglutination test (SAT) for detection of antibodies to *Brucella* infection in slaughter cattle in Sokoto, Nigeria. Sahel Journal of Veterinary Sciences 5, 9–12. URL: <https://www.ajol.info/index.php/sjvs/article/view/41460>.
- Junaidu, A., Salihu, M., Gulumbe, M., 2006. Seroprevalence of brucellosis in sheep in Sokoto city abattoir. Pakistan Journal of Biological Sciences 9, 2696–2698. [10.3923/pjbs.2006.2696.2698](https://doi.org/10.3923/pjbs.2006.2696.2698).
- Junaidu, A.U., Oboegbulem, S., Salihu, M.D., 2011. Serological survey of *Brucella* antibodies in breeding herds. Journal of Microbiology and Biotechnology Research 1, 60–65.
- Kaltungo, B., Saidu, S., Kudi, C., Ismail, M., Jacob, R., Salisu, U., Baba, Y., Buhari, H., 2019. Geo-spatial distribution of *Brucella melitensis* infection in selected local government areas of Katsina and Sokoto States, Nigeria. Nigerian Veterinary Journal 40, 95–103. [10.4314/nvj.v40i2.1](https://doi.org/10.4314/nvj.v40i2.1).
- Kaltungo, B., Saidu, S., Sackey, A., Kazeem, H., 2013. Serological evidence of brucellosis in goats in Kaduna North senatorial district of Kaduna State, Nigeria. International Scholarly Research Notices 2013. [10.1155/2013/963673](https://doi.org/10.1155/2013/963673).
- Kaltungo, B.Y., Saidu, S.N.A., Sackey, A.K.B., Kazeem, H.M., 2015. Sero-prevalence of brucellosis in sheep in north senatorial district of NKaduna State, igeria. Asian Pacific Journal of Tropical Disease 5, 163–168. [10.1016/S2222-1808\(14\)60646-0](https://doi.org/10.1016/S2222-1808(14)60646-0).
- Kamani, J., Irene, S., Qasim, A.M.M.M., Olubade, T.A., Abasiama, M.S., Gajibo, A., Balami, P.U., Shands, M., Harrus, S., 2022a. Nucleotide sequence types (ntSTs) of *Anaplasma marginale* in cattle in Nigeria based on the major surface protein 5 (msp5) gene. Acta Tropica 233, 106544. [10.1016/j.actatropica.2022.106544](https://doi.org/10.1016/j.actatropica.2022.106544).
- Kamani, J., Mani, A.U., Egwu, G.O., 2010a. Seroprevalence of *Toxoplasma gondii* infection in domestic sheep and goats in niborno state, geria. Tropical Animal Health and Production 42, 793–797. [10.1007/s11250-009-9488-3](https://doi.org/10.1007/s11250-009-9488-3).
- Kamani, J., Sannusi, A., Egwu, O., Dogo, G., Tanko, T., Kemza, S., Tafarki, A., Gbise, D., 2010b. Prevalence and significance of haemoparasitic infections of cattle in North- Central, Nigeria. Veterinary World 3, 445. [10.5455/vetworld.2010.445-448](https://doi.org/10.5455/vetworld.2010.445-448).
- Kamani, J., Schaer, J., Umar, A.G., Pilarshimwi, J.Y., Bukar, L., González-Miguel, J., Harrus, S., 2022b. Molecular detection and genetic characterization of *Anaplasma marginale* and *Anaplasma platys* in cattle in Nigeria. Ticks and Tick-borne Diseases 13, 101955. [10.1016/j.ttbdis.2022.101955](https://doi.org/10.1016/j.ttbdis.2022.101955).
- Kamga-Waladjo, A.R., Gbati, O.B., Kone, P., Lapo, R.A., Chatagnon, G., Bakou, S.N., Pangui, L.J., Diop, P.E.H., Akakpo, J.A., Tainturier, D., 2010. Seroprevalence of *Neospora caninum* antibodies and its consequences for reproductive parameters in dairy cows from Dakar-Senegal, West Africa. Tropical Animal Health and Production 42, 953–959. [10.1007/s11250-009-9513-6](https://doi.org/10.1007/s11250-009-9513-6).
- Karimuribo, E.D., Fitzpatrick, J.L., Bell, C.E., Swai, E.S., Kambarage, D.M., Ogden, N.H., Bryant, M.J., French, N.P., 2006. Clinical and subclinical mastitis in smallholder dairy farms in Tanzania: Risk, intervention and knowledge transfer. Preventive Veterinary Medicine 74, 84–98. [10.1016/j.prevetmed.2006.01.009](https://doi.org/10.1016/j.prevetmed.2006.01.009).
- Lawal, A., Lasisi, O., Emikpe, B., Ogundipe, G., 2011. Outbreak of peste des petits ruminants in West African Dwarf goats in Eruwa, Southwestern Nigeria. Nigerian Veterinary Journal 32. URL: <https://www.ajol.info/index.php/nvj/article/view/85619>.
- Lawal, N., Egwu, G., Tambuwal, F., Junaidu, A., Abubakar, M., Magaji, A., Rabi'u, M., Saulawa, M., Mamuda, A., Jibrin, M., 2012. Prevalence of *Brucella abortus* antibodies in bovine serum from Gusau Modern Abattoir, Zamfara state, Nigeria. Scientific Journal of Microbiology 1, 91–96.
- Lawal-Adebowale, O., 2012. Dynamics of ruminant livestock management in the context of the Nigerian agricultural system, in: Javed, K. (Ed.), Livestock Production. InTech, p. 160. [10.5772/52923](https://doi.org/10.5772/52923).
- Longbottom, D., Coulter, L.J., 2003. Animal chlamydioses and zoonotic implications. Journal of Comparative Pathology 128, 217–244. [10.1053/jcpa.2002.0629](https://doi.org/10.1053/jcpa.2002.0629).
- Lorusso, V., Wijnveld, M., Majekodunmi, A.O., Dongkum, C., Fajinmi, A., Dogo, A.G., Thrusfield, M., Mugenyi, A., Vau-mourin, E., Igweh, A.C., Jongejan, F., Welburn, S.C., Picozzi, K., 2016. Tick-borne pathogens of zoonotic and veterinary importance in Nigerian cattle. Parasites & Vectors 9, 217. [10.1186/s13071-016-1504-7](https://doi.org/10.1186/s13071-016-1504-7).
- Luka, P.D., Erume, J., Mwiine, F.N., Ayebazibwe, C., Shamaki, D., 2011. Molecular characterization and phylogenetic study of peste des petits ruminants viruses from north central states of Nigeria. BMC Veterinary Research 7, 1–7. [10.1186/1746-6148-7-32](https://doi.org/10.1186/1746-6148-7-32).
- Luther, N., Umoh, J., Majiyagbe, K., Shamaki, D., Nwosuh, D., Dogo, G., 2005. Studies on the prevalence of antibodies to peste des petits ruminants virus (PPRV) among goats in Bauchi state. Nigerian Veterinary Journal 27, 17–22. URL: <https://www.ajol.info/index.php/nvj/article/view/3500>.
- Magadu, R., Thompson, P.N., 2023. Seroprevalence and factors associated with *Coxiella burnetii* exposure in goats in moretele. The Onderstepoort Journal of Veterinary Research 90, e1–e7. [10.4102/ojvr.v90i1.2071](https://doi.org/10.4102/ojvr.v90i1.2071).
- Mai, H.M., Irons, P.C., Kabir, J., Thompson, P.N., 2012. A large seroprevalence survey of brucellosis in cattle herds under diverse production systems in northern Nigeria. BMC Veterinary Research 8, 1–14. [10.1186/1746-6148-8-144](https://doi.org/10.1186/1746-6148-8-144).
- Mai, H.M., Irons, P.C., Kabir, J., Thompson, P.N., 2013. Herd-level risk factors for campylobacter fetus infection, *Brucella* seropositivity and within-herd seroprevalence of brucellosis in cattle in northern Nigeria. Preventive Veterinary Medicine 111, 256–267. [10.1016/j.prevetmed.2013.05.016](https://doi.org/10.1016/j.prevetmed.2013.05.016).
- Mainar, R.C., de la Cruz, C., Asensio, A., Domínguez, L., Vázquez-Boland, J.A., 1996. Prevalence of agglutinating antibodies to *Toxoplasma gondii* in small ruminants of the Madrid region, Spain, and identification of factors influencing seropositivity by multivariate analysis. Veterinary Research Communications 20, 153–159. [10.1007/BF00385636](https://doi.org/10.1007/BF00385636).
- Maley, S.W., Livingstone, M., Rodger, S.M., Longbottom, D., Buxton, D., 2009. Identification of *Chlamydia abortus* and the development of lesions in placental tissues of experimentally infected sheep. Veterinary Microbiology 135, 122–127. [10.1016/j.vetmic.2008.09.031](https://doi.org/10.1016/j.vetmic.2008.09.031).
- Mantip, S., Sigismeau, A., Shamaki, D., Woma, T.Y., Kwiatek, O., Libeau, G., Farougou, S., Bataille, A., 2022. Molecular epidemiology of peste des petits ruminants virus in Nigeria: An

- update. *Transboundary and Emerging Diseases* 69, 1634–1640. [10.1111/tbed.14073](https://doi.org/10.1111/tbed.14073).
- Maurice, N.A., 2013. Bacteriological and serological studies of bovine brucellosis in Obudu cattle ranch, cross river state. Thesis. Master. Department Of Veterinary Microbiology, Ahmadu Bello University, Zaria, Kaduna State, Nigeria. URL: <https://kubanni.abu.edu.ng/items/feb069f4-a3c5-464a-87fc-7c35e8c09dfc>.
- Mbuk, E., Ajogi, I., Bale, J., Umoh, J., 2011. Prevalence of *Brucella* antibodies in migratory fulani cattle herds in Kaduna State, Nigeria. *Nigerian Veterinary Journal* 32, 26–29. [10.4314/nvj.v32i1.68999](https://doi.org/10.4314/nvj.v32i1.68999).
- McDermott, J.J., Arimi, S., 2002. Brucellosis in sub-Saharan Africa: Epidemiology, control and impact. *Veterinary Microbiology* 90, 111–134. [10.1016/S0378-1135\(02\)00249-3](https://doi.org/10.1016/S0378-1135(02)00249-3).
- Mertens, P.P.C., Maan, N.S., Belaganahalli, M.N., Singh, K.P., Nomikou, K., Maan, S., 2013. Full genome sequence of a western reference strain of bluetongue virus serotype 16 from Nigeria. *Genome Announcements* 1. [10.1128/genomeA.00684-13](https://doi.org/10.1128/genomeA.00684-13).
- Mohammed, F., Ibrahim, S., Musa, G., Kaltungo, B., Danbirni, S., Kwaga, J., 2020. *Brucella* infection in migratory cattle herds in Jigawa State Nigeria: A cross-sectional study. *Sokoto Journal of Veterinary Sciences* 18, 191–194. [10.4314/sokjvs.v18i4.2](https://doi.org/10.4314/sokjvs.v18i4.2).
- Mohammed, F.U., Ibrahim, S., Ajogi, I., Olaniyi, B.J., 2011. Prevalence of bovine brucellosis and risk factors assessment in cattle herds in Jigawa State. *International Scholarly Research Notices* 2011. [10.5402/2011/132897](https://doi.org/10.5402/2011/132897).
- Moher, D., Shamseer, L., Clarke, M., Ghersi, D., Liberati, A., Petticrew, M., Shekelle, P., Stewart, L.A., Group, P.P., 2015. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. *Systematic Reviews* 4, 1. [10.1186/2046-4053-4-1](https://doi.org/10.1186/2046-4053-4-1).
- Mpundu, P., Muma, J.B., Mukumbuta, N., Mukubesa, A.N., Muleya, W., Kapila, P., Hang'ombe, B.M., Munyeme, M., 2022. Isolation, discrimination, and molecular detection of *Listeria* species from slaughtered cattle in Namwala District, Zambia. *BMC Microbiology* 22, 160. [10.1186/s12866-022-02570-6](https://doi.org/10.1186/s12866-022-02570-6).
- Mshelia, G.D., Amin, J.D., Egwu, G.O., Woldehiwet, Z., Murray, R.D., 2012. The prevalence of bovine venereal campylobacteriosis in cattle herds in the Lake Chad basin of Nigeria. *Tropical Animal Health and Production* 44, 1487–1489. [10.1007/s11250-012-0092-6](https://doi.org/10.1007/s11250-012-0092-6).
- Muse, E.A., Karimuribo, E.D., Gitao, G.C., Misinzo, G., Mellau, L.S.B., Msoffe, P.L.M., Swai, E.S., Albano, M.O., 2012. Epidemiological investigation into the introduction and factors for spread of peste des petits ruminants, southern Tanzania. *The Onderstepoort Journal of Veterinary Research* 79, 457. [10.4102/ojvr.v79i2.457](https://doi.org/10.4102/ojvr.v79i2.457).
- Neils, J., Abdu, P., Sackey, A., 2011. Some reproductive disorders of the indigenous sheep and goats in Zaria, Northern Nigeria. *Nigerian Veterinary Journal* 30. [10.4314/nvj.v30i1.65160](https://doi.org/10.4314/nvj.v30i1.65160).
- Ngbede, E., Raji, M., Kwanashie, C., Okolocha, E., 2013. Serosurvey of *Leptospira* spp serovar hardjo in cattle from zaria, Nigeria. *Revue de Medecine Veterinaire* 164, 85–89.
- Ngbede, E.O., Raji, M.A., Kwanashie, C.N., Okolocha, E.C., Momoh, A.H., Adole, E.B., Giegbefumwen, A.O., Omontese, B., Okoro, L.K., Ehwe, M., 2012. Risk practices and awareness of leptospirosis in an abattoir in northwestern Nigeria. *Scientific Journal of Veterinary Advances* 1, 65–69.
- Niaz, S., Ur Rahman, Z., Ali, I., Cossío-Bayúgar, R., Amaro-Estrada, I., Alanazi, A.D., Khattak, I., Zeb, J., Nasreen, N., Khan, A., 2021. Molecular prevalence, characterization and associated risk factors of *Anaplasma* spp. and *Theileria* spp. in small ruminants in Northern Pakistan. *Parasite* 28, 3. [10.1051/parasite/2020075](https://doi.org/10.1051/parasite/2020075).
- Njiro, S.M., Kidanemariam, A.G., Tsotetsi, A.M., Katsande, T.C., Mnisi, M., Lubisi, B.A., Potts, A.D., Baloyi, F., Moyo, G., Mpfu, J., Kalake, A., Williams, R., 2011. A study of some infectious causes of reproductive disorders in cattle owned by resource-poor farmers in Gauteng Province, South Africa. *Journal of the South African Veterinary Association* 82, 213–218. [10.4102/jsava.v82i4.76](https://doi.org/10.4102/jsava.v82i4.76).
- Nwobodo, H., Ezeifeke, G., Ezejiyor, C., Onyianta, O., 2013. Seroprevalence of peste des petits ruminants among goats and sheep in Enugu State of Nigeria. *Bulletin of Animal Health and Production in Africa* 61, 613–616. URL: <https://www.ajol.info/index.php/bahpa/article/view/105298>.
- Ocholi, R., Kwaga, J., Ajogi, I., Bale, J., 2004. Phenotypic characterization of *Brucella* strains isolated from livestock in Nigeria. *Veterinary Microbiology* 103, 47–53. [10.1016/j.vetmic.2004.06.012](https://doi.org/10.1016/j.vetmic.2004.06.012).
- Ocholi, R., Kwaga, J., Ajogi, I., Bale, J., 2005. Abortion due to *Brucella abortus* in sheep in Nigeria. *Scientific and Technical Review. International Office of Epizootics* 24, 973–980. URL: <https://pubmed.ncbi.nlm.nih.gov/16642768/>.
- Odugbo, M.O., Ogunjumo, S.O., Chukwukere, S.C., Kumbish, P.R., Musa, A., Ekundayo, S.O., Okewole, P.A., Nwankpa, N.D., Itodo, A.E., Haruna, G., 2009. The first report of *Histophilus somni* pneumonia in Nigerian dairy cattle. *The Veterinary Journal* 181, 340–342. [10.1016/j.tvjl.2008.03.007](https://doi.org/10.1016/j.tvjl.2008.03.007).
- Ogugua, A.J., Akinseye, V.O., Ayoola, M.C., Oyesola, O.O., Shima, F.K., Tijjani, A.O., Musa, A.N.A., Adesokan, H.K., Perrett, L., Taylor, A., Stack, J.A., Moriyon, I., Cadmus, S.I.B., 2014. Seroprevalence and risk factors of brucellosis in goats in selected states in Nigeria and the public health implications. *African Journal of Medicine and Medical Sciences* 43, 121–129. URL: <https://www.ncbi.nlm.nih.gov/pubmed/26689681>.
- Ogugua, A.J., Akinseye, V.O., Cadmus, E.O., Jolaoluwa Awosanya, E.A., Alabi, P.I., Idowu, O.S., Akinade, S.A., Dale, E.J., Perrett, L., Taylor, A., Ignocio, M., Cadmus, S.I.B., 2018. Prevalence and risk factors associated with bovine brucellosis in herds under extensive production system in southwestern Nigeria. *Tropical Animal Health and Production* 50, 1573–1582. [10.1007/s11250-018-1597-4](https://doi.org/10.1007/s11250-018-1597-4).
- Okoh, A.E., 1986. Enzootic abortion of ewes in Nigeria: An investigation into the naturally occurring disease in a research animal facility. *Revue d'Elevage et de Medecine Veterinaire des Pays Tropicaux* 39, 181–184. URL: <https://www.ncbi.nlm.nih.gov/pubmed/3589063>.
- Okoli, I., 2003. Incidence and modulating effects of environmental factors on trypanosomosis, peste des petit ruminants (PPR) and bronchopneumonia of West African dwarf goats in Imo state, Nigeria. *Livestock Research for Rural Development* 15, 112–119. URL: <https://www.lrrd.cipav.org.co/lrrd15/9/okoli159.htm>.
- Okwelum, N., Iposi, S.O., Sowande, O.S., Smith, O.F., Takeet, M.I., 2021. Epidemiological survey of haemoparasitic infection in trade cattle at slaughter in Lafenwa abattoir, Ogun State, Nigeria. *Nigerian Journal of Animal Production* 41, 187–195. [10.51791/njap.v41i2.803](https://doi.org/10.51791/njap.v41i2.803).
- Oladimeji, O.D., Kelechi, A.I., 2019. Presence of bluetongue virus antibodies in cattle and sheep in Ogun and Osun States, Nigeria. *International Journal of Tropical Disease & Health* , 1–10. [10.9734/ijtdh/2019/v38i330187](https://doi.org/10.9734/ijtdh/2019/v38i330187).
- Olatunde, O.H., Cornelius, J.O., Peter, A.O., Jessica, A.B., Ayobamidele, O.F., 2014. Evaluation of hemoparasites in trade cattle slaughtered in Jos Abattoir, Plateau State-Nigeria. *International Journal of Livestock Research* 4, 113–120. URL: <https://api.semanticscholar.org/CorpusID:70612842>.
- Olorunshola, I., Daodu, O., Kolapo, T., Fadunsin, S., Sanda, I., Folaranmi, D., 2020. The prevalence of spirochaetes and other haemoparasites in small ruminants slaughtered at abattoirs in Ilorin Kwara State, Nigeria. *Alexandria Journal of Veterinary Sciences* 65, 1. [10.5455/ajvs.65780](https://doi.org/10.5455/ajvs.65780).
- Olufemi, O.T., Dantala, D.B., Shinggu, P.A., Dike, U.A., Otolorin, G.R., Nwuku, J.A., Baba-Onoja, E.B.T., Jatau, T.D., Amama, F.I., 2018. Seroprevalence of brucellosis and associated risk factors among indigenous breeds of goats in Wukari, Taraba State, Nigeria. *Journal of Pathogens* 2018, 5257926. [10.1155/2018/5257926](https://doi.org/10.1155/2018/5257926).
- Oluwayelu, D., Meseko, C., Adebisi, A., 2015. Serological screening for Schmallenberg virus in exotic and indigenous cattle in Nigeria. *Sokoto Journal of Veterinary Sciences* 13, 14–18. [10.4314/sokjvs.v13i3.3](https://doi.org/10.4314/sokjvs.v13i3.3).
- Oluwayelu, D.O., Olatoye, O., Akanbi, M., Hoffmann, B., 2011. Re-emergence of bluetongue virus infection in Oyo State, Nigeria. *Journal of Commonwealth Veterinary Association* 27, 237–241. URL: <https://www.cabdirect.org/cabdirect/abstract/20143268774>.

- Onoja, I., Ajani, A., Mshelia, W., Andrew, A., Ogunkoya, A., Achi, C., Sambo, K., 2008. Brucellosis outbreak in a flock of seventeen sheep in Zaria. *Sokoto Journal of Veterinary Sciences* 7. URL: <https://www.ajol.info/index.php/sokjvs/article/view/72756>.
- Onyiche, T.E., Ademola, I.O., 2015. Seroprevalence of anti-*Toxoplasma gondii* antibodies in cattle and pigs in Ibadan, Nigeria. *Journal of Parasitic Diseases* 39, 309–314. [10.1007/s12639-013-0350-1](https://doi.org/10.1007/s12639-013-0350-1).
- Onyiche, T.E., Mofokeng, L.S., Thekisoe, O., MacLeod, E.T., 2022. Molecular survey for tick-borne pathogens and associated risk factors in sheep and goats in Kano Metropolis, Nigeria. *Veterinary Parasitology, Regional Studies and Reports* 33, 100753. [10.1016/j.vprsr.2022.100753](https://doi.org/10.1016/j.vprsr.2022.100753).
- Oruene, I.S., Bekwele, B.B., 2020. Incidence of leptospirosis in household goats in some villages in Rivers State. *International Journal of Innovative Research and Advanced Studies* 7, 1–4. URL: <https://www.ijiras.com/2020/Vol.7-Issue.5/paper.1.pdf>.
- Osue, H., Okoh, K., Abdullahi, M., Ahmed, D., 2022. Haematophagous flies, haemoparasites and ecological variables impinging livestock health in three private farms within southern parts of Kano State, Nigeria. *Science World Journal* 17, 368–374. URL: <https://www.ajol.info/index.php/swj/article/view/236026>.
- Otlu, S., ŞAHİN, M., Unver, A., Celebi, O., 2007. Detection of *Brucella melitensis* and *Chlamydomphila abortus* antibodies. *Bulletin of the Veterinary Institute in Pulawy* 51, 493–495.
- Owai, P.U., 2007. Pest of small ruminants as a major constraint to small ruminant production in Cross River State, Nigeria. *Journal of Food, Agriculture and Environment* 5, 102. [10.1234/4.2007.740](https://doi.org/10.1234/4.2007.740).
- Pal, M., 2015. Growing role of fungi in mycotic abortion of domestic animal. *Journal of Bacteriology and Mycology* 2, 1009. URL: <https://mail.austinpublishinggroup.com/bacteriology/fulltext/bacteriology-v2-id1009.pdf>.
- Pam, V., Ogbu, K., Igeh, C., Bot, C., Vincent, G., 2013. The occurrence of gastrointestinal and haemo parasites of cattle in Jos of Plateau State, Nigeria. *Journal of Animal Science Advances* 3, 97–102. URL: <https://catalog.ihnsn.org/citations/64328>.
- Pandian, S.J., Ray, P.K., Chandran, P.C., Kumar, M., 2015. Seroprevalence of *Brucella abortus* and *Leptospira hardjo* in cattle. *Veterinary World* 8, 217–220. [10.14202/vetworld.2015.217-220](https://doi.org/10.14202/vetworld.2015.217-220).
- Parthiban, S., Malmarugan, S., Murugan, M.S., Johnson Rajeswar, J., Pothiappan, P., 2015. Review on emerging and reemerging microbial causes in bovine abortion. *International Journal of Nutrition and Food Sciences* 4, 1. [10.11648/j.ijnfs.s.2015040401.11](https://doi.org/10.11648/j.ijnfs.s.2015040401.11).
- Parvizi, O., Akinyemi, K., Roesler, U., Neubauer, H., Mertens-Scholz, K., 2020. Retrospective study of small ruminants in countries of North Africa and the Middle East. *Scientific and Technical Review (International Office of Epizootics)* 39, 1053–1068. [10.20506/rst.39.3.3196](https://doi.org/10.20506/rst.39.3.3196).
- Peter, A.T., 2000. Abortions in dairy cows: New insights and economic impact. *Proceedings of Western Canadian Dairy Seminar, Red Deer, Alberta, Canada. Advanced Dairy Technology* 12, 233–244.
- Peter, I., El-Yuguda, A., Mshelia, G., 2016. Detection of IgG antibodies to bovine viral diarrhoea virus in domestic ruminants in Maiduguri, Nigeria. *Nigerian Veterinary Journal* 37, 217–222. URL: <https://www.ajol.info/index.php/nvj/article/view/163812>.
- Peter, I.D., 2015. A retrospective study of small ruminant diseases identified at the state veterinary hospital Maiduguri, Nigeria. *Journal of Animal Health and Production* 3, 88–93. [10.14737/journal.jahp/2015/3.4.88.93](https://doi.org/10.14737/journal.jahp/2015/3.4.88.93).
- Pukuma, S.M., James-Rugu, N.N., Sale, M., 2011. A study on tick-borne infections of cattle in Yola locality of Adamawa State. *African Journal of Agricultural Research* 6, 6208–6211. [10.5897/AJAR10.236](https://doi.org/10.5897/AJAR10.236).
- Quinn, P.J., Markey, B.K., Carter, M.E., Donnelly, W.J., Leonard, F.C., 2001. *Veterinary Microbiology and Microbial Disease*. 1 ed., Wiley-Blackwell.
- Quinn, P.J., Markey, B.K., Leonard, F.C., Hartigan, P., Fanning, S., Fitzpatrick, E.S., 2011. *Veterinary Microbiology and Microbial Disease*. 2 ed., Wiley.
- Radostits, O.M., Done, S.H., 2007. *Veterinary Medicine: A textbook of the diseases of cattle, horses, sheep, pigs and goats (Radostits, Veterinary Medicine)*. 10 ed., Saunders Ltd., New York.
- Rahman, M., Faruk, M., Her, M., Kim, J., Kang, S., Jung, S., 2011. Prevalence of brucellosis in ruminants in Bangladesh. *Veterinárni Medicína* 56, 379–385. [10.17221/1555-VETMED](https://doi.org/10.17221/1555-VETMED).
- Salihu, M.D., Abdulkadir, J.U., Oboegbulem, S.I., Egwu, G.O., Magaji, A.A., Lawal, M., Hassan, Y., 2009. Isolation and prevalence of *Campylobacter* species in cattle from Sokoto state, Nigeria. *Vet. Ital* 45, 501–505. URL: <https://pubmed.ncbi.nlm.nih.gov/20391414/>.
- Sani, M., Amanloo, H., 2007. Heat stress effect on open days in holstein dairy cattle in yazd province, Iran. *The 3rd Congress of Animal Science, Mashhad, Iran*.
- Santos, C.S., Piatti, R.M., Azevedo, S.S., Alves, C.J., Higino, S.S., Silva, M.L., Brasil, A.W., Gennari, S.M., 2012. Seroprevalence and risk factors associated with *Chlamydomphila abortus* infection in dairy goats in the Northeast of Brazil. *Pesquisa Veterinária Brasileira* 32, 1082–1086. [10.1590/S0100-736X2012001100002](https://doi.org/10.1590/S0100-736X2012001100002).
- Sanusi, M.B., Aminu, I.D., Umar, M.C., Mikailu, B.A., Abdurrahman, H.J., Aghedo, F., 2016. Detection of antibodies to bovine viral diarrhoea virus in cattle presented for slaughter at Sokoto Metropolitan abattoir, Nigeria. *Journal of Veterinary Medicine and Animal Health* 8, 11–14. [10.5897/JVMAH2015.0445](https://doi.org/10.5897/JVMAH2015.0445).
- Sen, A., Saravanan, P., Balamurugan, V., Rajak, K.K., Sudhakar, S.B., Bhanuprakash, V., Parida, S., Singh, R.K., 2010. Vaccines against peste des petits ruminants virus. *Expert Review of Vaccines* 9, 785–796. [10.1586/erv.10.74](https://doi.org/10.1586/erv.10.74).
- Shu'aibu, G., Kabir, J., Umoh, J., Raji, M., Tijjani, A., Umaru, G., 2018. Seroprevalence of brucellosis in sheep in maigana and birnin gwari agro-ecological zones of Kaduna State, Nigeria. *Nigerian Veterinary Journal* 39, 240–249. [10.4314/nvj.v39i3.7](https://doi.org/10.4314/nvj.v39i3.7).
- Shyaka, A., Ugirabe, M.A., Wensman, J.J., 2021. Serological evidence of exposure to peste des petits ruminants in small ruminants in Rwanda. *Frontiers in Veterinary Science* 8, 651978. [10.3389/fvets.2021.651978](https://doi.org/10.3389/fvets.2021.651978).
- Sibhat, B., Ayelet, G., Gebremedhin, E.Z., Skjerve, E., Asmare, K., 2018. Seroprevalence of schmallenberg virus in dairy cattle in Ethiopia. *Acta Tropica* 178, 61–67. [10.1016/j.actatropica.2017.10.024](https://doi.org/10.1016/j.actatropica.2017.10.024).
- Siddaramppa, S., Inzana, T.J., 2004. *Haemophilus somnus* virulence factors and resistance to host immunity. *Animal Health Research Reviews* 5, 79–93. [10.1079/ahr200466](https://doi.org/10.1079/ahr200466).
- Solis-Calderon, J.J., Segura-Correa, V.M., Segura-Correa, J.C., 2005. Bovine viral diarrhoea virus in beef cattle herds of Yucatan, Mexico: Seroprevalence and risk factors. *Preventive Veterinary Medicine* 72, 253–262. [10.1016/j.prevetmed.2005.06.004](https://doi.org/10.1016/j.prevetmed.2005.06.004).
- St, K., Alenius, S., 2012. BVDV control and eradication in Europe—An update. *The Japanese Journal of Veterinary Research* 60 Suppl, S31–9. URL: <https://www.ncbi.nlm.nih.gov/pubmed/22458198>.
- Stephen, J., Paul, B.T., Ma'aruf, M.I., Dawurung, J., Bukar, M.M., Mshelia, G.D., 2022. Serological evidence of *Leptospira hardjo* antibodies and the incidence of reproductive disorders in selected smallholder cattle and goat farms from Maiduguri, Nigeria. *Journal of Advanced Veterinary Research* 12, 1–5. URL: <https://advetresearch.com/index.php/AVR/article/view/811>.
- Stuen, S., Longbottom, D., 2011. Treatment and control of chlamydial and rickettsial infections in sheep and goats. *The Veterinary Clinics of North America. Food Animal Practice* 27, 213–233. [10.1016/j.cvfa.2010.10.017](https://doi.org/10.1016/j.cvfa.2010.10.017).
- Taylor, W.P., Okeke, A.N., Shidali, N.N., 1977. Prevalence of bovine virus diarrhoea and infectious bovine rhinotracheitis antibodies in Nigerian sheep and goats. *Tropical Animal Health and Production* 9, 171–175. [10.1007/BF02236592](https://doi.org/10.1007/BF02236592).
- Ukwueze, K.O., Ishola, O.O., Dairo, M.D., Awosanya, E.J., Cadmus, S.I., 2020. Seroprevalence of brucellosis and associated factors among livestock slaughtered in Oko-Oba abattoir, Lagos State, southwestern Nigeria. *Pan African Medical Journal* 36. [10.11604/pamj.2020.36.53.21094](https://doi.org/10.11604/pamj.2020.36.53.21094).
- Ularanu, H., Owolodun, O., Woma, T., Audu, B., Aaron, G., Cholom, S., Shamaki, D., 2012. Molecular diagnosis of recent suspected outbreaks of peste des petits ruminants (PPR) in Yola, Adamawa State, Nigeria. *African Journal of Biotechnology* 11, 1158–1162. [10.5897/AJB10.1419](https://doi.org/10.5897/AJB10.1419).

- Underwood, W.J., Blauwiekel, R., Delano, M.L., Gillesby, R., Mischler, S.A., Schoell, A., 2015. Biology and diseases of ruminants (sheep, goats, and cattle), in: *Laboratory Animal Medicine*. Elsevier, pp. 623–694. [10.1016/B978-0-12-409527-4.00015-8](https://doi.org/10.1016/B978-0-12-409527-4.00015-8).
- Uwanibe, E.L., Olugbo, A.O., Ugwumba, N.S.C., Ifeanyi, S.S., Adenike, F., Ignatius, N.C., 2014. Detection of plasmid in *Listeria monocytogenes* isolated from fresh raw milk in Nigeria. *Animal and Veterinary Sciences* 2, 169. [10.11648/j.avs.20140206.12](https://doi.org/10.11648/j.avs.20140206.12).
- Victor, I., Akuve, B., Buba, E., Helen, A., 2017. Risk factors associated with pestes des petits ruminants (PPR) in sheep and goats in Makurdi, Benue State. *Archives on Veterinary Science and Technology* 2. [10.29011/2637-9988/100020](https://doi.org/10.29011/2637-9988/100020).
- Vijayachari, P., Sugunan, A.P., Shriram, A.N., 2008. Leptospirosis: An emerging global public health problem. *Journal of Biosciences* 33, 557–569. [10.1007/s12038-008-0074-z](https://doi.org/10.1007/s12038-008-0074-z).
- Vijayachari P, S.S., Vijayachari, P., Sugunan, A., Roy, S., Natara-jaseenivasan, K., 2014. Seroprevalence and carrier status for leptospirosis in cattle and goats in Andaman Island, India. *Journal of Veterinary Science & Technology* 05. [10.4172/2157-7579.1000205](https://doi.org/10.4172/2157-7579.1000205).
- de Vries, S.G., Visser, B.J., Nagel, I.M., Goris, M.G.A., Hartskeerl, R.A., Grobusch, M.P., 2014. Leptospirosis in Sub-Saharan Africa: A systematic review. *International Journal of Infectious Diseases* 28, 47–64. [10.1016/j.ijid.2014.06.013](https://doi.org/10.1016/j.ijid.2014.06.013).
- Wachida, N., Mtsem, C., Makama, P.D., 2018. Peste des petits ruminants infection and infertility in West African Dwarf (WAD) goats in makurdi, benue state, Nigeria. *Vom Journal of Veterinary Science* 13, 26 – 31. [10.13140/RG.2.2.17642.21445](https://doi.org/10.13140/RG.2.2.17642.21445).
- Walker, B., 2005. Diseases causing reproductive losses in breeding cattle. Agfact , 1–5 URL: https://www.dpi.nsw.gov.au/_data/assets/pdf_file/0006/160368/cattle-reproductive-disease.pdf.
- Wernike, K., Silaghi, C., Nieder, M., Pfeffer, M., Beer, M., 2014. Dynamics of Schmallenberg virus infection within a cattle herd in Germany, 2011. *Epidemiology and Infection* 142, 1501–1504. [10.1017/S0950268813002525](https://doi.org/10.1017/S0950268813002525).
- Woma, T.Y., Kalla, D.J.U., Ekong, P.S., Ularanu, H.G., Chollom, S.C., Lamurde, I.I., Bajehson, D.B., Tom, N.D., Aaron, G.B., Shamaki, D., Bailey, D., Diallo, A., Quan, M., 2015. Serological evidence of camel exposure to peste des petits ruminants virus (PPRV) in Nigeria. *Tropical Animal Health and Production* 47, 603–606. [10.1007/s11250-014-0747-6](https://doi.org/10.1007/s11250-014-0747-6).
- Wungak, Y., Aworh, M., Maurice, N., Balami, A., Danmarwa, A., Danthe, H., 2011. Serological survey of antibodies against *Bru-cella abortus* in cattle in Jos-South Local Government Area. *Vom Journal of Veterinary Science* 8, 39–42.
- Yaeger, M.J.H., 2007. Bacterial causes of bovine infertility and abortion, in: *Current therapy in large animal theriogenology*. Elsevier, pp. 389–399.
- Yakubu, Y., Salihu, M., Faleke, O., Abubakar, M., Junaidu, A., Magaji, A., Gulumbe, M., Aliyu, R., 2012. Prevalence and antibiotic susceptibility of *Listeria monocytogenes* in raw milk from cattle herds within Sokoto Metropolis, Nigeria. *Sokoto Journal of Veterinary Sciences* 10. [10.4314/sokjvs.v10i2.3](https://doi.org/10.4314/sokjvs.v10i2.3).
- Yousef, M.R., Al-Eesa, A.A., Al-Blowi, M.H., 2012. High seroprevalence of bluetongue virus antibodies in sheep, goats, cattle and camel in different districts of Saudi Arabia. *Veterinary World* 5, 389. [10.5455/vetworld.2012.389-393](https://doi.org/10.5455/vetworld.2012.389-393).
- Youssefi, M., Arabkhazaeli, F., Hassan, A.T.M., 2009. Seroprevalence of *Neospora caninum* infection in rural and industrial cattle in northern Iran. *Iranian Journal of Parasitology* 4, 15–18. URL: <https://www.sid.ir/FileServer/JE/114920090103.pdf>.
- Yusuf, Y., Abdurashheed, A., 2019. Survey on bovine brucellosis in Sokoto Metropolitan Abattoir, Nigeria. *Journal of Animal and Veterinary Advances* 18, 143–146. [10.36478/javaa.2019.143.146](https://doi.org/10.36478/javaa.2019.143.146).
- Zahur, A.B., Irshad, H., Ullah, A., Afzal, M., Latif, A., Ullah, R.W., Farooq, U., Samo, M.H., Jahangir, M., 2014. Peste des petits ruminants vaccine (Nigerian strain 75/1) confers protection for at least 3 years in sheep and goats. *Journal of Biosciences and Medicines* 02, 27–33. [10.4236/jbm.2014.26005](https://doi.org/10.4236/jbm.2014.26005).
- Zanna, M.Y., El-Yuguda, A.D., Shettima, Y.M., Abubakar, M.B., Baba, S.S., 2021. Detection of bluetongue virus nucleic acid in ruminants and camels (*Camelus dromedarius*) in Maiduguri, Nigeria. *Journal of Agriculture and Veterinary Science* 4, 37–43. URL: <https://www.iosrjournals.org/iosr-javs/papers/Vol14-issue4/Series-1/F1404013743.pdf>.
- Zhao, G.H., 2012. Seroprevalence of chlamydial infection in dairy goats in Shaanxi Province, Northwestern China. *African Journal of Biotechnology* 11. [10.5897/AJB11.2308](https://doi.org/10.5897/AJB11.2308).
- Zhao, H., Njeumi, F., Parida, S., Benfield, C.T.O., 2021. Progress towards eradication of peste des petits ruminants through vaccination. *Viruses* 13. [10.3390/v13010059](https://doi.org/10.3390/v13010059).
- Zubairu, A., Ardo, M., Mai, H., 2014. Seroprevalence of ruminant brucellosis in three selected local government areas of Taraba state. *Sokoto Journal of Veterinary Sciences* 12, 51. [10.4314/sokjvs.v12i1.8](https://doi.org/10.4314/sokjvs.v12i1.8).