

Mitteilungen und Nachrichten

Aus den Arbeitskreisen der Deutschen Phytomedizinischen Gesellschaft (DPG):

Bericht zum Jahrestreffen 2012 der Arbeitskreise „Mykologie“ und „Wirt-Parasit-Beziehungen“ der Deutschen Phytomedizinischen Gesellschaft

Das Jahrestreffen 2012 der Arbeitskreise „Wirt-Parasit-Beziehungen“ und „Mykologie“ der Deutschen Phytomedizinischen Gesellschaft e.V. fand am 22. und 23. März 2012 an der Universität Stuttgart-Hohenheim statt. Gastgeber war Herr Professor Dr. Ralf VÖGELE, der das Treffen bereits kurz nach seiner Berufung nach Hohenheim mit seinen Mitarbeitern hervorragend organisierte. Wie in den vergangenen Jahren wurde das Jahrestreffen der beiden Arbeitskreise in einer gemeinsamen Arbeitssitzung am ersten Tag und in zwei getrennten Sitzungen am zweiten Tag durchgeführt. Insgesamt waren ca. 120 Wissenschaftlerinnen und Wissenschaftler zum gemeinsamen Treffen nach Hohenheim gekommen. Unter ihnen war der wissenschaftliche Nachwuchs mit aktiven Beiträgen besonders stark vertreten. Insgesamt wurden 39 Vorträge gehalten. Darüber hinaus wurden Poster (fünf im Arbeitskreis „Wirt-Parasit-Beziehungen“ und neun im Arbeitskreis „Mykologie“) präsentiert, die an beiden Tagen intensiv diskutiert wurden. Die in den Beiträgen vorgestellten Themen waren vielfältig. Sie umfassten Veränderungen der Genexpression, sowie Proteom- und Sekretomanalysen bei der Pathogenese von Pilzen, Oomyceten und Nematoden. Darüber hinaus behandelten sie die Identität und Wirkung von pilzlichen Effektoren und Pathogenitätsfaktoren, Wirt- und Nicht-Wirt-Resistenz von Pflanzen, sowie die Induzierte Resistenz. Außerdem wurden die Synthese, das Spektrum und die Verteilung von Mykotoxinen in infizierten Pflanzen erläutert und taxonomische Aspekte, das Auftreten und die Charakterisierung von neuen Schadpilzen, sowie das Vorkommen und die Ursachen von Fungizid-Resistenzen besprochen.

Das nächste gemeinsame Jahrestreffen der Arbeitskreise „Wirt-Parasit-Beziehungen“ und „Mykologie“ wird am 21. und 22. März 2013 an der Georg-August-Universität Göttingen stattfinden; Gastgeber wird Prof. Dr. A. VON TIEDEMANN sein.

2012 Report on the Annual Meeting of the Working Groups 'Host-Parasite Interactions' and 'Mycology'

This year's Annual Meeting of the Study Groups 'Host-Parasite Interactions' and 'Mycology' of the German Phytomedical Society (Deutsche Phytomedizinische Gesellschaft e.V.) was held on March 22–23, 2012 at Hohenheim University. The meeting was hosted, and perfectly organized, by Prof. Ralf T. VÖGELE and associates. As for many years, the meeting was set up as a joint workshop with one combined and two separate sessions. In total, 120 senior and junior scientists attended the joint meeting presenting 39 talks and 14 posters (five on 'Host-Parasite Interactions' and nine on 'Mycology'). Talks and posters covered diverse topics, such as gene expression in pathogens and attacked plants, fungal effector molecules, proteome and secretome analyses, pathogenicity factors, host and nonhost disease resistance, induced resistance, mycotoxins, fungicide sensitivity, taxonomy, and new pathogens, to name but a few.

The next joint meeting of the Working Groups 'Host-Parasite Interactions' and 'Mycology' will be held on March 21–22, 2013

at Georg-August-University Göttingen. It will be hosted by Professor Dr. A. VON TIEDEMANN.

Für den AK Wirt-Parasit-Beziehungen
Prof. Dr. Uwe CONRATH (Aachen)
Für den AK Mykologie PD Dr. Ulrike STEINER (Bonn)

Einige Abstracts/Zusammenfassungen der Arbeitskreise werden nachfolgend aufgeführt:

Abstracts AK Wirt-Parasit-Beziehungen 2012

1) Do effectors of *Sporisorium reilianum* induce phytoalexin biosynthesis in *Sorghum bicolor*?

Jan UTERMARK, Katja ZUTHER, Jan SCHIRAWSKI

Georg-August-Universität Göttingen, Albrecht-von-Haller Institut für Pflanzenwissenschaften Abt. Molekularbiologie der Pflanzen-Mikroben-Interaktion Schwann-Schleiden-Forschungszentrum, Julia-Lermontowa-Weg 3, 37077 Göttingen

The basidiomycete fungus *Sporisorium reilianum* occurs in two varieties (*S. reilianum* f. sp. *reilianum* [SRS] and c.f.sp. *zeae* [SRZ]) that cause head smut disease on sorghum and maize, respectively, replacing inflorescences with fungal spores. Infection of sorghum with SRZ leads to a strong host response: The phytoalexin biosynthesis gene SbDFR3 is highly induced, which leads to accumulation of the red phytoalexin luteolinidin in leaves colonized by SRZ but not by SRS. Luteolinidin has a growth-inhibitory effect on *S. reilianum*, suggesting that regulation of luteolinidin biosynthesis determines infection success of *S. reilianum* on sorghum.

To better understand the factors underlying the host specificity of the two *S. reilianum* varieties, we search for effector proteins of SRZ capable of inducing luteolinidin biosynthesis in sorghum. We bioinformatically identified in the genome of SRZ > 200 candidate genes encoding effector-like secreted proteins, several of which are only expressed during plant colonization. To test whether any of these is involved in luteolinidin induction, we establish three different experimental assays. We express fungal effector genes in sorghum protoplasts and test SbDFR3 expression. We use biolistic transformation to deliver *S. reilianum* effector genes to sorghum leaves. The third assay involves delivery of *S. reilianum* effectors to sorghum using the bacterial type III secretion system from *Xanthomonas* sp. and *Pseudomonas* sp. Identification of the effector proteins that trigger phytoalexin biosynthesis in sorghum will provide new insights into the evolution of virulence and host range of *S. reilianum*.

(DPG AK Wirt-Parasit-Beziehungen)

2) Secretome Analysis of the Asian Soybean Rust *Phakopsora pachyrhizi*

Matthias KOHLNDORFER, Tobias LINK, Ralf T. VÖGELE

Universität Hohenheim, Fakultät Agrarwissenschaften, Institut für Phytomedizin, FG Phytopathologie, Otto-Sander-Str. 5, 70599 Stuttgart

Rust fungi have drawn the interest of scientists for decades, not only for their economic importance, but also because of some peculiarities of their lifestyle. The most fascinating aspect in this lifestyle is their obligate biotrophic growth, which is based on establishing a long lasting interaction with a living host plant. The hallmark of obligate biotrophy is the formation of

specialized hyphae inside the plant cell, so-called haustoria. These structures represent the most intimate contact zone between pathogen and host and play a vital role in the uptake of nutrients and the exchange of information. In order to suppress plant defense mechanisms, rust fungi produce specialized proteins, so-called effectors. These proteins are secreted into the extrahaustorial matrix or even into the plant cell, where they are expected to interact with plant proteins. Knowledge about these mechanisms might provide valuable clues for plant protection in the future, especially with respect to RNAi based approaches. Our work is focused on the identification and analysis of novel effector proteins from different rust species. To select for putative secreted proteins, we are working with a method called „Signal Sequence Trap“ (JACOBS et al., 1999). This method has already been used for *Uromyces fabae* (LINK and VOEGELE, 2008) and *Uromyces appendiculatus*. We are using cDNA libraries of *Phakopsora pachyrhizi* generated from infected soybean leaves (*Glycine max*). To enrich the cDNA for fungal cDNA we employ Subtractive Hybridization. Comparing the different rust secretomes across species might lead to the identification of novel effector proteins, essential for virulence.

JACOBS, K.A., et al., 1999: A genetic selection for isolating cDNA clones that encode signal peptides. *Methods Enzymol.* **303**, 468-479.

LINK, T.I., R.T. VOEGELE, 2008: Secreted proteins of *Uromyces fabae*: similarities and stage specificity. *Mol. Plant Pathol.* **9**, 59-66.

(DPG AK Wirt-Parasit-Beziehungen)

3) Haustorial transcriptomes of *Uromyces appendiculatus* and *Phakopsora pachyrhizi* – identification of families of candidate effectors

Tobias LINK

Universität Hohenheim, Fakultät Agrarwissenschaften,
Institut für Phytomedizin, FG Phytopathologie, Otto-Sander-Str. 5,
70599 Stuttgart

Rust fungi are biotrophic pathogens, which mean that they do not kill their respective host plants but are dependent on living tissue for propagation. Among them are species with major economic impact like *Phakopsora pachyrhizi* and *Uromyces appendiculatus*, infecting soybean and common bean respectively. A hallmark feature of biotrophic fungi are haustoria, which were shown to be the interface for nutrient uptake in rust fungi and probably are the place where effector proteins are transferred to the plant. Effector proteins can suppress the host resistance response and may have other functions in influencing the plant host, making establishment and maintenance of the biotrophic interaction possible. To identify effector candidates we did transcriptome sequencing using the next generation sequencing technology 454 pyrosequencing. For this cDNA was prepared from isolated haustoria of both *U. appendiculatus* and *P. pachyrhizi*. Comparing our annotation results with those for prebiotrophic structures we could corroborate findings that haustoria have indeed important functions in energy and amino acid metabolism. Blasting our sequences against other rust and basidiomycete genome sequences, predicting secreted proteins and building gene families through a clustering analysis, we could identify genes and gene families that are secreted and that are specific to rust fungi or subclades to the rust fungi. In addition to this, interesting motifs and expression patterns make these genes and gene families good candidate effectors.

(DPG AK Wirt-Parasit-Beziehungen)

Abstracts Arbeitskreis Mykologie 2012

1) Ash dieback in southwest Germany and genetic investigations on the causal agent *Hymenoscyphus pseudoalbidus*

Rasmus ENDERLE¹, Jörg GRÜNER², Siegfried FINK², Berthold METZLER¹

¹ Forstliche Versuchs- und Forschungsanstalt Baden-Württemberg, Abt. Waldschutz, Wonnhaldestr. 4, 79100 Freiburg

² Universität Freiburg, Professur für Forstbotanik, Bertoldstr. 17, 79085 Freiburg

Ash dieback is observed in southwest Germany since 2009 and causes increasingly damage. At present, planting of ashes is not recommended. The question arises in which extent *Fraxinus excelsior* will be suitable for forestry purposes in future. Characteristic symptoms of this new disease include wilting of leaves, premature leaf fall, shoot dieback and bark necroses. In young trees, the disease leads to bushy or dwarf growth, whereas the wood quality of older trees is endangered by formation of secondary shoots at the stems. Data concerning the occurrence of the disease in Baden-Württemberg was collected every year since 2009. Ash dieback occurred on 2505 ha in autumn 2009, 4106 ha in autumn 2010 and 8526 ha in autumn 2011. On 3133 ha the disease was rated as threatening whole stands. Particularly affected are stands in the Rhine valley. The genetic variation of 64 isolates of the causal agent *Hymenoscyphus pseudoalbidus* from southwest Germany was analysed by RAMS-fingerprinting (random amplified microsatellites) using the primer VDV(CT)₇C. A considerable amount of genetic variation among these strains was detected. That implies that the fungus has a heterothallic life cycle. Furthermore, the findings suggest a fast increase in genetic variation, although more research is needed to confirm this result. Nevertheless, it is a cause of concern, keeping in mind that the pathogen may be able to overcome the reported individual resistance of *F. excelsior* in future. Also, this could explain the contradiction between the indications of *H. pseudoalbidus* as an invasive species on the one hand and its high genetic variation on the other hand. There was no evidence for a connection between the geographic origin and the genetic distance of the strains. This is an indication that the dispersal of the pathogen's ascospores by the wind takes place over large distances. But also trading of nursery material may have contributed to this phenomenon.

(DPG AK Mykologie)

2) Grey mould isolates from German strawberry fields show multiple fungicide resistance and represent a novel clade between *B. cinerea* and *B. fabae*

Matthias HAHN¹, Michaela LEROCH¹, Cecilia PLESKEN¹, Zinnia NAOSHIN¹, Roland WEBER²

¹ TU Kaiserslautern, FB Biologie, Postfach 3049, 67663 Kaiserslautern

² ESTEBURG-Obstbauzentrum Jork, Moorende 53, 21635 Jork

Grey mold is a major problem of fruit and vegetable production worldwide. For control of Botrytis, strawberries receive multiple fungicide treatments each year. Grey mold populations from different German strawberry growing regions were tested for their sensitivity against Botrytis fungicides and for their genetic variability. Fungicide resistance was observed, including many isolates with multiple resistance against the majority of fungicides tested. A novel, stronger variant of the previously described multidrug resistance phenotype MDR1 was observed, called MDR1 h, conferring threefold higher, partial resistance to cyprodinil and fludioxonil. The majority of strawberry iso-