

Genome-wide identification of putative disease resistance genes (R genes) in carrot (*Daucus carota* subsp. *sativus*) by homology-based gene prediction

Anastassia Boudichevskaia¹, Thomas Berner², Jens Keilwagen², Frank Dunemann^{3*}

¹KWS SAAT SE & Co. KGaA, Grimsehlstr. 31, 37574 Einbeck, Germany

²Julius Kühn-Institut (JKI), Federal Research Centre for Cultivated Plants, Institute for Biosafety in Plant Biotechnology, Erwin-Baur-Str. 27, 06484 Quedlinburg, Germany

³Julius Kühn-Institut (JKI), Federal Research Centre for Cultivated Plants, Institute for Breeding Research on Horticultural Crops, Erwin-Baur-Str. 27, 06484 Quedlinburg, Germany

*correspondence to: frank.dunemann@julius-kuehn.de

Abstract

The cultivated carrot (*Daucus carota* ssp. *sativus*) is one of the most important root vegetable crops grown worldwide. Carrots are highly susceptible to several pests and diseases, and disease resistance is currently among the main breeding aims. The inheritance of resistance has been reported for a few carrot foliar diseases and root-knot nematodes, but no functionally characterized resistance gene (R gene) has yet been linked as a candidate gene to any resistance locus in carrot. Knowledge about the inventory of NLR genes (nucleotide-binding leucine-rich repeat receptors) and other R genes encoding transmembrane proteins such as receptor-like proteins (RLPs) and kinases (KIN) would be necessary to associate major QTLs (quantitative trait loci) identified by bi-parental QTL analyses or GWAS (genome-wide association analysis) with functional candidate R genes. In this study, we describe a combination of a genome-wide inventory of putative full-length carrot R genes based on a homology-based gene prediction approach called GeMoMa and subsequent classification by usage of the recent version of PRGdb 4.0 database (Calle-Garcia et al. 2022). A total of 320 putative carrot R genes were identified and bioinformatically characterized, including 72 newly identified gene models, that have not yet been annotated in the currently available carrot whole genome sequence. Based on the DRAGO 3 pipeline, totally 137 putative NLR genes were found, whereas 162 putative functional RLP and KIN genes were identified in the carrot genome. About one third of the R genes was found to be organized in clusters consisting only of NLR, RLP or KIN genes. To determine the evolutionary relationships of carrot R gene predictions, we generated a phylogenetic tree based on the alignment of all 320 R proteins. Three large clades (NLR, RLP and KIN) and a small clade (RLK) were identified, which reflect well the classification obtained after DRAGO 3 analysis. The presented carrot R gene inventory might be useful for resistance gene isolation, the development of (functional) molecular markers and resistance breeding in carrot.

Keywords: Carrot genome, resistance gene, annotation, gene prediction, R gene inventory, R gene map, gene cluster

Background

The cultivated carrot (*Daucus carota* ssp. *sativus*) is one of the most important root vegetable crops grown worldwide. Carrot production can be affected by a wide range of pests and pathogens. At least five diseases of carrot are caused by bacterial pathogens, 36 by fungal and oomycete pathogens, two by phytoplasmas, and 13 by viruses (Du Toit et al. 2019). Additionally, seven genera of nematodes and several insect and mite pests can impair carrot cultivation (Du Toit et al. 2019). Among the most important foliar diseases are leaf blight caused by the fungus *Alternaria dauci*, leaf spot disease (*Cercospora carotae*), bacterial blight (*Xanthomonas hortorum* pv. *carotae*), and powdery mildew (*Erysiphe heraclei*) (Davis and Raid 2002). The most widespread soilborne root pathogens of carrot are cavity spot (caused by several species of *Pythium*), white mold (*Sclerotinia sclerotiorum*), and root-knot nematodes (various species of *Meloidogyne*) (Davis and Raid 2002). Storage diseases are mainly caused by necrotrophic fungi, such as *Botrytis cinerea* (gray mold), and *Mycocentrospora acerina* (liquorice rot disease).

Despite the numerous biotic stress factors that can impact carrot growth and storage, comparatively less is known about putative resistances and, therefore, about the genetics of resistances. The inheritance of resistance has been reported for a few carrot foliar diseases and root-knot nematodes. For instance, monogenic resistance to two foliar diseases, *Cercospora* leaf spot and powdery mildew, was reported, but resistance genes have not yet been mapped (Simon 2019). Several quantitative trait loci (QTLs) for leaf blight resistance (*Alternaria dauci*) were identified by Le Clerc et al. (2015, 2019). For resistance to the root knot nematode *Meloidogyne javanica* two loci called as *Mj-1* and *Mj-2* were mapped on carrot chromosome 8 at different positions (Ali et al. 2014, Parsons et al. 2015). With a single exception, no predicted or functionally characterized resistance gene (R gene) has been associated yet as a candidate gene with any resistance in carrot. A cluster consisting of four putative R genes spanning a region of only 50 kb co-localized in the same region of chromosome 8 as the *Mj-1* locus (Iorizzo et al. 2016). The occurrence of the first carrot whole genome sequence (Iorizzo et al. 2016) and the progress made in candidate gene identification by SNP-based genome-wide association analyses (GWAS) of carrot natural substances (Keilwagen et al. 2017, Ellison et al. 2018) implicate such approaches also for carrot disease resistance research. As a first basis, a preliminary characterization of the carrot R gene inventory in the carrot genome has yielded a large number of more than 600 putative R gene sequences (Iorizzo et al. 2016).

As plants are immobile, they have evolved two major types of disease resistance, basal defense and R gene mediated defense (Gururani et al. 2012). Basal defense, which can be a constituent of both non-host and host resistance, provides the first line of defense to the infection by a wide range of pathogens. R gene-mediated pathogen resistance is mainly based on effector molecules encoded by Avr (avirulence) genes of the pathogen, which are delivered directly into the plant cells during initial stage of infection. The effectors found in viral pathogens, bacteria, oomycetes, fungi, nematodes or insects cause a plant pathogen to elicit a resistance response in a host plant. These receptors coded by R genes are mainly intracellular, and they can specifically interact with pathogen effectors coded by the Avr genes following the gene-for-gene relationship model (Borelli et al. 2018). Plant R genes have been identified from numerous species through genetic approaches, and these loci are associated with resistance to a wide spectrum of pathogens. Although plants are confronted by such a large variety of pathogens with different modes of pathogenesis (e.g., biotrophs versus necrotrophs), most R genes belong to a limited set of proteins that are made up of a conserved set of domains, whose organization is used to define them (Dangl and Jones 2001, Hammond-Kosack and Parker 2003). Through a comprehensive review, Kourelis and van der Horn (2018) identified a number of 314 cloned functional

plant R genes. Only 128 of the 314 gene products have a proposed mechanism, and the majority of these R genes encode cell surface or intracellular receptors (Kourelis and van der Horn 2018).

More than two thirds of plant disease resistance genes encode nucleotide-binding leucine-rich repeat receptors (NLRs), and most plant genomes carry a repertoire of hundreds of NLR genes (Van de Weyer et al. 2019). Based on their N-terminal structures, these R proteins can be further divided into two subclasses: TIR-NBS-LRR (TNL) that possesses a domain homologous to the Toll and interleukin-1 receptor (TIR), and non-TNL. Most non-TNL R proteins have a coiled-coil (CC) structure at the N terminal and are often called CC-NBS-LRR (CNL) R proteins (Wei et al 2018). The LRRs (Leucine rich repeats) represent the components that play an important role in recognition specificity, and these domains are present in the majority of R proteins (Jones 2001). There are a few further classes including groups that contain neither LRRs nor NBS domains but other functional domains, such as an intracellular serine-threonine kinase domain (Gururani et al. 2012). Transmembrane receptor proteins containing kinase and LRR domains, such as receptor-like proteins (RLP) and the receptor-like kinases (RLK), are also involved (Osuna-Cruz 2018). In the last years, several online omics platforms have been offered to facilitate the exploration, inventory and use of plant resistance genes. Among these platforms, the Pathogen Recognition Genes database (PRGdb) (Sanseverino et al. 2010) represents a user-friendly reference site and repository for plant geneticists interested in structure and putative function of genes involved in plant disease resistance.

In this study, we describe a combination of a genome wide inventory of predicted full-length carrot R genes based on a homology-based gene prediction approach called GeMoMa (Keilwagen et al. 2016) and usage of the PRGdb 4.0 database (Calle-Garcia et al. 2022) for classifying the predicted genes. A total of 320 putative carrot R genes were identified and bioinformatically characterized, including 72 gene models that have not yet been annotated in the current version of the carrot genome presented by Iorizzo et al. (2016).

Methods

For detecting resistance genes in carrot, homology-based gene prediction was utilized using a wide range of resistance genes from diverse plant species. Based on publicly available sequences of 110 known plant resistance genes, we collected the information about resistance transcript isoforms from eight reference genome annotations including *Arabidopsis thaliana*, *Glycine max*, *Malus domestica*, *Populus trichocarpa*, *Solanum lycopersicum*, *Solanum tuberosum*, *Sorghum bicolor*, and *Vitis vinifera*. In total, 810 putative R gene sequences selected from the reference genomes were used for homology-based gene prediction using the software GeMoMa. GeMoMa (version 1.4.3) was run for each reference species in combination with RNA-seq evidence (Keilwagen et al. 2017, 2018). The predictions of all reference species were combined using the module GAF yielding 602 gene predictions for carrot. The Integrative Genomics Viewer (IGV) (Robinson et al. 2011) was used as a visualization tool for the interactive exploration of predicted transcripts and annotated *Daucus* genome loci (DCAR sequences, Iorizzo et al. 2016). After this first round of screening, 375 gene models were selected for a second round of analysis to examine if the gene models are indeed complete. Translated protein sequences were analyzed manually by a BLAST P search at NCBI using the non-redundant protein sequence database, and only predicted full-length sequences with putative R gene function were selected for the final R gene inventory. DRAGO 3, the tool for automatic annotation and prediction of plant resistance genes implemented in PRGdb 4.0 database was used to identify the LRR, Kinase, NBS and TIR domains from 60 HMM modules created for this purpose using HMMER v3 package (Calle-Garcia et al. 2022). DRAGO 3 is also able to detect CC and TM domains using COILS 2.2 and TMHMM 2.0c

programs (Calle-Garcia et al. 2022). A phylogenetic analysis of 320 deduced *Daucus carota* R proteins was performed with MEGA-X after multiple sequence alignment by MUSCLE. Neighbor-joining method and n=1000 replicates for bootstrapping were applied. The software *MapChart* 2.2. (Plant Research International, Biometris, Wageningen, Netherlands) was used for R gene map visualization. R genes were considered clustered if at least three genes from the classes NLR, RLP or KIN were located within a genomic region < 200 kb.

Results and Discussion

The objective of this research was to conduct a genome-wide (re)-identification of R genes in the cultivated carrot. Initially, we predicted 602 putative R gene models after GeMoMa analysis. After manual evaluation using the IGV browser and elimination of too short and obviously fragmentary sequences, followed by BLAST P-based analyses for full-length CDS and predicted functions, 320 putative R gene models remained (Suppl. Table 1). Among these predicted genes, 72 of them (23%) have been newly identified since they have not yet been annotated in the current version of the carrot genome (Iorizzo et al. 2016). In addition, some new R gene clusters, such as the six-gene cluster at the beginning of chromosome Chr_3 (*DcRG_082* - *DcRG_87*) or the four-gene cluster on Chr_7 (*DcRG_233* - *DcRG_236*) were detected. Based on the DRAGO 3 pipeline, 137 putative NLR genes (43%) were found, whereas 172 putative genes encoding transmembrane proteins (54%) were identified in the carrot genome (Table 1). Among the NLR genes, the largest sub-class is the NBS-LRR (NL) type, followed by the TIR-NBS-LRR (TNL) type (Table 1). A few genes showing rare or unknown domain combinations such as CK or CL (Osuna-Cruz et al. 2018) were indeed genes possibly involved in plant resistance. For instance, genes *DcRG_169* and *DcRG_192* (CTNL class) were predicted in the *D. carota* genome as TMV resistance protein N-like isoforms X1 and X2 (Table 2). Only two genes (*DcRG_079* and *DcRG_222*) were not classified by DRAGO domain recognition but were predicted by BLAST as putative EDR2s (Table 2). These carrot genes are similar (amino acid identity ~ 80%) to the *A. thaliana* *EDR2* resistance gene (AT4G19040, Vorwerk et al. 2007). Because loss-of-function mutations in the *EDR2* gene confer enhanced disease resistance to powdery mildew in *A. thaliana*, *EDR2* probably functions as a negative regulator of powdery mildew resistance (Vorwerk et al. 2007). In the work of Iorizzo et al. (2016), MATRIX-R pipeline was used to automatically retrieve, annotate, and classify carrot R genes. Based on this pipeline, 634 putative R genes were predicted, which is comparable to the 602 gene models detected after GeMoMa analysis. However, based on sequence analyses of a random sample, we noticed that not all predicted sequences appeared to represent putative functional R genes (data not shown). In the study of Iorizzo et al. (2016), 295 R genes were assigned to the NLR classes of cytoplasmic proteins, and 339 genes were classified as transmembrane receptors. The NLR sub-class with the highest number of genes was the NL class (63 genes), which is in accordance with our finding of 64 NL genes in this sub-class (Table 1). The proportion of the NLRs at the total number of R genes was 46%, which is similar to the value found after GeMoMa (43%).

Over the past two decades, sequencing technologies have been rapidly developed and used to assess plant-microbe interactions (Lee et al. 2015). Their use enables genome-wide analyses of NLR genes based on the NB-ARC domain (Meyers et al. 2005). Because of their importance in ecology and breeding, there has been much interest in defining inventories of NLR genes at different taxonomic levels. These efforts have revealed that the number of NLR genes across species varies from less than a hundred to over a thousand (van de Weyer et al. 2019). The number of NLR genes in flowering plants is largely variable without any clear correlation to the phylogeny, suggesting species-specific mechanisms in NLR genes expansion and/or contraction (Jacob et al. 2013). *Arabidopsis thaliana* has

159 NLR genes including 43 CNLs and 83 TNLs (Guo et al. 2011). Solanaceae plants carry more than twice the number of NLR genes than *Arabidopsis* and possess more CNLs than TNLs (Lee et al. 2015). As reviewed by Jacob et al. (2013), species with large numbers of NLRs are rice and grape with each about 460 NLR genes. Nearly 1000 NLR genes have been identified in apple, although its genome size is only approximately 740 Mb (Velasco et al. 2010). Furthermore, there is also variation in the gene copy numbers of the two subclasses, TIR-type NLRs (TNL) and non-TIR-type NLRs (CNL). In our study, we observed a CNL-type NLR to TNL-type NLR ratio of 1:2. The same ratio was found in Brassicaceae species, in potato and grapevine it was 4:1, whereas a ratio of 1:1 has been reported for apple (Borelli et al. 2018). The importance of the NLR-type R genes for plant disease resistance is also documented by the number of cloned R genes. Out of the more than 300 cloned genes, 61% encode NLRs, but only 19% of the cloned R genes encode RLPs and RLKs (Kourelis and van der Horn 2018).

The predicted carrot R genes are not evenly distributed over the nine chromosomes. The number of R genes varied between only 19 on chromosome Chr_6 and 52 on Chr_3 (Table 3). As demonstrated by the R gene map presented in Figure 1, some genomic regions appear to have no or a fewer number of R genes. Especially dense R gene regions were observed on chromosomes Chr_1 (lower end), Chr_2 (whole chromosome), Chr_3 (upper and lower part), Chr_4 (middle part), Chr_5 (lower end), Chr_7 (lower half), Chr_8 (lower half), and Chr_9 (whole chromosome). NLRs, RLPs and kinases (KIN) showed a clear tendency for clustering (Figure 1, Suppl. Table 1). The size of clusters is rather variable. The largest cluster on Chr_8 contains 11 NLRs within an interval of about 250 kb, and the second-largest NLR cluster was found on Chr_7 with 9 genes within a 340 kb-interval. Both clusters are heterogenous and contain a mixture of diverse NLRs, i.e. NL, CNL, TN, and TNL. Smaller homogenous NLR clusters were also found, as for instance the TNL-gene cluster on Chr_2 or the NL-cluster on Chr_3 (Suppl. Table 1). The in total 8 RLP and 7 KIN clusters (Table 3) were, with two exceptions on Chr_1 and Chr_9, homogeneously in terms that they do not contain genes from the cytoplasmic NLR sub-classes. The three largest RLP clusters with each seven genes are located on chromosomes Chr_1, Chr_3 and Chr_5. NLR genes are known to be unevenly located in plant genomes and are often found in multigene clusters (Meyers et al. 2003). This clustered distribution likely arose by tandem duplications and further sequence divergence and depicts a large genetic reservoir for evolution of new specificities to pathogens (Duplessis et al. 2009). The homogeneous cluster type is most probably generated by tandem duplication, whereas the heterogenous cluster type is derived from ectopic duplications, transpositions, and/or large-scale segmental duplications with subsequent local rearrangements (Jacob et al. 2013). In the carrot genome, also other major R gene classes such as RLP and KIN were found to be organized in gene clusters.

To determine the evolutionary relationships of carrot R gene predictions, we generated a phylogenetic tree based on the alignment of all 320 deduced R proteins. Three large clades (NLR, RLP and KIN) and a small clade (RLK) were identified (Fig. 2). The predicted classification of the R genes is well represented by the phylogenetic relationships. Only a single predicted NL gene (*DcRG_010*) was placed among the RLP genes. The three genes with the unknown domain combination CK (*DcRG_101*, *DcRG_218* and *DcRG_253*) were assigned to the KIN clade, and the single CL gene was assigned to the RLP clade, respectively. The two EDR2 genes are closely related to the two CNL genes *DcRG_093* and *DcRG_158* and function in this phylogenetic analysis like an outgroup (Fig. 2). The similarity of the sequences of clustered R genes is often reflected by their positions in the phylogenetic tree (Fig. 2). Examples are the NLR cluster on Chr_2, which contains the six NL genes *DcRG_041* - *DcRG_046*, and the eleven-gene cluster on Chr_8 with the six genes *DcRG_269* - *DcRG_274*, which also formed a common NL sub-group. The other five genes from this cluster, however, were grouped in a common

TNL gene sub-group at a different position in the phylogenetic tree (Fig. 2). There are also some RLP and KIN gene clusters that obviously contain structurally similar R genes. For instance, RLP genes *DcRG_113 - DcRG_119* (Chr_3) and KIN genes *DcRG_258 - DcRG_262* (Chr_8) were located each in the same sub-group after phylogenetic analysis indicating common evolutionary origins.

Conclusion

The presented carrot R gene inventory consisting of more than 300 putative functional genes might be useful for resistance research and resistance breeding in carrot. Studies on the genotypic and phenotypic variability of resistances against several carrot diseases, supplemented by bi-parental QTL analyses or GWAS based on larger carrot genotype collections, may identify major QTLs and associated candidate R genes. In case of conventional breeding for resistance, introgression of resistance genes from resistant wild carrot relatives by repeated backcrossing is a long-lasting process. The use of molecular markers developed directly from R genes or R gene clusters might strongly support the backcrossing program. In addition, a deeper knowledge about the structure and function of isolated R gene candidates might be used in future for biotechnological approaches, such as CRISPR/Cas9-based genome editing, with the aim to generate new or to integrate known resistance alleles in the genome of the cultivated carrot.

Author's contributions

AB, TB, and JK performed the bioinformatic analyses. FD performed the evaluation and classification of predictions and compiled the carrot R gene inventory. FD drafted the manuscript. AB, JK, and FD contributed to the discussion and interpretation of results and read and approved the final manuscript.

References

- Ali A, Matthews WC, Cavagnaro PF, Iorizzo M, Roberts PA, Simon PW (2014) Inheritance and mapping of *Mj-2*, a new source of root-knot nematode (*Meloidogyne javanica*) resistance in carrot. *J Heredity* 105:288–291
- Andolfo G, Ercolano MR (2015) Plant innate immunity multicomponent model. *Front Plant Sci.* 6:987–994
- Borelli GM, Mazzucotelli E, Marone D, Crosatti C, Michelotti V, Valè G, Mastrangelo AM (2018) Regulation and Evolution of NLR Genes: A close interconnection for plant immunity. *Int J Mol Sci* 19:1662, DOI: 10.3390/ijms19061662
- Calle-Garcia J, Guadagno A, Paytuy-Gallart A, Saera-Vila A, Amoroso CG, D'Esposito D, Andolfo G, Cigliano RA, Sanseverino W, Ercolano MR (2022) PRGdb 4.0: an updated database dedicated to genes involved in plant disease resistance process. *Nucleic Acids Research* 50:D1483–D1490, DOI: 10.1093/nar/gkab1087
- Dangl JL, Jones JD (2001) Plant pathogens and integrated defense responses to infection. *Nature* 411:826–833
- Davis RM, Raid RN (eds) (2002) Compendium of umbelliferous crop diseases. American Phytopathological Society, APS Press, St. Paul, USA, 75pp.
- Duplessis S, Major I, Martin F, Séguin A (2009) Poplar and pathogen interactions: Insights from *Populus* genome-wide analyses of resistance and defense gene families and gene expression Profiling. *Critical Reviews in Plant Science* 28:309–334, DOI: 10.1080/07352680903241063
- Du Toit LJ, Le Clerc V, Briard M (2019) Genetics and Genomics of Carrot Biotic Stress. In: Simon P, Iorizzo M, Grzebelus D, Baranski R, editors. *The Carrot Genome*. Cham: Springer Nature Switzerland. 2019. p. 317–367, DOI: 10.1007/978-3-030-03389-7_18
- Ellison SL, Luby CH, Corak KE, Coe KM, Senalik D, Iorizzo M, et al. (2018) Carotenoid presence is associated with the *Or* gene in domesticated carrot. *Genetics* 210:1497–1508

- Guo YL, Fitz J, Schneeberger K, et al. (2011) Genome-wide comparison of nucleotide-binding site-leucine-rich repeat-encoding genes in *Arabidopsis*. *Plant Physiology* 157:757–769
- Gururani MA, Venkatesh J, Upadhyaya CP, Nookaraju A, Pandey SK, Park SW (2012) Plant disease resistance genes: Current status and future directions. *Physiological and Molecular Plant Pathology* 78:51e65, DOI: 10.1016/j.pmpp.2012.01.002
- Hammond-Kosack KE, Parker JE (2003) Deciphering plant-pathogen communication: fresh perspectives for molecular resistance breeding. *Current Opinions in Biotechnology* 14:177–193
- Iorizzo M, Ellison S, Senalik D, Zeng P, Satapoomin P, et al. (2016) A high-quality carrot genome assembly provides new insights into carotenoid accumulation and asterid genome evolution. *Nature Genetics* 48:657–666
- Jacob F, Vernaldi S, Maekawa T (2013) Evolution and conservation of plant NLR functions. *Front Immun* 4:Article 297, DOI: 10.3389/fimmu.2013.00297
- Jones JD (2001) Putting knowledge of plant disease resistance genes to work. *Current Opinions in Plant Biology* 4:281e7
- Keilwagen J, Wenk M, Erickson JL, Schattat MH, Grau J, Hartung F (2016) Using intron position conservation for homology-based gene prediction. *Nucleic Acids Research* 44:e89, DOI: 10.1093/nar/gkw092
- Keilwagen J, Lehnert H, Berner T, Budahn H, Nothnagel T, Ulrich D, Dunemann F (2017) The terpene synthase gene family of carrot (*Daucus carota* L.): Identification of QTLs and candidate genes associated with terpenoid volatile compounds. *Front Plant Sci* 8:1930, DOI: 10.3389/fpls.2017.01930
- Keilwagen J, Hartung F, Paulini M, Twardziojk SO, Grau J (2018) Combining RNA-seq data and homology-based gene prediction for plants, animals and fungi. *BMC Bioinformatics* 19:189
- Kourelis J, van der Hoorn RAL (2018) Defended to the nines: 25 years of resistance gene cloning identifies nine mechanisms for R protein function. *The Plant Cell* 30:285–299, DOI: 10.1105/tpc.17.00579
- Le Clerc V, Marques S, Suel A, Huet S, Hamama L, Voisine L, Auperpin E, Jourdan M, Barrot L, Prieur R (2015) QTL mapping of carrot resistance to leaf blight with connected populations: stability across years and consequences for breeding. *Theor Appl Genet* 128:2177–2187
- Le Clerc V, Aubert C, Cottet V, Yovanopoulos C, Piquet M, Suel A, et al. (2019) Breeding for carrot resistance to *Alternaria dauci* without compromising taste. *Molecular Breeding* 39:59, DOI: 10.1007/s11032-019-0966-7
- Lee H-A, Yeom S-I (2015) Plant NB-LRR proteins: tightly regulated sensors in a complex manner. *Briefings in Functional Genomics* 14:233–242, DOI: 10.1093/bfpg/elv012
- Meyers BC, Kozik A, Griego A, Kuang H, Michelmore RW (2003) Genome-wide analysis of NBS-LRR-encoding genes in *Arabidopsis*. *Plant Cell* 15:809–834
- Meyers BC, Kaushik S, Nandety RS (2005) Evolving disease resistance genes. *Current Opinion in Plant Biology* 8:129–134
- Osuna-Cruz CM, Paytavi-Gallart A, Di Donato A, Sundesha V, Andolfo G, Cigliano RA, Sanseverino W, Ercolano MR (2018) PRGdb 3.0: a comprehensive platform for prediction and analysis of plant disease resistance genes. *Nucleic Acids Research* 46, Database issue D1197–D1201, DOI: 10.1093/nar/gkx1119
- Parsons J, Matthews WC, Iorizzo M, Roberts PA, Simon PW (2015) *Meloidogyne incognita* nematode resistance QTL in carrot. *Molecular Breeding* 35:114
- Robinson JT, Thorvaldsdóttir H, Winckler W, Guttman M, Lander ES, Gatz G, et al. (2011) Integrative Genomics Viewer. *Nature Biotechnology* 29:24–26, DOI: 10.1038/nbt.1754
- Sanseverino W, Roma G, De Simone M, Faino L, Melito S, Stupka E, Frusciante L, Ercolano MR (2010) PRGdb: a bioinformatics platform for plant resistance gene analysis. *Nucleic Acids Research* 38: D814–D821
- Simon PW (2019) Classical and molecular carrot breeding. In: Simon P, Iorizzo M, Grzebelus D, Baranski R, editors. *The Carrot Genome*. Cham: Springer Nature Switzerland. 2019. p. 137–147, DOI: 10.1007/978-3-030-03389-7_9
- Van de Weyer A-L, Monteiro F, Furzer OJ, Nishimura MT, Cevik V, Witek K, Jones JD, et al. (2019) A species-wide inventory of NLR genes and alleles in *Arabidopsis thaliana*. *Cell* 178:1260–1272, DOI: 10.1016/j.cell.2019.07.038

Velasco R, Zharkikh A, Affourtit J, Dhingra A, Cestaro A, Kalyanaraman A, Fontana P, Bhatnagar SK, Troggio M, et al. (2010) The genome of the domesticated apple (*Malus x domestica* Borkh.). *Nature Genetics* 42:833–839

Vorwerk S, et al. (2007) EDR2 negatively regulates salicylic acid-based defenses and cell death during powdery mildew infections of *Arabidopsis thaliana*. *BMC Plant Biology* 7:35

Wei WC, Chen J, Kuang H, et al. (2018) Dramatic number variation of R genes in Solanaceae species accounted for by a few R gene subfamilies. *PLoS ONE* 11:e0148708, DOI:10.1371/journal.pone.0148708

Table 1. Predicted putative R genes identified in the carrot genome

R protein type	Class	No. of genes
<i>NLR</i>		
CC-NBS-LRR	CNL	19
TIR-NBS-LRR	TNL	40
NBS-LRR	NL	64
CC-NBS	CN	1
TIR-NBS	TN	10
NBS	N	3
<i>Transmembrane protein</i>		
Receptor-like kinase	RLK	10
Receptor-like protein	RLP	82
Kinase	KIN	80
<i>Unknown domain combination</i> ²⁾	CK, CL, CTNL	9
<i>Other</i> ²⁾		
Enhanced disease resistance 2	EDR2	2
		320

¹⁾ according to PRGdb 4.0 database (Calle-Garcia et al. 2022)

²⁾ for details, see Table 2

Table 2. R genes with unknown domain combinations (CK, CL, CTNL) or no domain recognition (EDR2) by DRAGO 3 predicted with highest max. score by NCBI BLAST P (best hit)






Gene name	Chromosome	Class	Locus name	Prediction NCBI BLAST P (<i>D. carota ssp. sativus</i>)	Max. Score	Sequence ID
<i>DcRG_002</i>	Chr_1	CTNL	NA ¹⁾	TMV resistance protein N-like isoform X3	2270	XP_017226921.1
<i>DcRG_006</i>	Chr_1	CTNL	DCAR_001436	Disease resistance protein TAO1-like	1477	XP_017224931.1
<i>DcRG_011</i>	Chr_1	CTNL	DCAR_002255	TMV resistance protein N-like	2292	XP_017230126.1
<i>DcRG_033</i>	Chr_1	CL	DCAR_004675	Receptor-like protein 12	1410	XP_017255570.1
<i>DcRG_079</i>	Chr_2	EDR2	DCAR_008189	Protein ENHANCED DISEASE RESISTANCE 2 isoform X1	1448	XP_017237032.1
<i>DcRG_101</i>	Chr_3	CK	NA ¹⁾	Wall-associated receptor kinase-like 20	1200	XP_017237511.1
<i>DcRG_169</i>	Chr_5	CTNL	DCAR_017232	TMV resistance protein N-like isoform X1	1283	XP_017222904.1
<i>DcRG_192</i>	Chr_6	CTNL	DCAR_022321	TMV resistance protein N-like isoform X2	2292	XP_017254942.1
<i>DcRG_218</i>	Chr_7	CK	DCAR_024644	Serine/threonine-protein kinase PBS1-like	872	XP_017218963.1
<i>DcRG_222</i>	Chr_7	EDR2	DCAR_024947	Protein ENHANCED DISEASE RESISTANCE 2-like	1449	XP_017219098.1
<i>DcRG_253</i>	Chr_8	CK	DCAR_028743	Probable serine/threonine-protein kinase NAK	770	XP_017222609.1

¹⁾ NA - not annotated in carrot genome vers.2

Table 3. Number of predicted R genes on carrot chromosomes and number of R gene clusters according to the inventory list (Suppl. Table 1)

Chromosome	No. of R genes	No. of R gene clusters ¹⁾		
		NLR	RLP	KIN
Chr_1	35	1	2	0
Chr_2	46	2	0	1
Chr_3	52	2	2	1
Chr_4	30	0	1	1
Chr_5	26	0	1	0
Chr_6	19	0	0	0
Chr_7	27	1	1	2
Chr_8	33	1	0	1
Chr_9	40	2	1	1
unknown	2			
	320	9	8	7

¹⁾ three or more genes, 200 kb interval definition (see Methods)






Figure 1. Carrot R gene map. Schematic map presentation of the genomic localization of the 320 carrot R genes listed in Suppl. Table 1. Figures on the left side of the bars show the start position of the CDS of each R gene in Mb (mega base pairs). R gene clusters are presented in coloured boxes (NLR - blue, RLP - yellow, KIN - green).




Figure 2. Phylogenetic analysis of 320 deduced *Daucus carota* R proteins and their grouping in clades (NLR, RLP, RLK and KIN). Multiple sequence alignment by MUSCLE (MEGA-X). Neighbor-joining method and n=1000 replicates for bootstrapping were applied (MEGA-X). Bootstrap values are shown next to each node.

Suppl. Table 1. List of 320 *Daucus carota* R gene models sorted by their physical position on the assembled nine carrot chromosomes according to the carrot whole genome sequence assembly vers.2 (Iorizzo et al. 2016). NLR gene clusters are shaded in blue, RLP clusters in yellow, and KIN clusters in green, respectively (for cluster definition, see methods). NA – Not annotated in carrot genome sequence vers.2

Gene name ¹⁾	Chrom.	Class ²⁾	Genomic coordinates ³⁾			Locus name ⁴⁾	Prediction (GeMoMa) ⁵⁾	Exon no.	AA ⁶⁾
			Strand	Start	Stop				
<i>DcRG_001</i>	Chr_1	TNL	-	3334635	3351268	DCAR_001225	GLYMA.03G087500.2.WM82.A2.V1_R3	6	817
<i>DcRG_002</i>	Chr_1	CTNL	-	3374751	3378828	NA	SOLYC05G006630.2.1.ITAG2.4_R9	5	1153
<i>DcRG_003</i>	Chr_1	TNL	-	3386037	3390016	NA	AT1G63870.1.TAIR10_R9	4	1025
<i>DcRG_004</i>	Chr_1	NL	-	4017929	4029504	DCAR_001279	POTRI.005G042900.1.V3.0_R0	3	1247
<i>DcRG_005</i>	Chr_1	KIN	+	14125232	14131946	DCAR_001366	GLYMA.09G181400.1.WM82.A2.V1_R6	3	659
<i>DcRG_006</i>	Chr_1	CTNL	+	15285224	15290985	DCAR_001436	POTRI.019G001700.1.V3.0_R1	6	1156
<i>DcRG_007</i>	Chr_1	TNL	-	15948244	15953280	NA	AT5G40100.1.TAIR10_R4	5	965
<i>DcRG_008</i>	Chr_1	TNL	+	19308075	19312726	DCAR_001691	SOLYC09G007710.2.1.ITAG2.4_R7	5	1226
<i>DcRG_009</i>	Chr_1	RLP	-	21268604	21271345	DCAR_001832	POTRI.009G112000.1.V3.0_R9	2	884
<i>DcRG_010</i>	Chr_1	NL	+	24550403	24551521	DCAR_002050	GLYMA.06G319700.1.WM82.A2.V1_R5	1	373
<i>DcRG_011</i>	Chr_1	CTNL	-	27479966	27485197	DCAR_002255	POTRI.005G206400.1.V3.0_R5	6	1180
<i>DcRG_012</i>	Chr_1	NL	+	30042291	30045532	NA	AT1G58410.1.TAIR10_R0	2	949
<i>DcRG_013</i>	Chr_1	RLP	-	31167439	31169202	DCAR_002613	AT5G23400.1.TAIR10_R0	1	588
<i>DcRG_014</i>	Chr_1	KIN	+	33733961	33736555	DCAR_002886	AT5G39030.1.TAIR10_R8	1	865
<i>DcRG_015</i>	Chr_1	RLK	+	38784490	38787184	DCAR_003427	POTRI.013G158800.1.V3.0_R6	2	639
<i>DcRG_016</i>	Chr_1	RLK	+	39874385	39877096	DCAR_003536	POTRI.013G158800.1.V3.0_R9	2	647
<i>DcRG_017</i>	Chr_1	NL	-	40335091	40337429	DCAR_003581	SOLYC07G039390.1.1.ITAG2.4_R8	4	405
<i>DcRG_018</i>	Chr_1	NL	-	40354145	40356865	DCAR_003582	SOLYC04G009660.2.1.ITAG2.4_R7	1	907
<i>DcRG_019</i>	Chr_1	RLP	-	41620093	41623047	DCAR_003743	AT2G34930.1.TAIR10_R1	1	985
<i>DcRG_020</i>	Chr_1	RLP	-	41625658	41628633	DCAR_003744	AT2G34930.1.TAIR10_R0	1	992
<i>DcRG_021</i>	Chr_1	RLP	-	41630254	41633226	DCAR_003745	AT2G34930.1.TAIR10_R2	1	991
<i>DcRG_022</i>	Chr_1	KIN	-	44989248	44999805	DCAR_004129	GLYMA.07G096100.1.WM82.A2.V1_R4	3	725
<i>DcRG_023</i>	Chr_1	NL	-	47059885	47063792	DCAR_004352	POTRI.002G056100.1.V3.0_R1	4	796

Gene name ¹⁾	Chrom.	Class ²⁾	Genomic coordinates ³⁾			Locus name ⁴⁾	Prediction (GeMoMa) ⁵⁾	Exon no.	AA ⁶⁾
			Strand	Start	Stop				
<i>DcRG_024</i>	Chr_1	TNL	-	47313350	47317926	NA	SOLYC09G092410.2.1.ITAG2.4_R3	5	1099
<i>DcRG_025</i>	Chr_1	TN	+	47833334	47835138	NA	GLYMA.02G023800.1.WM82.A2.V1_R3	2	339
<i>DcRG_026</i>	Chr_1	KIN	-	48680869	48683436	DCAR_004501	GLYMA.05G237100.1.WM82.A2.V1_R1	3	654
<i>DcRG_027</i>	Chr_1	KIN	+	49412713	49413804	DCAR_004576	GLYMA.07G095100.1.WM82.A2.V1_R9	1	364
<i>DcRG_028</i>	Chr_1	RLP	+	50256662	50259553	NA	AT1G47890.1.TAIR10_R8	1	964
<i>DcRG_029</i>	Chr_1	RLP	+	50267963	50270839	NA	POTRI.001G389100.1.V3.0_R2	1	959
<i>DcRG_030</i>	Chr_1	RLP	+	50281594	50284452	DCAR_004673	POTRI.011G104900.1.V3.0_R5	1	953
<i>DcRG_031</i>	Chr_1	RLP	+	50300950	50303859	NA	AT1G47890.1.TAIR10_R7	1	970
<i>DcRG_032</i>	Chr_1	RLP	+	50321690	50331579	NA	POTRI.001G389100.1.V3.0_R1	2	1053
<i>DcRG_033</i>	Chr_1	CL	+	50337331	50340131	DCAR_004675	POTRI.001G389100.1.V3.0_R9	3	899
<i>DcRG_034</i>	Chr_1	RLP	+	50352172	50355033	NA	POTRI.001G389100.1.V3.0_R0	1	954
<i>DcRG_035</i>	Chr_1	RLP	-	50383733	50386504	DCAR_004682	POTRI.001G389100.1.V3.0_R5	2	899
<i>DcRG_036</i>	Chr_2	NL	+	58662	61556	DCAR_004830	AT5G43470.1.TAIR10_R5	2	884
<i>DcRG_037</i>	Chr_2	NL	-	211201	213883	DCAR_004846	AT1G10920.1.TAIR10_R2	3	841
<i>DcRG_038</i>	Chr_2	NL	-	280061	283927	DCAR_004853	SOLYC12G096920.1.1.ITAG2.4_R4	3	955
<i>DcRG_039</i>	Chr_2	NL	-	938672	942000	DCAR_004872	AT1G59780.1.TAIR10_R5	2	946
<i>DcRG_040</i>	Chr_2	NL	-	1152442	1155392	DCAR_004880	AT1G10920.1.TAIR10_R5	2	855
<i>DcRG_041</i>	Chr_2	NL	-	2492204	2495294	DCAR_004969	GLYMA.08G247900.1.WM82.A2.V1_R5	5	866
<i>DcRG_042</i>	Chr_2	NL	+	2503242	2506744	DCAR_004970	AT1G10920.1.TAIR10_R9	3	850
<i>DcRG_043</i>	Chr_2	N	+	2541539	2544772	DCAR_004976	AT5G43470.1.TAIR10_R9	2	952
<i>DcRG_044</i>	Chr_2	NL	+	2559352	2562670	DCAR_004978	AT1G58410.1.TAIR10_R8	2	947
<i>DcRG_045</i>	Chr_2	NL	+	2610685	2620387	DCAR_004985	AT1G58807.1.TAIR10_R2	4	976
<i>DcRG_046</i>	Chr_2	NL	+	2625176	2628516	DCAR_004988	AT1G59780.1.TAIR10_R6	3	909
<i>DcRG_047</i>	Chr_2	KIN	+	4062037	4063385	DCAR_005050	PGSC0003DMT400023005.V4.03_R5	5	259
<i>DcRG_048</i>	Chr_2	TNL	+	8605381	8610012	NA	AT5G17970.1.TAIR10_R8	4	790
<i>DcRG_049</i>	Chr_2	RLP	+	8887832	8890630	DCAR_005187	AT3G24982.1.TAIR10_R9	1	933
<i>DcRG_050</i>	Chr_2	TNL	+	9014191	9018101	DCAR_005188	SOLYC09G092410.2.1.ITAG2.4_R6	6	1014

Gene name ¹⁾	Chrom.	Class ²⁾	Genomic coordinates ³⁾			Locus name ⁴⁾	Prediction (GeMoMa) ⁵⁾	Exon no.	AA ⁶⁾
			Strand	Start	Stop				
<i>DcRG_051</i>	Chr_2	TNL	+	9061979	9065369	DCAR_005190	GLYMA.16G085700.1.WM82.A2.V1_R6	4	985
<i>DcRG_052</i>	Chr_2	TNL	-	9847210	9852422	DCAR_005216	POTRI.019G001700.1.V3.0_R5	7	968
<i>DcRG_053</i>	Chr_2	KIN	+	11987487	11989819	DCAR_005345	PGSC0003DMT400023009.V4.03_R2	6	406
<i>DcRG_054</i>	Chr_2	KIN	+	17857017	17859611	DCAR_005745	AT5G39030.1.TAIR10_R6	1	865
<i>DcRG_055</i>	Chr_2	RLP	+	18187570	18190523	DCAR_005778	POTRI.009G112000.1.V3.0_R8	3	859
<i>DcRG_056</i>	Chr_2	NL	+	18270956	18274006	DCAR_005785	POTRI.015G121800.1.V3.0_R2	1	1017
<i>DcRG_057</i>	Chr_2	RLP	+	18742636	18743808	DCAR_005830	POTRI.011G023500.1.V3.0_R2	1	391
<i>DcRG_058</i>	Chr_2	NL	-	18805419	18808028	DCAR_005833	SOLYC04G009110.1.1.ITAG2.4_R6	1	870
<i>DcRG_059</i>	Chr_2	NL	-	20184081	20186635	DCAR_005938	PGSC0003DMT400047507.V4.03_R3	2	842
<i>DcRG_060</i>	Chr_2	KIN	-	20426714	20430070	NA	PGSC0003DMT400023009.V4.03_R1	6	402
<i>DcRG_061</i>	Chr_2	TNL	+	23902080	23906583	NA	AT5G45210.1.TAIR10_R2	4	754
<i>DcRG_062</i>	Chr_2	RLP	-	24115847	24116929	DCAR_006300	GLYMA.13G005800.1.WM82.A2.V1_R4	1	361
<i>DcRG_063</i>	Chr_2	TNL	-	25095215	25099092	NA	POTRI.T127700.1.V3.0_R9	4	934
<i>DcRG_064</i>	Chr_2	TNL	-	25318472	25323739	NA	SOLYC05G006630.2.1.ITAG2.4_R2	5	1301
<i>DcRG_065</i>	Chr_2	KIN	+	25932552	25937554	DCAR_006481	PGSC0003DMT400002807.V4.03_R2	5	402
<i>DcRG_066</i>	Chr_2	TNL	+	26026896	26031279	DCAR_006494	POTRI.019G001700.1.V3.0_R9	5	1143
<i>DcRG_067</i>	Chr_2	TNL	+	26037273	26039281	NA	POTRI.T001400.1.V3.0_R9	3	572
<i>DcRG_068</i>	Chr_2	TNL	-	26320625	26324534	NA	SOLYC05G007850.1.1.ITAG2.4_R6	4	1076
<i>DcRG_069</i>	Chr_2	RLP	-	26326546	26331048	NA	POTRI.T005300.1.V3.0_R3	4	512
<i>DcRG_070</i>	Chr_2	TNL	-	26349468	26354227	NA	POTRI.T001500.1.V3.0_R7	5	1245
<i>DcRG_071</i>	Chr_2	RLK	+	29218504	29220472	DCAR_006860	POTRI.013G158800.1.V3.0_R7	2	605
<i>DcRG_072</i>	Chr_2	KIN	-	32816767	32820124	DCAR_007260	SOLYC03G119240.2.1.ITAG2.4_R4	4	658
<i>DcRG_073</i>	Chr_2	NL	+	35123060	35125756	DCAR_007540	GLYMA.01G010500.1.WM82.A2.V1_R2	1	899
<i>DcRG_074</i>	Chr_2	CNL	+	38170622	38174623	DCAR_007914	GLYMA.03G075200.1.WM82.A2.V1_R0	1	1334
<i>DcRG_075</i>	Chr_2	KIN	-	38765036	38767053	DCAR_007999	GLYMA.08G044300.1.WM82.A2.V1_R8	2	642
<i>DcRG_076</i>	Chr_2	KIN	-	38772867	38774833	DCAR_008001	AT1G66920.1.TAIR10_R5	2	622
<i>DcRG_077</i>	Chr_2	KIN	-	38784684	38786650	DCAR_008003	AT1G66920.1.TAIR10_R4	2	621

Gene name ¹⁾	Chrom.	Class ²⁾	Genomic coordinates ³⁾			Locus name ⁴⁾	Prediction (GeMoMa) ⁵⁾	Exon no.	AA ⁶⁾
			Strand	Start	Stop				
<i>DcRG_078</i>	Chr_2	KIN	-	39842813	39844439	DCAR_008139	PGSC0003DMT400002807.V4.03_R1	5	398
<i>DcRG_079</i>	Chr_2	EDR2*	+	40247417	40257126	DCAR_008189	SOLYC12G096050.1.1.ITAG2.4_R0	22	723
<i>DcRG_080</i>	Chr_2	NL	+	40338741	40341735	DCAR_008200	GLYMA.15G168500.2.WM82.A2.V1_R0	2	938
<i>DcRG_081</i>	Chr_2	NL	-	43131647	43134988	DCAR_008568	GLYMA.15G168500.2.WM82.A2.V1_R4	2	947
<i>DcRG_082</i>	Chr_3	TN	-	431859	433745	NA	AT1G72890.1.TAIR10_R4	2	418
<i>DcRG_083</i>	Chr_3	TNL	-	452369	457048	NA	AT2G14080.1.TAIR10_R3	4	1158
<i>DcRG_084</i>	Chr_3	TNL	-	593784	597788	NA	AT1G63870.1.TAIR10_R7	4	1014
<i>DcRG_085</i>	Chr_3	TNL	+	749747	764287	NA	SOLYC05G006630.2.1.ITAG2.4_R6	5	1300
<i>DcRG_086</i>	Chr_3	TNL	+	803446	807170	NA	AT1G63870.1.TAIR10_R5	4	1073
<i>DcRG_087</i>	Chr_3	TN	+	808952	810503	NA	AT1G66090.1.TAIR10_R2	2	486
<i>DcRG_088</i>	Chr_3	TNL	+	1576802	1580958	NA	AT5G17680.1.TAIR10_R9	4	1120
<i>DcRG_089</i>	Chr_3	CNL	-	1713748	1718604	DCAR_008832	POTRI.018G003500.1.V3.0_R3	3	889
<i>DcRG_090</i>	Chr_3	NL	-	1721885	1725366	DCAR_008833	PGSC0003DMT400079343.V4.03_R0	4	597
<i>DcRG_091</i>	Chr_3	RLP	+	2131140	2133680	DCAR_008878	AT2G33060.1.TAIR10_R4	2	815
<i>DcRG_092</i>	Chr_3	CN	+	2687353	2689026	DCAR_008924	PGSC0003DMT400028241.V4.03_R0	1	558
<i>DcRG_093</i>	Chr_3	CNL	-	4465596	4470353	DCAR_009112	POTRI.003G101000.1.V3.0_R1	1	1586
<i>DcRG_094</i>	Chr_3	RLK	-	5580948	5584234	DCAR_009226	POTRI.019G131500.1.V3.0_R2	2	657
<i>DcRG_095</i>	Chr_3	NL	+	7912809	7916087	DCAR_009477	AT1G59620.1.TAIR10_R5	3	951
<i>DcRG_096</i>	Chr_3	NL	-	15204254	15206854	NA	AT3G46730.1.TAIR10_R4	1	867
<i>DcRG_097</i>	Chr_3	NL	+	16594988	16596481	DCAR_010221	PGSC0003DMT400005733.V4.03_R3	1	498
<i>DcRG_098</i>	Chr_3	NL	+	16600738	16602243	DCAR_010221	PGSC0003DMT400005733.V4.03_R1	1	502
<i>DcRG_099</i>	Chr_3	NL	+	16602695	16605097	DCAR_010221	PGSC0003DMT400047507.V4.03_R1	1	801
<i>DcRG_100</i>	Chr_3	KIN	-	22467470	22470490	DCAR_010403	PGSC0003DMT400023009.V4.03_R0	6	384
<i>DcRG_101</i>	Chr_3	CK	+	23628357	23630908	NA	POTRI.004G096900.1.V3.0_R7	2	619
<i>DcRG_102</i>	Chr_3	RLK	-	26522846	26524976	DCAR_010637	POTRI.019G131500.1.V3.0_R3	2	617
<i>DcRG_103</i>	Chr_3	CNL	+	29959620	29962208	DCAR_010879	PGSC0003DMT400043963.V4.03_R6	1	863
<i>DcRG_104</i>	Chr_3	NL	-	30494490	30502892	DCAR_010928	POTRI.007G039300.1.V3.0_R1	5	822

Gene name ¹⁾	Chrom.	Class ²⁾	Genomic coordinates ³⁾			Locus name ⁴⁾	Prediction (GeMoMa) ⁵⁾	Exon no.	AA ⁶⁾
			Strand	Start	Stop				
<i>DcRG_105</i>	Chr_3	KIN	+	30686966	30691050	DCAR_010945	AT1G18390.1.TAIR10_R7	3	702
<i>DcRG_106</i>	Chr_3	KIN	+	33645917	33648382	DCAR_011214	PGSC0003DMT400079581.V4.03_R1	6	542
<i>DcRG_107</i>	Chr_3	KIN	+	35029913	35032685	DCAR_011361	GLYMA.07G094500.5.WM82.A2.V1_R1	2	637
<i>DcRG_108</i>	Chr_3	KIN	-	35034031	35036812	DCAR_011362	SOLYC12G036330.1.1.ITAG2.4_R7	5	629
<i>DcRG_109</i>	Chr_3	KIN	-	35038969	35041349	DCAR_011363	SOLYC03G119240.2.1.ITAG2.4_R3	3	621
<i>DcPG_110</i>	Chr_3	NL	+	35496072	35497775	DCAR_011401	PGSC0003DMT400005733.V4.03_R6	1	568
<i>DcPG_111</i>	Chr_3	KIN	+	39490394	39493207	NA	PGSC0003DMT400023009.V4.03_R4	6	367
<i>DcPG_112</i>	Chr_3	NL	-	41574966	41577572	DCAR_011885	PGSC0003DMT400047504.V4.03_R2	2	826
<i>DcRG_113</i>	Chr_3	RLP	+	43463639	43469780	DCAR_012065	POTRI.001G389100.1.V3.0_R8	3	846
<i>DcRG_114</i>	Chr_3	RLP	-	43613175	43615967	DCAR_012081	POTRI.001G389100.1.V3.0_R7	2	896
<i>DcRG_115</i>	Chr_3	RLP	-	43619007	43621433	NA	AT3G28890.1.TAIR10_R1	1	809
<i>DcRG_116</i>	Chr_3	RLP	-	43622571	43625189	DCAR_012082	POTRI.001G389100.1.V3.0_R6	2	842
<i>DcRG_117</i>	Chr_3	RLP	-	43625963	43628533	DCAR_012082	AT3G11080.1.TAIR10_R3	1	857
<i>DcRG_118</i>	Chr_3	RLP	-	43630268	43632838	DCAR_012083	AT3G11080.1.TAIR10_R4	1	857
<i>DcRG_119</i>	Chr_3	RLP	-	43633910	43636552	DCAR_012084	AT4G13880.1.TAIR10_R9	2	803
<i>DcRG_120</i>	Chr_3	NL	+	45238656	45241851	NA	GLYMA.18G269500.1.WM82.A2.V1_R1	2	949
<i>DcRG_121</i>	Chr_3	NL	+	45250607	45253911	NA	GLYMA.15G168500.2.WM82.A2.V1_R6	2	948
<i>DcRG_122</i>	Chr_3	NL	-	45518004	45521342	NA	AT1G58410.1.TAIR10_R9	2	963
<i>DcRG_123</i>	Chr_3	NL	-	45523712	45527297	NA	AT1G50180.1.TAIR10_R9	4	857
<i>DcRG_124</i>	Chr_3	RLP	-	46618509	46622364	DCAR_012405	POTRI.001G389100.1.V3.0_R4	2	964
<i>DcRG_125</i>	Chr_3	RLP	-	46950016	46951215	DCAR_012446	POTRI.011G023500.1.V3.0_R1	1	400
<i>DcRG_126</i>	Chr_3	RLP	+	48014155	48017131	NA	AT5G25910.1.TAIR10_R3	2	904
<i>DcRG_127</i>	Chr_3	RLP	-	48361703	48366009	NA	AT5G25910.1.TAIR10_R4	3	852
<i>DcRG_128</i>	Chr_3	RLP	-	48383310	48386225	NA	AT5G25910.1.TAIR10_R2	3	859
<i>DcRG_129</i>	Chr_3	RLP	+	48532959	48535924	NA	AT5G25910.1.TAIR10_R1	2	922
<i>DcRG_130</i>	Chr_3	RLP	+	48914598	48922543	DCAR_012595	AT3G24900.1.TAIR10_R7	2	846
<i>DcRG_131</i>	Chr_3	RLP	+	49027920	49034372	NA	AT2G32660.1.TAIR10_R9	2	675

Gene name ¹⁾	Chrom.	Class ²⁾	Genomic coordinates ³⁾			Locus name ⁴⁾	Prediction (GeMoMa) ⁵⁾	Exon no.	AA ⁶⁾
			Strand	Start	Stop				
<i>DcRG_132</i>	Chr_3	RLP	+	49238871	49241530	DCAR_012608	AT3G23120.1.TAIR10_R4	2	855
<i>DcRG_133</i>	Chr_3	RLP	+	49320867	49324434	DCAR_012610	AT5G40170.1.TAIR10_R7	3	860
<i>DcRG_134</i>	Chr_4	KIN	-	9400357	9401528	DCAR_015659	PGSC0003DMT400023008.V4.03_R8	4	246
<i>DcRG_135</i>	Chr_4	NL	-	10196825	10199416	DCAR_015581	AT3G46710.1.TAIR10_R3	1	864
<i>DcRG_136</i>	Chr_4	RLP	-	10912755	10915271	DCAR_015528	POTRI.011G104900.1.V3.0_R9	1	839
<i>DcRG_137</i>	Chr_4	CNL	-	10979036	10983290	DCAR_015522	POTRI.007G039300.1.V3.0_R2	5	840
<i>DcRG_138</i>	Chr_4	TNL	+	11162302	11167515	NA	SOLYC05G006630.2.1.ITAG2.4_R8	5	1098
<i>DcRG_139</i>	Chr_4	TNL	+	11345889	11350028	NA	AT5G17680.1.TAIR10_R5	4	1109
<i>DcRG_140</i>	Chr_4	KIN	+	13387215	13389872	DCAR_015328	AT5G39030.1.TAIR10_R3	1	886
<i>DcRG_141</i>	Chr_4	KIN	+	15162639	15172670	DCAR_015152	SOLYC12G036330.1.1.ITAG2.4_R1	4	659
<i>DcRG_142</i>	Chr_4	KIN	-	15178142	15186369	DCAR_015151	GLYMA.07G094100.1.WM82.A2.V1_R3	3	756
<i>DcRG_143</i>	Chr_4	KIN	-	15188694	15191151	DCAR_015150	GLYMA.05G237100.1.WM82.A2.V1_R0	3	670
<i>DcRG_144</i>	Chr_4	KIN	-	15210137	15213261	DCAR_015147	GLYMA.05G237100.1.WM82.A2.V1_R4	3	582
<i>DcRG_145</i>	Chr_4	NL	-	15444997	15447666	DCAR_015124	POTRI.003G149800.1.V3.0_R0	1	890
<i>DcRG_146</i>	Chr_4	KIN	+	16125082	16127526	DCAR_015038	PGSC0003DMT400079581.V4.03_R2	7	567
<i>DcRG_147</i>	Chr_4	KIN	-	16820179	16821932	DCAR_014957	PGSC0003DMT400002806.V4.03_R5	4	351
<i>DcRG_148</i>	Chr_4	KIN	+	17397624	17398755	DCAR_014899	PGSC0003DMT400023008.V4.03_R7	4	251
<i>DcRG_149</i>	Chr_4	RLK	-	17794504	17797891	DCAR_014853	POTRI.013G158800.1.V3.0_R4	2	633
<i>DcRG_150</i>	Chr_4	KIN	-	18649658	18655105	NA	SOLYC03G119240.2.1.ITAG2.4_R0	4	673
<i>DcRG_151</i>	Chr_4	NL	-	18705101	18707740	NA	SOLYC04G009260.1.1.ITAG2.4_R0	1	880
<i>DcRG_152</i>	Chr_4	NL	+	19038730	19041485	DCAR_014749	SOLYC04G009150.1.1.ITAG2.4_R2	2	861
<i>DcRG_153</i>	Chr_4	CNL	-	19280907	19283498	DCAR_014736	SOLYC04G009660.2.1.ITAG2.4_R0	1	864
<i>DcRG_154</i>	Chr_4	KIN	-	23072876	23075938	DCAR_014295	AT5G38240.1.TAIR10_R6	2	664
<i>DcRG_155</i>	Chr_4	KIN	-	23088254	23091497	DCAR_014293	AT5G38240.1.TAIR10_R3	2	637
<i>DcRG_156</i>	Chr_4	RLK	-	24239998	24243673	DCAR_014170	POTRI.013G158800.1.V3.0_R8	2	625
<i>DcRG_157</i>	Chr_4	KIN	-	25285981	25290620	NA	PGSC0003DMT400023009.V4.03_R9	6	402
<i>DcRG_158</i>	Chr_4	CNL	+	29504495	29508370	DCAR_013527	POTRI.003G101000.1.V3.0_R0	1	1292

Gene name ¹⁾	Chrom.	Class ²⁾	Genomic coordinates ³⁾			Locus name ⁴⁾	Prediction (GeMoMa) ⁵⁾	Exon no.	AA ⁶⁾
			Strand	Start	Stop				
<i>DcRG_159</i>	Chr_4	RLP	-	32742140	32744629	DCAR_013113	AT1G74190.1.TAIR10_R9	1	830
<i>DcRG_160</i>	Chr_4	RLP	+	34485335	34488235	DCAR_012890	POTRI.016G126900.1.V3.0_R2	1	967
<i>DcRG_161</i>	Chr_4	RLP	+	34492526	34495417	DCAR_012889	POTRI.016G126900.1.V3.0_R4	1	964
<i>DcRG_162</i>	Chr_4	RLP	+	34499027	34501933	DCAR_012888	POTRI.016G126900.1.V3.0_R0	1	969
<i>DcRG_163</i>	Chr_4	RLP	-	34985004	34987841	DCAR_012836	SOLYC07G008590.1.1.ITAG2.4_R6	1	946
<i>DcRG_164</i>	Chr_5	CNL	+	381436	384306	DCAR_016111	POTRI.015G121800.1.V3.0_R0	1	957
<i>DcRG_165</i>	Chr_5	RLP	-	4035831	4037048	DCAR_016491	GLYMA.20G127300.1.WM82.A2.V1_R2	1	406
<i>DcRG_166</i>	Chr_5	NL	-	9192370	9195015	NA	AT3G46730.1.TAIR10_R0	1	882
<i>DcRG_167</i>	Chr_5	CNL	+	10929855	10933017	DCAR_017046	GLYMA.15G168500.2.WM82.A2.V1_R1	2	921
<i>DcRG_168</i>	Chr_5	CNL	-	12380413	12383112	DCAR_017175	POTRI.006G147100.1.V3.0_R0	3	866
<i>DcRG_169</i>	Chr_5	CTNL	+	13287323	13292347	DCAR_017232	GLYMA.03G088100.2.WM82.A2.V1_R5	8	884
<i>DcRG_170</i>	Chr_5	KIN	+	22006152	22007384	DCAR_017641	PGSC0003DMT400067987.V4.03_R5	1	411
<i>DcRG_171</i>	Chr_5	KIN	+	28134725	28137265	DCAR_018279	AT5G39030.1.TAIR10_R7	1	847
<i>DcRG_172</i>	Chr_5	KIN	+	31110250	31113746	DCAR_018558	PGSC0003DMT400002807.V4.03_R8	5	404
<i>DcRG_173</i>	Chr_5	RLK	-	31240587	31243000	DCAR_018578	POTRI.013G158800.1.V3.0_R0	2	641
<i>DcRG_174</i>	Chr_5	KIN	-	31851624	31854293	DCAR_018647	AT5G39030.1.TAIR10_R2	1	890
<i>DcRG_175</i>	Chr_5	RLP	-	32105374	32108775	DCAR_018680	POTRI.009G112000.1.V3.0_R6	1	1134
<i>DcRG_176</i>	Chr_5	RLP	-	32111721	32114933	DCAR_018681	AT1G47890.1.TAIR10_R9	1	1071
<i>DcRG_177</i>	Chr_5	NL	+	33584182	33586779	DCAR_018824	SOLYC04G009660.2.1.ITAG2.4_R3	2	846
<i>DcRG_178</i>	Chr_5	KIN	-	34451354	34452550	DCAR_018917	PGSC0003DMT400067987.V4.03_R1	1	399
<i>DcRG_179</i>	Chr_5	KIN	+	39512287	39514695	DCAR_019492	POTRI.012G054700.1.V3.0_R9	2	396
<i>DcRG_180</i>	Chr_5	KIN	-	40781784	40784888	DCAR_019649	GLYMA.07G094500.5.WM82.A2.V1_R4	2	601
<i>DcRG_181</i>	Chr_5	KIN	+	40785361	40787483	DCAR_019650	AT1G66910.1.TAIR10_R8	2	636
<i>DcRG_182</i>	Chr_5	NL	-	40921822	40924557	DCAR_019675	POTRI.002G216800.1.V3.0_R0	1	912
<i>DcRG_183</i>	Chr_5	RLP	+	41036760	41039498	DCAR_019688	POTRI.016G126900.1.V3.0_R6	1	913
<i>DcRG_184</i>	Chr_5	RLP	+	41045834	41048543	DCAR_019690	POTRI.016G126900.1.V3.0_R7	2	893
<i>DcRG_185</i>	Chr_5	RLP	+	41072410	41075146	DCAR_019691	GLYMA.09G145600.1.WM82.A2.V1_R5	2	871

Gene name ¹⁾	Chrom.	Class ²⁾	Genomic coordinates ³⁾			Locus name ⁴⁾	Prediction (GeMoMa) ⁵⁾	Exon no.	AA ⁶⁾
			Strand	Start	Stop				
<i>DcRG_186</i>	Chr_5	RLP	+	41079818	41082706	DCAR_019692	GLYMA.09G145600.1.WM82.A2.V1_R3	1	963
<i>DcRG_187</i>	Chr_5	RLP	+	41085142	41087892	DCAR_019693	POTRI.016G126900.1.V3.0_R5	1	917
<i>DcRG_188</i>	Chr_5	RLP	+	41104317	41112751	DCAR_019698	AT4G13810.1.TAIR10_R7	2	860
<i>DcRG_189</i>	Chr_5	RLP	-	41178696	41181500	DCAR_019710	POTRI.016G126900.1.V3.0_R1	1	935
<i>DcRG_190</i>	Chr_6	TNL	-	1672431	1677920	DCAR_023109	AT4G12010.1.TAIR10_R4	5	1202
<i>DcRG_191</i>	Chr_6	CNL	-	10198845	10201457	DCAR_022573	SOLYC04G009150.1.1.ITAG2.4_R4	1	871
<i>DcRG_192</i>	Chr_6	CTNL	-	14082355	14092519	DCAR_022321	POTRI.019G001700.1.V3.0_R6	5	1109
<i>DcRG_193</i>	Chr_6	RLP	+	14714744	14722626	DCAR_022265	AT2G34930.1.TAIR10_R5	2	1021
<i>DcRG_194</i>	Chr_6	NL	+	14785298	14787241	DCAR_022256	SOLYC07G039390.1.1.ITAG2.4_R4	4	393
<i>DcRG_195</i>	Chr_6	RLP	+	17725227	17726369	DCAR_021995	GLYMA.20G127300.1.WM82.A2.V1_R0	1	381
<i>DcRG_196</i>	Chr_6	KIN	+	18417261	18419669	NA	PGSC0003DMT400023006.V4.03_R9	4	234
<i>DcRG_197</i>	Chr_6	KIN	+	22339780	22342317	DCAR_021517	AT5G39030.1.TAIR10_R9	1	846
<i>DcRG_198</i>	Chr_6	RLP	-	22565102	22567021	DCAR_021489	AT5G23400.1.TAIR10_R3	1	640
<i>DcRG_199</i>	Chr_6	RLP	-	22574733	22575995	DCAR_021488	GLYMA.20G127300.1.WM82.A2.V1_R1	1	421
<i>DcRG_200</i>	Chr_6	NL	+	22963077	22967507	DCAR_021434	POTRI.007G039300.1.V3.0_R3	6	809
<i>DcRG_201</i>	Chr_6	NL	+	22968646	22973544	DCAR_021433	POTRI.007G039300.1.V3.0_R0	5	827
<i>DcRG_202</i>	Chr_6	RLP	-	25879400	25882168	DCAR_021097	POTRI.T092000.1.V3.0_R9	2	810
<i>DcRG_203</i>	Chr_6	RLP	-	25886193	25889105	DCAR_021096	POTRI.016G126900.1.V3.0_R3	1	971
<i>DcRG_204</i>	Chr_6	NL	-	27171479	27174536	DCAR_020939	GLYMA.14G079500.1.WM82.A2.V1_R0	5	832
<i>DcRG_205</i>	Chr_6	N	-	29899947	29900948	DCAR_020607	AT4G19060.1.TAIR10_R0	1	334
<i>DcRG_206</i>	Chr_6	KIN	-	32480969	32483315	NA	AT1G66910.1.TAIR10_R9	2	663
<i>DcRG_207</i>	Chr_6	RLK	-	32614297	32617063	DCAR_020297	POTRI.013G158800.1.V3.0_R3	2	627
<i>DcRG_208</i>	Chr_6	RLP	+	32666721	32669779	DCAR_020292	SOLYC07G008600.1.1.ITAG2.4_R0	2	989
<i>DcRG_209</i>	Chr_7	NL	-	2743772	2747464	DCAR_023561	GLYMA.01G171000.1.WM82.A2.V1_R1	2	870
<i>DcRG_210</i>	Chr_7	KIN	+	3388041	3392288	DCAR_023631	GLYMA.09G116300.2.WM82.A2.V1_R1	3	629
<i>DcRG_212</i>	Chr_7	KIN	-	9880464	9882796	DCAR_024142	PGSC0003DMT400079581.V4.03_R0	6	431
<i>DcRG_213</i>	Chr_7	NL	-	10329028	10331628	DCAR_024175	SOLYC04G009120.1.1.ITAG2.4_R2	4	671

Gene name ¹⁾	Chrom.	Class ²⁾	Genomic coordinates ³⁾			Locus name ⁴⁾	Prediction (GeMoMa) ⁵⁾	Exon no.	AA ⁶⁾
			Strand	Start	Stop				
<i>DcRG_215</i>	Chr_7	NL	+	14970860	14974237	DCAR_024486	AT3G14470.1.TAIR10_R4	1	1126
<i>DcRG_216</i>	Chr_7	KIN	-	16165740	16171443	DCAR_024551	PGSC0003DMT400002807.V4.03_R4	5	500
<i>DcRG_217</i>	Chr_7	KIN	+	16633579	16636386	DCAR_024598	SOLYC02G086210.2.1.ITAG2.4_R3	3	600
<i>DcRG_218</i>	Chr_7	CK	-	17576626	17584590	DCAR_024644	PGSC0003DMT400002807.V4.03_R3	5	444
<i>DcRG_219</i>	Chr_7	KIN	-	20141986	20142993	DCAR_024858	GLYMA.07G095700.1.WM82.A2.V1_R6	1	336
<i>DcRG_220</i>	Chr_7	KIN	+	20162811	20166302	DCAR_024862	GLYMA.07G094500.5.WM82.A2.V1_R5	2	614
<i>DcRG_221</i>	Chr_7	KIN	-	20196041	20199899	NA	GLYMA.07G095000.1.WM82.A2.V1_R7	2	602
<i>DcRG_222</i>	Chr_7	EDR2*	-	21158149	21170460	DCAR_024947	POTRI.003G100600.1.V3.0_R0	22	725
<i>DcRG_223</i>	Chr_7	CNL	+	23266220	23269705	DCAR_031533	AT3G14470.1.TAIR10_R6	1	1162
<i>DcRG_224</i>	Chr_7	N	+	23906942	23907955	DCAR_025118	PGSC0003DMT400050981.V4.03_R0	1	338
<i>DcRG_225</i>	Chr_7	NL	+	23910837	23914322	DCAR_025119	GLYMA.03G075200.1.WM82.A2.V1_R2	1	1162
<i>DcRG_226</i>	Chr_7	NL	+	24033063	24036572	NA	POTRI.018G003500.1.V3.0_R6	3	926
<i>DcRG_227</i>	Chr_7	NL	+	24063911	24067470	DCAR_025136	POTRI.001G029300.1.V3.0_R9	2	1165
<i>DcRG_228</i>	Chr_7	NL	+	24071521	24075364	DCAR_025137	POTRI.015G121800.1.V3.0_R1	3	926
<i>DcRG_229</i>	Chr_7	CNL	-	24182183	24185614	DCAR_025154	PGSC0003DMT400043963.V4.03_R2	2	1102
<i>DcRG_230</i>	Chr_7	CNL	-	24204879	24208310	DCAR_025158	POTRI.001G029300.1.V3.0_R6	1	1144
<i>DcRG_231</i>	Chr_7	CNL	-	24245002	24248466	DCAR_025161	POTRI.001G029300.1.V3.0_R3	1	1155
<i>DcRG_232</i>	Chr_7	CNL	-	24249990	24253496	DCAR_025162	PGSC0003DMT400043963.V4.03_R3	2	1121
<i>DcRG_233</i>	Chr_7	RLP	+	24458287	24466889	NA	SOLYC12G005620.1.1.ITAG2.4_R1	2	614
<i>DcRG_234</i>	Chr_7	RLP	+	24557336	24560593	NA	POTRI.015G024800.1.V3.0_R1	2	1065
<i>DcRG_235</i>	Chr_7	RLP	-	24673090	24676308	NA	POTRI.015G028600.1.V3.0_R7	3	1005
<i>DcRG_236</i>	Chr_7	RLP	-	24714263	24717556	NA	POTRI.015G024600.1.V3.0_R1	1	1098
<i>DcRG_237</i>	Chr_7	CNL	-	26125618	26129129	DCAR_025331	POTRI.018G138500.1.V3.0_R7	3	968
<i>DcRG_238</i>	Chr_7	CNL	-	26263229	26266696	DCAR_025337	POTRI.018G138500.1.V3.0_R4	3	948
<i>DcRG_239</i>	Chr_7	KIN	-	26793064	26795577	DCAR_025396	AT5G39020.1.TAIR10_R5	1	838
<i>DcRG_240</i>	Chr_7	KIN	-	26800147	26802678	DCAR_025397	AT5G39030.1.TAIR10_R5	1	844
<i>DcRG_241</i>	Chr_7	KIN	-	26997007	26999559	DCAR_025418	AT5G39030.1.TAIR10_R1	1	851

Gene name ¹⁾	Chrom.	Class ²⁾	Genomic coordinates ³⁾			Locus name ⁴⁾	Prediction (GeMoMa) ⁵⁾	Exon no.	AA ⁶⁾
			Strand	Start	Stop				
<i>DcRG_242</i>	Chr_7	KIN	-	27009412	27011979	DCAR_025419	AT5G39030.1.TAIR10_R0	1	856
<i>DcRG_243</i>	Chr_7	KIN	+	27264645	27267206	DCAR_025444	AT5G39030.1.TAIR10_R4	1	854
<i>DcRG_244</i>	Chr_7	KIN	-	29559054	29562867	DCAR_025671	GLYMA.09G116300.2.WM82.A2.V1_R7	3	694
<i>DcRG_245</i>	Chr_7	NL	+	29874881	29877481	DCAR_025716	SOLYC04G009660.2.1.ITAG2.4_R6	2	856
<i>DcRG_246</i>	Chr_7	NL	+	30082257	30083785	NA	POTRI.T001400.1.V3.0_R4	4	406
<i>DcRG_247</i>	Chr_7	KIN	+	31600066	31605352	DCAR_025904	SOLYC02G086270.2.1.ITAG2.4_R0	4	948
<i>DcRG_248</i>	Chr_8	RLK	+	52996	55126	DCAR_028931	POTRI.019G131500.1.V3.0_R1	2	642
<i>DcRG_249</i>	Chr_8	KIN	+	1005839	1006858	DCAR_028838	PGSC0003DMT400067987.V4.03_R0	1	340
<i>DcRG_250</i>	Chr_8	RLP	+	1491967	1495891	NA	POTRI.015G025800.1.V3.0_R6	3	1047
<i>DcRG_251</i>	Chr_8	RLP	+	1497877	1500759	NA	POTRI.015G024600.1.V3.0_R5	1	961
<i>DcRG_252</i>	Chr_8	RLP	+	1526473	1530802	NA	POTRI.015G025800.1.V3.0_R8	4	845
<i>DcRG_253</i>	Chr_8	CK	-	1988272	1990892	DCAR_028743	PGSC0003DMT400023009.V4.03_R3	6	373
<i>DcRG_254</i>	Chr_8	NL	+	2409633	2411897	DCAR_028713	PGSC0003DMT400047507.V4.03_R4	1	755
<i>DcRG_255</i>	Chr_8	RLP	+	3277998	3280631	DCAR_028661	AT5G25910.1.TAIR10_R0	2	846
<i>DcRG_256</i>	Chr_8	RLP	-	15081403	15084282	DCAR_028105	POTRI.009G112000.1.V3.0_R2	1	960
<i>DcRG_257</i>	Chr_8	RLP	+	20277198	20279895	NA	AT3G23110.1.TAIR10_R9	2	822
<i>DcRG_258</i>	Chr_8	KIN	-	21527561	21534304	DCAR_027586	GLYMA.07G094500.2.WM82.A2.V1_R9	4	631
<i>DcRG_259</i>	Chr_8	KIN	+	21534833	21537254	DCAR_027585	GLYMA.07G094100.1.WM82.A2.V1_R9	2	655
<i>DcRG_260</i>	Chr_8	KIN	-	21586442	21590197	DCAR_027574	GLYMA.09G182200.1.WM82.A2.V1_R5	4	538
<i>DcRG_261</i>	Chr_8	KIN	+	21594123	21596404	DCAR_027573	GLYMA.07G094500.5.WM82.A2.V1_R6	2	616
<i>DcRG_262</i>	Chr_8	KIN	-	21620875	21621960	DCAR_027568	GLYMA.07G095700.1.WM82.A2.V1_R9	1	362
<i>DcRG_263</i>	Chr_8	TN	+	22614613	22616761	NA	POTRI.T112700.1.V3.0_R4	2	544
<i>DcRG_264</i>	Chr_8	KIN	+	23518222	23521194	DCAR_027395	PGSC0003DMT400023009.V4.03_R5	6	421
<i>DcRG_265</i>	Chr_8	TNL	-	25975380	25979796	DCAR_027115	SOLYC09G092410.2.1.ITAG2.4_R9	5	1114
<i>DcRG_266</i>	Chr_8	KIN	-	26096075	26101852	DCAR_027102	PGSC0003DMT400002807.V4.03_R5	5	492
<i>DcRG_267</i>	Chr_8	RLP	-	28006751	28008529	DCAR_026875	AT5G23400.1.TAIR10_R1	1	593
<i>DcRG_268</i>	Chr_8	KIN	+	29153059	29155931	DCAR_026747	GLYMA.09G116300.2.WM82.A2.V1_R4	2	614

Gene name ¹⁾	Chrom.	Class ²⁾	Genomic coordinates ³⁾			Locus name ⁴⁾	Prediction (GeMoMa) ⁵⁾	Exon no.	AA ⁶⁾
			Strand	Start	Stop				
<i>DcRG_269</i>	Chr_8	NL	-	29981804	29985804	DCAR_026651	SOLYC04G009690.1.1.ITAG2.4_R7	2	859
<i>DcRG_270</i>	Chr_8	NL	-	29994173	29996725	DCAR_026650	POTRI.003G149800.1.V3.0_R8	1	851
<i>DcRG_271</i>	Chr_8	NL	-	30007354	30009897	DCAR_026649	POTRI.003G149800.1.V3.0_R7	1	848
<i>DcRG_272</i>	Chr_8	NL	-	30022204	30024783	DCAR_026647	SOLYC04G009250.1.1.ITAG2.4_R9	1	860
<i>DcRG_273</i>	Chr_8	NL	-	30030611	30033437	DCAR_026646	GLYMA.01G010500.1.WM82.A2.V1_R8	2	861
<i>DcRG_274</i>	Chr_8	NL	-	30041115	30043715	DCAR_026645	SOLYC04G009660.2.1.ITAG2.4_R9	2	850
<i>DcRG_275</i>	Chr_8	TN	-	30194345	30196181	NA	POTRI.T112700.1.V3.0_R1	2	528
<i>DcRG_276</i>	Chr_8	TNL	-	30196984	30202575	DCAR_026632	POTRI.005G206400.1.V3.0_R7	8	1102
<i>DcRG_277</i>	Chr_8	TNL	-	30207704	30213355	DCAR_026631	POTRI.005G206400.1.V3.0_R1	6	1173
<i>DcRG_278</i>	Chr_8	TNL	-	30218198	30222649	DCAR_026630	GLYMA.03G077400.1.WM82.A2.V1_R0	5	1161
<i>DcRG_279</i>	Chr_8	TNL	-	30227120	30233089	DCAR_026630	POTRI.005G206400.1.V3.0_R0	6	1123
<i>DcRG_280</i>	Chr_8	RLP	+	31346764	31350006	NA	POTRI.015G024600.1.V3.0_R2	1	1081
<i>DcRG_281</i>	Chr_9	KIN	-	4767644	4769636	DCAR_029326	AT1G66920.1.TAIR10_R6	2	626
<i>DcRG_282</i>	Chr_9	KIN	+	4769885	4772336	DCAR_029327	GLYMA.07G094500.5.WM82.A2.V1_R9	2	653
<i>DcRG_283</i>	Chr_9	TNL	+	5354137	5358211	DCAR_029356	SOLYC05G007850.1.1.ITAG2.4_R3	4	1130
<i>DcRG_284</i>	Chr_9	TNL	-	5406512	5411097	DCAR_029364	POTRI.005G206400.1.V3.0_R3	6	1135
<i>DcRG_285</i>	Chr_9	RLP	-	10975964	10979104	DCAR_029693	POTRI.015G024600.1.V3.0_R6	1	1047
<i>DcRG_286</i>	Chr_9	TNL	-	15150058	15155738	DCAR_029849	POTRI.T001500.1.V3.0_R8	5	994
<i>DcRG_287</i>	Chr_9	TNL	-	15288383	15292500	DCAR_029841	POTRI.005G206400.1.V3.0_R2	6	1145
<i>DcRG_288</i>	Chr_9	TNL	-	15293469	15297400	DCAR_029841	GLYMA.16G087100.1.WM82.A2.V1_R1	5	1115
<i>DcRG_289</i>	Chr_9	NL	+	17022468	17025047	DCAR_029804	SOLYC04G009260.1.1.ITAG2.4_R9	1	860
<i>DcRG_290</i>	Chr_9	KIN	-	18411219	18413262	NA	GLYMA.08G044300.1.WM82.A2.V1_R6	2	631
<i>DcRG_291</i>	Chr_9	KIN	+	18413446	18415951	DCAR_029973	GLYMA.05G237100.1.WM82.A2.V1_R7	2	670
<i>DcRG_292</i>	Chr_9	KIN	+	18417593	18420040	NA	GLYMA.20G119600.1.WM82.A2.V1_R7	3	575
<i>DcRG_294</i>	Chr_9	RLP	+	22378825	22381671	DCAR_030240	POTRI.015G024600.1.V3.0_R9	1	949
<i>DcRG_295</i>	Chr_9	RLP	+	22399093	22403137	DCAR_030241	POTRI.015G025100.1.V3.0_R6	3	987
<i>DcRG_296</i>	Chr_9	RLP	-	22765296	22768211	DCAR_030271	SOLYC07G008600.1.1.ITAG2.4_R9	1	972

Gene name ¹⁾	Chrom.	Class ²⁾	Genomic coordinates ³⁾			Locus name ⁴⁾	Prediction (GeMoMa) ⁵⁾	Exon no.	AA ⁶⁾
			Strand	Start	Stop				
<i>DcRG_297</i>	Chr_9	RLP	-	22806645	22809486	DCAR_030276	POTRI.015G028600.1.V3.0_R8	3	836
<i>DcRG_298</i>	Chr_9	CNL	-	22872395	22875076	DCAR_030282	SOLYC04G009660.2.1.ITAG2.4_R8	1	894
<i>DcRG_299</i>	Chr_9	RLP	+	23083373	23086000	DCAR_030294	AT2G34930.1.TAIR10_R9	2	862
<i>DcRG_300</i>	Chr_9	RLP	+	23113481	23116288	DCAR_030298	POTRI.015G024600.1.V3.0_R7	1	936
<i>DcRG_301</i>	Chr_9	TN	-	23678226	23679887	NA	AT1G66090.1.TAIR10_R0	2	483
<i>DcRG_302</i>	Chr_9	RLP	+	23914898	23916658	DCAR_030350	AT5G23400.1.TAIR10_R2	1	587
<i>DcRG_303</i>	Chr_9	TN	+	24270402	24275569	NA	AT5G48780.1.TAIR10_R1	3	641
<i>DcRG_304</i>	Chr_9	TN	-	24423185	24426677	NA	AT4G04110.1.TAIR10_R4	4	252
<i>DcRG_305</i>	Chr_9	TN	-	24443468	24445550	NA	AT1G72890.2.TAIR10_R0	5	476
<i>DcRG_306</i>	Chr_9	TN	+	24990017	24991343	NA	AT1G72890.1.TAIR10_R2	2	383
<i>DcRG_307</i>	Chr_9	TNL	+	25033379	25039407	DCAR_030397	AT5G41750.1.TAIR10_R9	4	1081
<i>DcRG_308</i>	Chr_9	RLP	+	25147901	25149565	DCAR_030412	SOLYC12G005620.1.1.ITAG2.4_R7	1	555
<i>DcRG_309</i>	Chr_9	RLP	+	25187555	25189219	NA	SOLYC12G005620.1.1.ITAG2.4_R8	1	555
<i>DcRG_310</i>	Chr_9	TNL	-	25266572	25272274	NA	AT5G45210.1.TAIR10_R1	4	704
<i>DcRG_311</i>	Chr_9	KIN	-	27687655	27688860	DCAR_030583	PGSC0003DMT400067987.V4.03_R4	1	402
<i>DcRG_312</i>	Chr_9	TNL	-	28722716	28727938	DCAR_030671	GLYMA.08G293600.10.WM82.A2.V1_R1	6	1284
<i>DcRG_313</i>	Chr_9	TNL	-	28730851	28739956	DCAR_030672	SOLYC05G006630.2.1.ITAG2.4_R4	6	1198
<i>DcRG_315</i>	Chr_9	KIN	+	30146336	30149634	DCAR_030820	SOLYC05G010530.2.1.ITAG2.4_R9	4	370
<i>DcRG_316</i>	Chr_9	RLP	-	31851146	31853776	DCAR_031009	SOBIC.009G012900.1.V3.1_R9	1	877
<i>DcRG_317</i>	Chr_9	RLP	+	31865894	31868533	DCAR_031012	AT2G34930.1.TAIR10_R7	1	880
<i>DcRG_318</i>	Chr_9	KIN	+	32206922	32209658	DCAR_031037	PGSC0003DMT400023009.V4.03_R6	5	420
<i>DcRG_319</i>	Chr_9	KIN	-	32350016	32352409	DCAR_031050	PGSC0003DMT400023009.V4.03_R7	5	426
<i>DcRG_320</i>	Chr_9	KIN	+	32455717	32457660	DCAR_031067	POTRI.010G035300.1.V3.0_R0	1	648
<i>DcRG_321</i>	Chr_9	TNL	+	32519349	32525701	DCAR_031076	SOLYC05G006620.2.1.ITAG2.4_R0	5	1428
<i>DcRG_322</i>	Chr_9	TNL	+	32890459	32894388	NA	AT1G72840.2.TAIR10_R5	4	1153
<i>DcRG_323</i>	Contig_S190.1	KIN	+	104964	106058	DCAR_031991	PGSC0003DMT400067987.V4.03_R9	1	365
<i>DcRG_324</i>	Contig_S72.1	NL	-	12023	13532	DCAR_032355	POTRI.T001400.1.V3.0_R6	4	418

- 1) Gene numbers 211, 214, 293, and 314 are not existing
- 2) Class assignment according to DRAGO 3 / PRGdb 4.0 database (Calle-Garcia et al. 2022). Unknown domain combinations are in italics (for details, see Table 2). *assignment after BLAST P
- 3) Genomic coordinates and ⁴⁾ DCAR locus names according to the carrot whole genome sequence assembly vers.2
- 5) AT - *Arabidopsis thaliana*; GLYMA - *Glycine max*, POTRI - *Populus trichocarpa*, SOLYC - *Solanum lycopersicum*; SOBIC - *Sorghum bicolor*, PGSC - *Solanum tuberosum*
- 6) Protein length (AA - Amino Acids)